

This is Author proof version of the paper published in Scientia Horticulturae <https://doi.org/10.1016/j.scienta.2021.110405> and is available under a CC-BY-NC-ND international licence.

The final authenticated version is available online at [10.1016/j.scienta.2021.110405](https://doi.org/10.1016/j.scienta.2021.110405)

Terms of use see Elsevier article sharing policy see open access options: <https://www.elsevier.com/journals/scientia-horticulturae/0304-4238/open-access-options> for archived author accepted manuscripts (AAMs) of subscription articles.

1 **Genetic diversity of common bean (*Phaseolus vulgaris* L.) germplasm from Serbia, as revealed**  
2 **by single sequence repeats (SSR)**

3

4 Aleksandra Savić<sup>a</sup>,

5 email: [aleksandra.savic@ifvcns.ns.ac.rs](mailto:aleksandra.savic@ifvcns.ns.ac.rs)

6 Barbara Pipan<sup>b</sup>,

7 email: [barbara.pipan@kis.si](mailto:barbara.pipan@kis.si)

8 Mirjana Vasić<sup>a</sup>,

9 email: [mirjana.vasic@ifvcns.ns.ac.rs](mailto:mirjana.vasic@ifvcns.ns.ac.rs)

10 Vladimir Meglič<sup>b</sup>

11 email: [vladimir.meglic@kis.si](mailto:vladimir.meglic@kis.si)

12

13 <sup>a</sup> Institute of Field and Vegetable Crops, Maksima Gorkog 30, 21000 Novi Sad, Serbia

14 <sup>b</sup> Agricultural Institute of Slovenia, Hacquetova ulica 17 SI-1000 Ljubljana, Slovenia

15

16 corresponding author: Aleksandra Savić, email: [aleksandra.savic@ifvcns.ns.ac.rs](mailto:aleksandra.savic@ifvcns.ns.ac.rs)

17 postal address: Institute of Field and Vegetable Crops, Maksima Gorkog 30, 21000 Novi Sad, Serbia

18

19 **Abstract**

20 Genetic diversity and structure of common bean (*Phaseolus vulgaris*) germplasm from Serbia,  
21 comprising 118 landraces and 18 cultivars, was assessed with the application of 27 Single Sequence  
22 Repeats (SSR) markers. Thirteen accessions from Agricultural Institute of Slovenia were used as  
23 references for gene pool determination. Main parameters of genetic diversity were calculated for each  
24 SSR loci, i.e. number of different and rare alleles, number of effective alleles, Shannon's information  
25 index, observed and expected heterozygosity and polymorphic information content. A total of 445  
26 allelic variants, with 16.5 alleles per locus on average, were detected. Mean gene diversity ( $H_e = 0.79$ )  
27 indicated sufficient reservoir of genetic variation preserved in studied bean germplasm. Landraces  
28 displayed higher variability compared to cultivars (405 in relation to 233 allelic variants). Genetic

29 structure and relatedness of accessions was assessed by model-based method and hierarchical clustering  
30 method in combination with genetic distance calculation. The Bayesian clustering model implemented  
31 in STRUCTURE software, on the primary level (K=2), revealed clear separation of accessions into two  
32 groups, corresponding to gene pool affiliation. Mesoamerican gene pool (M) was represented with  
33 23.5% of accessions, while Andean (A) was larger, composed of 68.4% of studied germplasm. Small  
34 group (8.1%) showed admixed genetic structure between two gene pools. Additional variation in respect  
35 to two recognized gene pools was revealed (K=3), whose basis was acknowledged to be within Andean  
36 gene pool. Further subdivision of accessions (K=8), mainly according to the seed forms, was observed.  
37 Genetic distance analysis associated with Neighbour-joining clustering method revealed grouping  
38 pattern of landraces and cultivars corresponding to the gene pool and their seed phenotypes.  
39 Classification and structuring of the bean accessions according to and beyond the gene pool of origin  
40 should facilitate conservation strategies and breeding of this material. Combining the information of  
41 phenotypic variation obtained in previous research and molecular data revealed in this study will assist  
42 in selection of parental components for breeding, or in the choice of smaller sample in order to further  
43 acknowledge their breeding value. In addition, obtained results of this work should serve as an  
44 additional information on common bean germplasm variation in Western Balkans and beyond, in  
45 Europe.

46

47 **Keywords:** *Phaseolus vulgaris*, SSR, genetic diversity, gene pool

48

## 49 **1. Introduction**

50

51 The common bean (*Phaseolus vulgaris* L.) is one of the most valuable vegetable crops for  
52 human consumption since it is rich in proteins, fibres, vitamins, minerals and many other antioxidant  
53 compounds (Broughton et al., 2003; Maras et al., 2015; Sitohy et al., 2020). Being a diverse crop in  
54 terms of cultivation methods, use, phenotypic diversity and range of environments to which it is  
55 adapted, the common bean is grown worldwide (De Ron et al., 2016). In many European countries,  
56 *Phaseolus vulgaris* is a significant component of a traditional diet and life.

57 *Phaseolus vulgaris* has undergone two independent domestication events in primary centres of  
58 origin: one in Middle America, and one in Andes. As a result, two highly differentiated gene pools were  
59 formed: Mesoamerican and Andean, which are distinguished and recognized according to their  
60 phenotypic, biochemical and genotypic differences (Gepts et al., 1986; Gepts, 1999; Šuštar-Vozlič et  
61 al., 2006; Raggi et al., 2013; Carović-Stanko et al., 2017; Gioia et al., 2019; Savić et al., 2020). These  
62 two gene pools are also characterized by geographic and partial reproductive barriers (Gepts and Bliss,  
63 1985; Gioia et al., 2013). Furthermore, within each gene pool, there is subdivision of this species to  
64 many eco-geographic races and seed forms (market classes). Inter-gene pool and interracial crosses of  
65 genotypes can exhibit negative combining ability and lethality problems, which aggravates the breeding  
66 of common bean (Singh et al., 1991; Kelly et al., 1998; Blair et al., 2007).

67 Gene pool affiliation has usually been determined based on variation of the main storage protein  
68 of the common bean, phaseolin (Gepts and Bliss, 1988; Šuštar-Vozlič et al., 2006; Logozzo et al., 2007;  
69 Carović-Stanko et al., 2017; Savić et al., 2020). Mesoamerican origin of common bean genotypes is  
70 associated with phaseolin types S (*Sanilac*), M (*Middle America*) and B (*Boyaca*), while genotypes with  
71 phaseolin types T (*Tendergreen*), C (*Contender*) and H (*Huevo de Huanchaco*) belong to the Andean  
72 gene pool. However, since Nani et al. (2011) identified three indel spanning markers SHP1-A, SHP1-  
73 B and SHP1-C, newer researches more rely on these marker systems for gene pool identification (Maras  
74 et al., 2015, 2016; Pipan and Meglič, 2019).

75 From primary centres of origin and domestication in America, common bean spread worldwide  
76 (Zeven, 1997; Maras et al., 2015). It is believed that the common bean arrived in Europe on two  
77 occasions; Mesoamerican beans through Spanish and Portuguese exploration of the Americas around  
78 1506; and Andean beans somewhere later, in 1528, during Pizarro's expeditions in Peru (Gioia et al.,  
79 2013). In Europe, common bean landraces and cultivars evolved under diverse environments, cropping  
80 systems and farmers preferences (Zeven, 1997; Carović-Stanko et al., 2017; Pipan and Meglič, 2019).  
81 In addition, outcrossing among Andean and Mesoamerican genotypes facilitated in the development of  
82 high genotypic and phenotypic diversity of the European common bean (Rodino et al., 2006; Gioia et  
83 al., 2013).

84           Angioi et al. (2010) and Gioia et al. (2013) focused their research on hybridization phenomena  
85 of Andean and Mesoamerican germplasm in Europe. The presence of hybrid genotypes in high  
86 proportion within the European germplasm was revealed with chloroplast microsatellites (cpSSRs) and  
87 two unlinked nuclear loci (phaseolin and PvSHP1). Hegay et al. (2013) and Sinkovič et al. (2019)  
88 observed signs of introgression on the phenotypic level. Zhang et al. (2008) believe that identification  
89 of putative hybrids is of special interest since these genotypes are a reflection of growing regions and  
90 are adapted to ecological conditions within that region.

91           In Serbia, people have grown common bean for centuries, establishing it as an important part  
92 of their diet and traditional life. Even though commercial cultivars have been developed and are widely  
93 represented in production, common bean landraces are still maintained and used by farmers and people  
94 in rural and marginal areas (Scarano et al., 2014; Mallor et al., 2018; Savić et al., 2020). In Europe  
95 (including Serbia) development of new cultivars was instituted with the aim to maintain phenotypes  
96 within each seed form (market class), usually by selecting parental components among elite material.  
97 This resulted in narrowing the genetic base of this elite germplasm, compromising long-term genetic  
98 gain (McClellan and Lee, 2007; Gioia et al., 2019). In order to address these problems, breeders sought  
99 to incorporate new variability, most commonly by exploring the existing diversity found among  
100 landraces and unadapted germplasm.

101           Genetic collections of the common bean in Serbia, consisting of seeds of traditional and modern  
102 cultivars and landraces, are maintained within breeding institutes (Vasić et al., 2009). Knowledge of  
103 genetic diversity preserved in these collections is crucial for proper conservation, further research,  
104 selection of parental components and for defining breeding strategies. Landraces are described as  
105 genetically diverse material with traits specific for growing regions. They are traditionally grown in low  
106 input systems, adapted to local agro-climatic conditions and display a high level of phenotypic diversity.  
107 All of this makes landraces interesting material for conservation, research and implementation in  
108 breeding programs (Carović-Stanko et al., 2017; Gioia et al., 2019).

109           Selected set of landraces and commercial cultivars from Serbia have already been characterized  
110 for diversity on a phenotype level, by assessing morphological traits chosen according to international  
111 descriptors. In addition, gene pool of origin was identified based on variation of phaseolin types (for

112 more information see Savić et al., 2020). However, in order to better understand genetic variation and  
113 relationships among landraces and cultivars on a molecular level, complementary study on previous  
114 research was performed. Single sequence repeats (SSR) markers were chosen for the analysis, because  
115 they are abundant and widely distributed in the genome, codominantly inherited, highly polymorphic  
116 and repeatable (Yu et al., 1999; Maras et al., 2015; Pipan and Meglič, 2019). Therefore, the aim of this  
117 study was to: (i) assess the allelic diversity of the common bean germplasm from Serbia and determine  
118 relationships among the accessions, and (ii) investigate the genetic structure and organization of genetic  
119 diversity of the studied germplasm within and beyond gene pools of origin.

120

## 121 **2. Material and Methods**

122

### 123 2.1. Plant material

124

125 A total of 136 accessions from the Serbian common bean genetic collection maintained at the  
126 Institute of Field and Vegetable Crops, Novi Sad (IFVCNS) were analysed in this paper. This included  
127 118 local landraces collected from 53 sites in Serbia (Supplementary Material 1) in timescale of 1970-  
128 2014 and 18 commercially available cultivars from Serbia: Rozalija (C1), Žutotrban (C2), Sremac (C3),  
129 Balkan (C4), Slavonski žutozeleni (C5), Pasuljica P-1 (C6), Biser (C7), Medijana (C8), Oplenac (C9),  
130 Panonski gradištanac (C10), Panonski tetovac (C11), Aster (C12), Pobjoljšani gradištanac (C13), Galeb  
131 (C14), Zlatko (C15), Dvadesetica (C16), Belko (C17) and Maksa (C18), Table 1. Part of the studied  
132 material was collected during field expeditions, while the rest was acquired via seed exchange with  
133 other institutions. Landraces and cultivars were classified according to seed traits in several forms, most  
134 commonly grown in Serbia: *Roseus* (pink seed colour), *Versicolor* (seed coat pattern), *Griseus*  
135 (greenish-yellow seed colour), *Aureus* (yellow and golden-yellow seed coat colour), *Albus* (white seed  
136 colour) and other (red, cream, brown, black seed colour).

137 As references for gene pool determination, accessions of familiar phaseolin type (type T –  
138 PHA131, PHA306, PHA309, PHA318, PHA336; type C – PHA181, PHA222, PHA29, PHA315,  
139 PHA390; type S – PHA245, PHA368, PHA371) from Agricultural Institute of Slovenia (AIS) were

140 included in the study (Supplementary Material 1). Phaseolin types T and C indicate germplasm origin  
141 from Andean, while the Mesoamerican gene pool is determined with type S. For detailed phenotypic  
142 characterization, see Savić et al. (2020).

143

## 144 2.2. SSR analysis

145

146 Plants were grown in greenhouse conditions until the phase of the first true leaves at  
147 Agricultural Institute of Slovenia. DNA was extracted from a total of 60 to 100 mg of bulked fresh plant  
148 tissue (4 plants per accession), using a BioSprint 15 DNA Plant Kit (Qaigen) on KingFisher (Thermo)  
149 isolation robot according to optimized manufacturer's instructions.

150 A set of 27 genome-specific SSR (single sequence repeats; microsatellites) markers distributed  
151 across all linkage groups was used for genetic diversity and genetic structure analysis of selected  
152 material (Supplementary Material 1). For the identification of genotypes gene pool affiliation  
153 (Mesoamerican/Andean) three indel spanning markers (SHP1-A, SHP1-B and SHP1-C) developed by  
154 Nanni *et al.* (2011) were used.

155 The final volume of PCR reaction was 11.5 µL, which included: 8.4 ng genomic DNA, 1 µL  
156 10x PCR buffer (Biotools), 0.2 µL of each 10 mM dNTP (Sigma-Aldrich), 0.5 µL of 50mM MgCl<sub>2</sub>  
157 (Biotools), 0.1 µL of 10 µM forward primer (Sigma-Aldrich), 0.25 µL of 10 µM reverse primer (Sigma-  
158 Aldrich), 0.183 µL of 10 µM 5'-fluorescently labelled primer (6-FAM, NED or HEX; Omega), and  
159 0.5 µL of 5 U Taq DNA polymerase (Biotools). The forward primer of each SSR had an added 18-bp  
160 tail sequence of 5'-TGTAACGACGGCCAGT-3' (M13(-21)).

161 PCR analyses were performed on a thermal cycler (Veriti, ThermoFisher Scientific) under  
162 touch-down conditions: 94°C for 4 min; 15 cycles at 94 °C for 1 min; decreased temperature from 60  
163 (62) °C to 49.5 (51.5) °C at 0.7 °C per cycle for 30 s; 72 °C for 1 min; followed by 23 cycles at 94 °C  
164 for 30 s; 53 °C for 30 s; 72 °C for 1 min; and final extension for 5 min at 72 °C, as described by Pipan  
165 and Meglič (2019). PCR conditions were dependent on each primer pair. Fragment analysis was  
166 performed on a genetic analyser (3130XL; Applied Biosystems). Allele lengths were determined by

167 comparison with an internal size standard (GeneScan-350 ROX; Applied Biosystems) using the  
 168 GeneMapper 4.0 software (Applied Biosystems).

169

170 Table 1. List of accessions from Serbian common bean genetic collection, used for genetic analysis

Accession	Type	Seed form	Accession	Type	Seed form
<b>L1</b>	III	other	<b>L56</b>	I	<i>Versicolor</i>
<b>L2</b>	III	<i>Aureus</i>	<b>L57</b>	I	<i>Griseus</i>
<b>C1</b>	I	<i>Roseus</i>	<b>L58</b>	I	<i>Albus</i>
<b>L3</b>	I	<i>Roseus</i>	<b>L59</b>	II	<i>Albus</i>
<b>L4</b>	I	<i>Roseus</i>	<b>L60</b>	I	<i>Griseus</i>
<b>L5</b>	I	<i>Roseus</i>	<b>C14</b>	I	<i>Albus</i>
<b>L6</b>	I	<i>Roseus</i>	<b>C15</b>	I	<i>Aureus</i>
<b>L7</b>	I	<i>Roseus</i>	<b>C16</b>	I	<i>Albus</i>
<b>L8</b>	I	<i>Roseus</i>	<b>C17</b>	I	<i>Albus</i>
<b>L9</b>	II	<i>Griseus</i>	<b>C18</b>	I	<i>Albus</i>
<b>L10</b>	I	<i>Griseus</i>	<b>L61</b>	I	<i>Griseus</i>
<b>L11</b>	II	<i>Albus</i>	<b>L62</b>	I	<i>Griseus</i>
<b>L12</b>	I	<i>Albus</i>	<b>L63</b>	I	<i>Griseus</i>
<b>C2</b>	I	<i>Versicolor</i>	<b>L64</b>	I	<i>Griseus</i>
<b>L13</b>	I	<i>Versicolor</i>	<b>L65</b>	I	<i>Aureus</i>
<b>L14</b>	I	<i>Versicolor</i>	<b>L66</b>	I	other
<b>L15</b>	I	<i>Versicolor</i>	<b>L67</b>	I	other
<b>L16</b>	I	<i>Versicolor</i>	<b>L68</b>	I	<i>Griseus</i>
<b>L17</b>	I	<i>Versicolor</i>	<b>L69</b>	I	<i>Griseus</i>
<b>L18</b>	I	<i>Versicolor</i>	<b>L70</b>	I	<i>Versicolor</i>
<b>L19</b>	I	<i>Versicolor</i>	<b>L71</b>	I	other
<b>L20</b>	I	<i>Griseus</i>	<b>L72</b>	I	other
<b>L21</b>	I	<i>Griseus</i>	<b>L73</b>	I	<i>Griseus</i>
<b>L22</b>	I	<i>Griseus</i>	<b>L74</b>	I	<i>Griseus</i>



<b>L23</b>	I	<i>Griseus</i>	<b>L75</b>	I	<i>Griseus</i>
<b>L24</b>	I	<i>Griseus</i>	<b>L76</b>	I	<i>Versicolor</i>
<b>L25</b>	I	<i>Griseus</i>	<b>L77</b>	I	<i>Albus</i>
<b>L26</b>	I	<i>Griseus</i>	<b>L78</b>	I	<i>Albus</i>
<b>L27</b>	I	<i>Griseus</i>	<b>L79</b>	III	<i>Albus</i>
<b>L28</b>	I	<i>Aureus</i>	<b>L80</b>	I	<i>Griseus</i>
<b>L29</b>	I	other	<b>L81</b>	I	<i>Griseus</i>
<b>L30</b>	I	other	<b>L82</b>	I	<i>Versicolor</i>
<b>L31</b>	I	<i>Griseus</i>	<b>L83</b>	I	<i>Albus</i>
<b>L32</b>	I	other	<b>L84</b>	I	<i>Griseus</i>
<b>L33</b>	I	other	<b>L85</b>	I	<i>Versicolor</i>
<b>L34</b>	I	<i>Griseus</i>	<b>L86</b>	I	<i>Griseus</i>
<b>L35</b>	I	other	<b>L87</b>	I	other
<b>L36</b>	I	other	<b>L88</b>	I	<i>Albus</i>
<b>L37</b>	I	<i>Griseus</i>	<b>L89</b>	III	<i>Albus</i>
<b>L38</b>	I	<i>Roseus</i>	<b>L90</b>	I	other
<b>L39</b>	I	<i>Roseus</i>	<b>L91</b>	I	<i>Albus</i>
<b>L40</b>	I	other	<b>L92</b>	II	<i>Aureus</i>
<b>L41</b>	I	other	<b>L93</b>	III	<i>Albus</i>
<b>L42</b>	I	<i>Roseus</i>	<b>L94</b>	II	other
<b>L43</b>	II	<i>Aureus</i>	<b>L95</b>	I	<i>Griseus</i>
<b>L44</b>	I	<i>Aureus</i>	<b>L96</b>	I	<i>Versicolor</i>
<b>L45</b>	I	<i>Aureus</i>	<b>L97</b>	I	other
<b>L46</b>	I	<i>Aureus</i>	<b>L98</b>	III	<i>Albus</i>
<b>C3</b>	I	<i>Griseus</i>	<b>L99</b>	I	<i>Versicolor</i>
<b>C4</b>	I	<i>Albus</i>	<b>L100</b>	I	other
<b>C5</b>	I	<i>Griseus</i>	<b>L101</b>	I	<i>Versicolor</i>
<b>C6</b>	II	<i>Albus</i>	<b>L102</b>	I	<i>Griseus</i>
<b>C7</b>	I	<i>Albus</i>	<b>L103</b>	I	<i>Griseus</i>
<b>C8</b>	II	<i>Albus</i>	<b>L104</b>	III	<i>Albus</i>

<b>L47</b>	I	<i>Albus</i>	<b>L105</b>	I	<i>Albus</i>
<b>L48</b>	I	<i>Albus</i>	<b>L106</b>	I	<i>Albus</i>
<b>L49</b>	I	<i>Albus</i>	<b>L107</b>	II	<i>Albus</i>
<b>L50</b>	I	<i>Griseus</i>	<b>L108</b>	I	<i>Versicolor</i>
<b>L51</b>	I	<i>Albus</i>	<b>L109</b>	I	<i>Albus</i>
<b>C9</b>	I	<i>Albus</i>	<b>L110</b>	I	<i>Versicolor</i>
<b>L52</b>	I	<i>Albus</i>	<b>L111</b>	I	<i>Versicolor</i>
<b>L53</b>	I	<i>Albus</i>	<b>L112</b>	I	<i>Albus</i>
<b>C10</b>	I	<i>Albus</i>	<b>L113</b>	I	<i>Albus</i>
<b>C11</b>	I	<i>Albus</i>	<b>L114</b>	I	<i>Griseus</i>
<b>L54</b>	II	<i>Aureus</i>	<b>L115</b>	I	<i>Versicolor</i>
<b>C12</b>	I	<i>Albus</i>	<b>L116</b>	III	<i>Albus</i>
<b>C13</b>	III	<i>Albus</i>	<b>L117</b>	I	<i>Albus</i>
<b>L55</b>	I	<i>Versicolor</i>	<b>L118</b>	I	<i>Albus</i>

---

Type – plant growth habit (I – determinate bush, II – indeterminate bush,

III – indeterminate prostrate or vining)

171

172

### 173 2.3. Data analysis

174

175 For each SSR locus, main parameters of genetic diversity were calculated in GenAlEx 6.1  
 176 (Peakall and Smouse, 2006) and Microsatellite-Toolkit (Park, 2001) software. They included number  
 177 of alleles (Na), number of alleles with frequency  $\geq 5\%$ , number of rare alleles, number of effective  
 178 alleles (Ne), Shannon's information index (I), observed heterozygosity (Ho), expected heterozygosity  
 179 (He) and polymorphic information content (PIC).

180 STRUCTURE 2.3.3 software was employed to establish genetic structure of common bean  
 181 collection. STRUCTURE uses Bayesian clustering approach, applying Markov Chain Monte Carlo  
 182 (MCMC) algorithm, to study difference in accessions distribution among population by placing  
 183 accessions into clusters that share similar variation patterns. Bayesian clustering approach is described  
 184 by the posterior probability that each accession belong to each genetic cluster, while each cluster (K) is  
 185 characterized by a subset of allelic frequencies identified in the data. Beside determining and assigning

186 accessions to genetic clusters, this method is used to identify admixed accessions by investigating  
187 hybridization zones of individuals from different clusters that give genetically recombined offspring  
188 (Pritchard *et al.*, 2009). In the study of common bean collection from Serbia, the most probable number  
189 of clusters (Q value) was determined with ten independent runs for each K (1 to 10) for the admixture  
190 model, with burning period of 50.000 followed by 500.000 Markov Chain Monte Carlo repeats. MCMC  
191 algorithm initiates by randomly assigning accessions to a pre-determined number of clusters. After that,  
192 allelic frequencies are estimated for each cluster and accessions are re-assigned based on those  
193 frequency assessments. According to Evanno delta K statistics (Evanno *et al.*, 2005), implemented in  
194 the software Structure Harvester (Earl and von Holdt, 2011), the real K value was determined based on  
195 the increase in the likelihood ratios between the runs. An accession is assigned to a specific cluster  
196 when the percentage of membership was  $Q \geq 80\%$ , while the accessions with membership coefficient  
197  $Q < 80\%$  are believed to be of admixed origin (putative hybrids).

198 For more detailed analysis of genetic structure and relationships among genotypes, DARwin  
199 software (<https://darwin.cirad.fr/>) was applied to perform cluster analysis based on similarity matrix  
200 and construct dendrogram using Neighbour-joining method (NJ).

201

### 202 **3. Results**

203

204 The whole set of 27 SSR markers chosen for the diversity study of the Serbian common bean  
205 genetic collection proved to be polymorphic, producing more than one allelic variants. A total of 445  
206 alleles were scored for the studied collection, which included cultivars, landraces and reference  
207 accessions. The average allele number per SSR was 16.5, ranging from six alleles for loci BM155,  
208 BM210 and BMd044 to 25 alleles for loci GATS91 and ATA002. The highest number of effective  
209 alleles (9.35) and the most alleles with frequency over  $>5\%$  were found in locus BMd001. Total number  
210 of rare alleles was 303, which accounted for 68.1% of all alleles detected in the studied germplasm  
211 (Table 2).

212

213 Table 2. Parameters of genetic variability of Serbian common bean collection and reference cultivars

<b>Locus</b>	<b>Na</b>	<b>Allele frequency ≥ 5%</b>	<b>Number of rare alleles</b>	<b>Ne</b>	<b>I</b>	<b>Ho</b>	<b>He</b>	<b>PIC</b>
ATA003	19	6	13	6.87	2.26	0.74	0.85	0.84
ATA004	15	4	11	3.82	1.77	0.44	0.74	0.71
ATA005	20	6	14	6.60	2.28	0.93	0.85	0.83
ATA007	17	5	12	3.42	1.74	0.30	0.71	0.68
ATA016	12	5	7	4.41	1.78	0.91	0.77	0.74
GATS91	25	5	20	8.38	2.49	1.00	0.88	0.87
ATA002	25	5	20	5.31	2.09	0.85	0.81	0.79
BM172	22	6	16	8.43	2.45	0.92	0.88	0.87
BMd001	24	9	15	9.35	2.46	0.99	0.89	0.88
ATA020	21	5	16	6.52	2.27	0.65	0.85	0.83
Pv-ag004	21	5	16	5.80	2.14	0.94	0.83	0.81
ATA010	13	5	8	4.25	1.75	0.31	0.77	0.73
BM155	6	4	2	3.53	1.36	0.87	0.72	0.67
BM170	20	6	14	8.07	2.40	0.95	0.88	0.87
BM183	8	5	3	3.26	1.41	0.93	0.69	0.64
BM210	6	2	4	2.14	0.85	0.97	0.53	0.42
BMd044	6	4	2	2.69	1.23	0.45	0.63	0.58
ATA009	22	8	14	8.18	2.45	0.75	0.87	0.87
ATA145	20	3	17	3.68	1.80	0.31	0.73	0.69
GA16	13	6	7	6.12	2.04	0.68	0.84	0.82
ATA006	22	7	15	8.98	2.52	0.79	0.89	0.88
BM157	14	6	8	5.88	2.00	0.60	0.83	0.81
BMd042	18	4	14	6.08	2.18	0.98	0.84	0.82
ATA289	17	4	13	3.87	1.74	0.97	0.74	0.71
PvSHP1-A	13	6	7	5.52	1.93	0.95	0.82	0.79
PvSHP1-B	11	6	5	5.89	1.89	1.00	0.83	0.81
PvSHP1-C	15	5	10	5.45	2.03	0.92	0.82	0.79

average	<b>16.5</b>	<b>5.2</b>	<b>11.2</b>	<b>5.65</b>	<b>1.97</b>	<b>0.78</b>	<b>0.79</b>	<b>0.77</b>
total	<b>445</b>	<b>142</b>	<b>303</b>					

214 Na – number of alleles, Ne – number of effective alleles, I - Shannon’s information index, Ho –observed  
215 heterozygosity, He - expected heterozygosity, PIC - polymorphic information content

216

217 Loci BMd001 and ATA006 generated the highest values of expected heterozygosity (0.89) and  
218 PIC (0.88). The locus ATA006 scored the highest value of Shannon’s information index (2.58).  
219 Observed heterozygosity ranged from 0.30 (locus ATA007) to 1.00 (loci GATS and PvSHP1-B).  
220 Average values of Shannon’s information index (1.97), expected heterozygosity (0.79) and PIC (0.77)  
221 indicated that all SSR markers showed sufficient polymorphism and are suitable for common bean  
222 diversity study.

223 Main genetic diversity parameters were calculated for each common bean form determined  
224 according to seed traits (Table 3). Overall, landraces and cultivars from the *Albus* group showed the  
225 greatest diversity for all parameters, except number of alleles with a frequency higher than 5%. On  
226 contrary, the lowest diversity was found among accessions from the *Rosues* group, with slightly larger  
227 expected heterozygosity (0.76). The highest average number of alleles with a frequency higher than 5%  
228 was observed in the *Aureus* form. Rare alleles were not found among common bean genotypes from  
229 *Roseus* and *Aureus* groups, while for the other groups ranged from 3 (*Versicolor*) to 4.96 (*Albus*). Gene  
230 diversity (He) ranged from 0.67 (*Griseus*) to 0.78 (*Albus*). Observed heterozygosity was the highest  
231 among accessions from other (0.82) and *Aureus* (0.81) groups. Almost twice as many alleles were  
232 scored among landraces (405) compared to cultivars (233). In addition, the percentage of rare alleles  
233 was much larger in landraces (67.6%) in relation to cultivars (29.6%). Average values of all the other  
234 parameters of genetic diversity (number of effective alleles, Shannon’s information index, observed and  
235 expected heterozygosity) were similar between the two groups.

236

237 Table 3. Genetic diversity calculated for 27 SSR loci considering six groups defined according to the  
238 seed form; cultivars and landraces separately

<b>Forms</b>	<b>Total number of accessions</b>	<b>Na</b>	<b>Na average</b>	<b>Allele frequency ≥ 5%</b>	<b>Number of rare alleles</b>	<b>Ne</b>	<b>I</b>	<b>Ho</b>	<b>He</b>
<i>Roseus</i>	10	144	5.33	5.33	0	3.39	1.33	0.66	0.76
<i>Versicolor</i>	21	205	7.59	4.59	3	4.22	1.57	0.72	0.76
<i>Grisues</i>	35	231	8.55	4.52	4.03	3.88	1.49	0.79	0.67
<i>Aureus</i>	10	175	6.48	6.48	0	4.33	1.56	0.81	0.73
<i>Albus</i>	42	294	10.89	5.93	4.96	5.17	1.84	0.77	0.78
other	18	242	8.96	5.06	3.52	5.06	1.76	0.82	0.76
cultivars	18	233	8.63	6.07	2.56	5.28	1.79	0.79	0.78
landraces	118	405	15.00	4.85	10.15	5.32	1.91	0.78	0.78

239 Na – number of alleles, Ne – number of effective alleles, I - Shannon's information index, Ho – observed  
240 heterozygosity, He - expected heterozygosity

241


242 According to the Bayesian clustering model implemented in STRUCTURE software, the most  
243 informative number of subgroups was two (K=2). The second largest peak of Delta K value was  
244 observed for three subgroups (K=3), while studied accessions also displayed classification at eight  
245 subgroups (K=8) (Fig. 1).

246 Affiliation to the gene pools was identified when maximum likelihood and Delta K values were  
247 two (K=2), assigning accession to Mesoamerican (M; marked red in Fig. 2a) or Andean (A; marked  
248 green) group. This was confirmed based on allocation of reference accessions for gene pool  
249 classification. Total of 91.9% genotypes had membership coefficients higher than 0.80, implying that  
250 the majority of samples were strongly assigned to the groups. In addition, 11 genotypes (8.1%) were  
251 with membership coefficients lower than 0.80, showing admixed genetic structure between two groups  
252 (gene pools) (Supplementary Material 1). First group (M) included 32 genotypes (23.5% of studied  
253 germplasm) and reference accessions with mainly Mesoamerican phaseolin type. It comprised 21  
254 landraces and 11 cultivars. A majority of genotypes in group M were from the *Albus* seed form (84%)  
255 with primarily medium (56%) and large (34%) seed weight. Second group (A) comprised 93 genotypes

256 (68.4% of studied germplasm) and reference accessions with mainly Andean phaseolin types (Fig. 2).  
 257 It included 88 landraces and five cultivars. Genotypes with coloured and medium to large seeds  
 258 predominated in this group (*Griseus*, *Aureus*, *Roseus* and *Versicolor* forms). Only 13 genotypes within  
 259 second group had white seed coats with large seed weight in general. Genotypes identified as potential  
 260 hybrids between the gene pools (11 accessions) according to STRUCTURE analysis were primarily  
 261 landraces (5 *Albus*, 3 *Aureus*, 1 *Griseus*, 1 *Versicolor* and 1 *other*) with phaseolin type T.

262 For K = 3 grouping pattern, further division of the Andean group into two additional subgroups  
 263 was observed. In this scenario, 88.9% of genotypes had membership coefficient higher than 0.80, while  
 264 the rest showed admixed origin between the three subgroups (11.1%). First subgroup, M (marked red  
 265 in Fig. 2b), remained the same, comprising 23.5% of studied germplasm. Second subgroup, A1 (marked  
 266 green), was composed of 45 landraces that belonged to mainly *Griseus*, *Roseus* and *Versicolor* forms,  
 267 two cultivars (Oplenac and Aster) of large white seeds and six reference accessions. Third subgroup,  
 268 A2 (marked blue), comprised 34 landraces of largely *Griseus* and *Versicolor* common bean forms, 4  
 269 cultivars (Rozalija, Žutotrban, Sremac and Slavonski žutozeleni) and three reference accessions.

270




271

272 Figure 1. Estimation of most likely subgroups number according to Evanno's method (Evanno et al.,  
 273 2005)

274

275 Another subdivision of a studied germplasm with classification criteria in which membership  
 276 coefficient was higher than 0.50, corresponding to seed traits patterns, was observed at K = 8. Common  
 277 bean landraces and cultivars from the *Albus* group clustered mainly in Mesoamerican M (red in Fig. 2c)  
 278 (84%) and Andean Ab (dark blue) (80%) subgroups. Genotypes of *Griseus* form predominated in  
 279 Andean Ae (light blue) (73.9%) and Aa (green) (41.2%) subgroups. *Versicolor* (47.1%) and *Roseus*  
 280 (42.1%) bean accessions were most numerous in Ac (yellow) subgroup. The most diverse was subgroup  
 281 Ad (pink), comprising all seed forms. The smallest number of accessions were classified in subgroups  
 282 Af (orange) (three accessions belonging to *other* forms) and Ag (brown) (two accessions from  
 283 *Versicolor* and one from *other* forms).




284  
 285  
 286  
 287 Figure 2. Estimation of population structure for common bean germplasm for a) K=2, b) K=3 and  
 288 c) K=8 using STRUCTURE software; group generated within Mesoamerican gene pool (M); groups  
 289 generated within Andean gene pool (A, A1, Aa, Ab, Ac, Ad, Ae, Af, Ag)

290  
 291 The relationships among the genotypes were assessed in more detail by hierarchical cluster  
 292 analysis (Figure 3). The Neighbour joining-based dendrogram divided 136 genotypes and reference



293 accessions into two main clusters (gene pools), Mesoamerican and Andean, with additional subclusters  
294 identified within each main cluster.



295  
296 Figure 3. Neighbour joining tree of 136 landraces and cultivars and 13 reference accessions based on  
297 similarity matrix (simple matching coefficient)

298  
299 In total, 32 genotypes (23.5%) were classified in Mesoamerican (M) cluster. Three subclusters  
300 are recorded within cluster M. Subcluster M1 included 9 landraces and 3 cultivars (Pasuljica P1,  
301 Panonski gradištanac, Belko) with seed weights from small to large. Subcluster M2 consisted of 12  
302 landraces and 6 cultivars (Balkan, Biser, Panonski tetovac, Poboļjšani gradištanac, Galeb, Maksa) with  
303 medium to large seeds. Two landraces and one cultivar (Medijana) clustered in subcluster M3. All  
304 accessions in this group had medium seed weight. Great majority of accessions in cluster M belonged  
305 to the *Albus* form.

306 The Andean (A) cluster comprised 75% of studied landraces and cultivars in total. Subcluster  
307 A1 consisted of 31 landraces and 2 cultivars (Sremac and Slavonski žutozeleni). Accessions largely  
308 belonged to *Grisues*, followed by *Versicolor* in form. In total, 79% of genotypes in subcluster A1  
309 corresponded to subgroup A2 identified in STRUCTURE analysis when K=3, while the rest were of

310 admixed origin. Therefore, it can be suggested that genotypes comprising these two subclusters  
311 represent novel variation, which was created in this region or was introduced from different sources.

312           Andean subcluster A2 in NJ dendrogram consisted of 40 landraces and 2 cultivars (Oplenac,  
313 Zlatko). Landraces and cultivars predominantly belonged to *Grisues* form (48%); however, other seed  
314 forms were observed in smaller number – *Roseus*, *Versicolor* and *Aureus*. Only two landraces had a  
315 white seed coat. Subcluster A3 comprised 22 landraces, three cultivars (Rozalija, Žutotrban, Aster) and  
316 seven reference accessions. The great majority of accessions belonged to *Versicolor* (32%) and *Albus*  
317 (28%) forms. Five landraces (L2, L54, L59, L109 and L110) and two reference accessions (PHA390  
318 and PHA131) deviated from this grouping pattern and were separated on the dendrogram.

319

#### 320 **4. Discussion**

321

322           For proper conservation, assessment of breeding value and organization and structuring of  
323 breeding programs, it is essential to identify levels of phenotypic and genetic diversity preserved in the  
324 germplasm of any crop species (Mhlaba et al., 2018). Common bean accessions examined in this paper  
325 have already been characterized for their phenotypic variability, while origin and dissemination of local  
326 bean germplasm from Serbia was assessed based on variation of phaseolin types (see Savić et al., 2020).  
327 Therefore, in order to further reveal genetic diversity preserved in studied germplasm, relatedness of  
328 landraces and cultivars on a molecular level, as well as their structure within and beyond gene pools of  
329 origin, the present study was performed, which should serve as complementary to the previous one.

330           Results of this research indicated that substantial allelic diversity was preserved in the  
331 germplasm from Serbia and reference accessions. Total number of alleles ( $N_a$ ) and gene diversity ( $H_e$ )  
332 were notably higher when compared to germplasm from Portugal, Italy and Croatia (Leitao et al., 2017;  
333 Carović-Stanko et al., 2017; Gioia et al., 2019). On the other hand, average number of alleles per locus  
334 was comparable to those reported in common bean core collection by Blair et al. (2009). Even though  
335 our material was collected from a relatively narrow geographic region and was smaller in size compared  
336 to the other research, obtained result could be due to high percentage of rare alleles (68.1%) identified  
337 in this paper. In addition, this research material included samples, mainly landraces, from both gene

338 pools belonging to various seed forms that represented great morphological diversity, which could have  
339 affected allelic variability recorded in present study. In addition, the geographic position of Serbia,  
340 which is a familiar trade and migration crossroad from east to west, might have been significant in  
341 shaping common bean diversity found in this region (Vasić et al., 2009).

342 Genetic variability was also measured as the amount of actual or potential heterozygosity.  
343 Observed heterozygosity ( $H_o$ ) represents the level of heterozygous individuals in the population  
344 compared with expected heterozygosity ( $H_e$ ) which reflects the genetic diversity at the specific loci  
345 along genotypes due to the degree of its out-crossing potential (Štajner, 2010; Pipan et al., 2013). In the  
346 case when  $H_o$  is equal to  $H_e$ , it means that the population is in Hardy-Weinberg equilibrium, among  
347 other random crosses. In addition, the small deviations between average  $H_o$  and  $H_e$  (0.01, Table 2) can  
348 indicate uniform abundance of alleles along Serbian common bean germplasm that could reflect their  
349 common genetic origin. Moreover, there are some loci where the deviation between  $H_o$  and  $H_e$  is higher  
350 than 0.3, i.e. loci ATA004, ATA007, ATA010, BM210 and ATA145 (Table 2). Those loci could be  
351 highly applicable to evaluate cross-pollination potential of common bean under field conditions.

352 Gene diversity ( $H_e$ ) recorded in this study, which is not dependable on the sample size, was  
353 much higher compared to the results of Leitao et al. (2017) and Carović-Stanko et al. (2017). On the  
354 other hand, it mostly corresponded to that found in bean germplasm from Western Balkan countries  
355 (former Yugoslav republics) by Maras et al. (2015). This could be due to frequent material exchange,  
356 gene flow between the countries that constituted former Yugoslavia, and a similar set of markers used.  
357 It was also revealed that gene diversity ( $H_e$ ) of each bean group generated according to seed form in  
358 this work was larger compared to that observed for common bean market classes from USA by Gioia  
359 et al. (2019). Bearing in mind that mentioned authors investigated elite advanced cultivars compared to  
360 landraces analysed in this research, it could be suggested that breeding interventions have narrowed the  
361 genetic basis of elite material in comparison to landraces. This was also proven with more allelic  
362 variants observed among landraces (405) compared to cultivars (233) in our study. Another interesting  
363 fact is that a larger percentage of rare alleles were found in landraces (67.6%) in relation to cultivars  
364 (29.6%), even though other parameters of genetic diversity were quite similar for these two groups,  
365 which also corresponded to values found for the entire collection. This revelation is important from a

366 breeding perspective, allowing breeders to use this unexplored variability preserved among landraces  
367 and cultivars in their advantage. Conversely, allelic variability differed among the groups. Accessions  
368 from *Albus* and *Griseus* forms proved to be the most variable, which could also be related to a  
369 proportionally larger number of accessions in these seed forms. Therefore, differences in allelic  
370 variability found in various other research could be in line with nature, number and variability of  
371 accessions assessed, geographic origin of studied material, DNA isolation and detection methods.

372         The genetic structure of the studied germplasm, on primary level, corresponded to familiar  
373 differentiation of common bean accessions according to gene pool affiliation, Mesoamerican or  
374 Andean. These results are in accordance with various investigations of *Phaseolus vulgaris* worldwide  
375 (Sicard et al., 2005; Kwak and Gepts, 2009; Blair et al., 2010; Raggi et al., 2013; Bitocchi et al., 2017).  
376 Separation of Serbian accessions into two recognized gene pools have already been shown based on  
377 phaseolin and phenotypic variation, with a considerably larger proportion of accessions belonging to  
378 the Andean gene pool (Savić et al., 2020), which is in accordance with results presented in this paper.  
379 The prevalence of Andean in contrast to Mesoamerican accessions, along with congruency in bean  
380 accessions clustering according to gene pool of origin (estimated at 95%) using different methods in  
381 these two research, was determined. It was also shown that a chosen set of SSR markers, including the  
382 combination of the PvSHP1 markers, proved their usefulness and efficiency in discriminating bean  
383 accessions according to the gene pool affiliation, as suggested by Nanni et al. (2011), Maras et al. (2015,  
384 2016), Pipan and Meglič (2019).

385         Moreover, application of molecular versus phenotypic markers revealed further subdivision of  
386 the studied germplasm from Serbia. Additional variation in respect to two recognized gene pools was  
387 identified in both STRUCTURE analysis when K=3 (subgroup A2) and based on genetic distance  
388 analysis combined with NJ dendrogram (subcluster A1). Accessions with membership coefficients of  
389 these two groups largely corresponded to each other. It was also acknowledged that this additional  
390 variation is concentrated within the Andean gene pool, which is in line with findings of Maras et al.  
391 (2015) for bean germplasm from Western Balkan countries, Raggi et al. (2013) for Italian and Leitao  
392 et al. (2017) for Portuguese beans. It is believed that potential sources of this distinctive variation could  
393 be accessions with admixed genetic bases derived from inter-gene pool crosses. Apart from that, it is

394 probable that unique variation was generated in this geographic area as a result of genotypic and  
395 phenotypic adaptation to local growing conditions during long period of cultivation. For the Serbian  
396 common bean germplasm, it was observed that mainly landraces with greenish-yellow (*Griseus*) and  
397 mottled (*Versicolor*) seeds comprised the mentioned clusters, and were marked as new variation. Since  
398 these seed types, apart from white-seeded cultivars and landraces, are favourite among the bean  
399 producers, it is possible that new variation is a result of farmer's selection towards most tolerant and  
400 high yielding landraces in those seed forms.

401 The theoretical speculation mentioned above is supported by the revelation of putative hybrid  
402 genotypes in STRUCTURE analysis. In the case where phaesolin type T predominated among these  
403 accessions, it was determined that genetic bases of this material was also within the Andean gene pool.  
404 Observed frequency of putative hybrids in this study was quite low (8.1%), but nonetheless in  
405 accordance with results of Blair et al. (2010), Gioia et al. (2013) and Maras et al. (2015). On the  
406 contrary, Angioi et al. (2010) observed larger percentage (44%) of accessions derived from inter-gene  
407 pool hybridization indicating a high contribution of admixed genotypes in the European common bean.  
408 Santalla et al. (2002), Logozzo et al. (2007) and Gioia et al. (2013) highlighted the significance of  
409 identification of such accessions in certain bean germplasm collections and their breeding value.  
410 Putative hybrids might possess new and interesting combination of traits created in inter-gene pool  
411 crosses, which could be related to higher adaptability to environmental stress, tolerance to pests and  
412 pathogens, better productivity and overcome negative correlation between seed weight and yield  
413 potential.

414 Relatedness of landraces and cultivars was discussed based on genetic distance analysis  
415 associated with Neighbour-joining clustering method, while subsequent genetic structure was revealed  
416 when K=8 was considered in STRUCTURE analysis. In both methods, it was obvious that landraces  
417 and cultivars formed subgroups according to their phenotype (with several exceptions). Arguably,  
418 however, a genetic distance-based method might better assist in differentiation among accessions of  
419 similar phenotypes and in selection of more genetically distant parenting components for breeding. In  
420 the case of common bean, it would support breeding of cultivars in specific seed type (form) with  
421 improved productivity. Different types of clustering methods in combinations with genetic distance

422 calculations were applied. Observing the fact that both landraces and cultivars were used within  
423 calculations, the combination of NJ and simple matching coefficient was the best choice to evaluate and  
424 visualise the genetic relations and distribution of Serbian genotypes according their genetic origin (gene  
425 pool determination). Moreover, the distribution of genotypes among clusters (A1-A3; M1-M3) on  
426 Figure 3 is also correlated with seed characteristics for both, Andean and Mesoamerican group,  
427 respectively.

428 A larger number of groups distinguished among landraces indicate existent diversity of this  
429 material, despite the fact that farmers have maintained landraces during longer period on the farms close  
430 to each other, exchanged seeds with surrounding farms and among themselves. Our results are in  
431 accordance with research of Masi et al. (2009) and Raggi et al. (2013) who made similar observations.  
432 Even though landraces clustered largely according to the seed traits, some deviations of this pattern  
433 were observed. The largest discrepancy was detected among accessions belonging to the *Albus* seed  
434 form. Even though these accessions are mainly of Mesoamerican origin, large white-seeded types were  
435 also observed in Andean groups. There are two possible explanations of this finding. Firstly, breeding  
436 intervention in Serbia were mostly done within white-seeded beans, which could have resulted in  
437 hybridization between the gene pools. Secondly, common bean landraces with white large seeds,  
438 belonging to Nueva Granada race were introduced and are grown in Serbia. In addition, accessions of  
439 *Roseus* and *Versicolor* group often clustered together. Although these accessions are usually  
440 distinguished by their phenotype, they were not well separated on molecular level in this research.

441 Distribution of Serbian cultivars in distinctive, genetically diverse subclusters in NJ  
442 dendrogram is a result of various selection and breeding criteria over time. It is well known that there  
443 is accepted tendency to satisfy market demands in term of making cultivars that phenotypically  
444 correspond to specific market class, or seed form (Geravandi et al., 2020). In various timeframes of  
445 common bean breeding in Serbia, different available material was used, which also included  
446 introduction of foreign germplasm for breeding purposes. This, together with adaptation of newly  
447 created material to environmental and growing conditions at the time, resulted in genetic divergence of  
448 Serbian common bean assortment. On the other hand, there were cases when two or more cultivars were

449 more closely positioned on the dendrogram. The most probable cause of this phenomenon is the  
450 common genetic origin they share, belonging to the same ancestral line.

451 In a conclusion, examined landraces and cultivars displayed marked genotypic variation, which  
452 allowed detailed description of diversity present in common bean germplasm from Serbia,  
453 accompanying the information on phenotypic variability previously assessed. Classification and  
454 structuring of the accessions in accordance within and beyond gene pool of origin should facilitate  
455 conservation strategies and breeding of this material. A combination of phenotypic and molecular data  
456 should allow researchers to make a selection of smaller number of accession for the core collections,  
457 which will further be assessed for their breeding value (productivity, nutritional value, tolerance to  
458 biotic and abiotic stress). As a result of the information presented in this research, it is hoped that the  
459 study of common bean genetic diversity in the Western Balkans and the rest of Europe has been  
460 purposefully expanded.

461

#### 462 **Acknowledgement**

463 This research was supported by the Ministry of Education, Science and Technological Development of  
464 the Republic of Serbia, Grant Number: 451-03-9/2021-14/200032, which covered experimental design,  
465 statistical data processing, manuscript writing and partially material analysis. It was also funded  
466 partially by the Slovenian Ministry of Agriculture, Forestry and Food and the Slovenian Research  
467 Agency, grants number P4-0072 and L4-7520, respectively. Under the applicative research project L4-  
468 7520 we have designed the study and performed analyses. The funding under Agrobiodiversity research  
469 pregame P4-0072 covered the writing procedure of the manuscript.

470

#### 471 **References**

472

473 **Angioi, S.A., Rau, D., Attend, G., Nanni, L., Bellucci, E., Logozzo, G., Negri, V., Spagnoletti Zeuli,**  
474 **P.L., Papa, R., 2010.** Beans in Europe: origin and structure of the European landraces of *Phaseolus*  
475 *vulgaris* L. Theor. Appl. Genet. 121, 829-843. <https://doi.org/10.1007/s00122-010-1353-2>

476 **Bitocchi, E., Rau, D., Bellucci, E., Rodriguez, M., Murgia, M., Gioia, T., Santo, D., Nanni, L.,**  
477 **Attene, G., Papa, R. 2017.** Beans (*Phaseolus* spp.) as a model for understanding crop evolution. Front.  
478 Plant Sci. 8: 722. <https://doi.org/10.3389/fpls.2017.00722>

479 **Blair, M., Diaz, J.M., Hidalgo, R., Diaz, L., Duque, M., 2007.** Microsatellite characterization of  
480 Andean races of common bean (*Phaseolus vulgaris* L.). Theor. Appl. Genet. 116 (1): 29-43.  
481 <https://doi.org/10.1007/s00122-007-0644-8>

482 **Blair, M., Diaz, L., Buendia, H., Duque, M. 2009.** Genetic diversity, seed size association and  
483 population structure of a core collection of common bean (*Phaseolus vulgaris* L.). Theor. Appl. Genet.  
484 119 (6), 955-972. <https://doi.org/10.1007/s00122-009-1064-8>

485 **Blair, M., Gonzales, L., Kimani, P., Butare, L. 2010.** Genetic diversity, inter-gene pool introgression  
486 and nutritional quality of common bean (*Phaseolus vulgaris* L.) from Central Africa. Theor. Appl.  
487 Genet. 121 (2), 237-248. <https://doi.org/10.1007/s00122-010-1305-x>

488 **Broughton, W.J., Hernandez, G., Blair, M., Beebe, S., Gepts, P., Vanderleyden, J., 2003.** Beans  
489 (*Phaseolus* spp.) - Model food legumes. Plant Soil 252, 55-128.  
490 <https://doi.org/10.1023/A:1024146710611>

491 **Carović-Stanko, K., Liber, Z., Vidak, M., Barešić, A., Grdiša, M., Lazarević, B., Šatović, Z. 2017.**  
492 Genetic diversity of Croatian common bean landraces. Frontiers in Plant Science 8, 604.  
493 <https://doi.org/10.3389/fpls.2017.00604>

494 **De Ron, A., Gonzalez, A., Rodino, A., Santalla, M., Godoy, L., Papa, R. 2016.** History of the  
495 common bean crop: its evolution beyond its areas of origin and domestication. Arbor 192-779. doi:  
496 10.3989/arbor.2016.779n3007

497 **Earl, D.A. and von Holdt, B.M.** STRUCTURE HARVESTER: a website and program for visualizing  
498 STRUCTURE output and implementing the Evanno method. Conserv. Genet. Resour. 3, 429-431.  
499 <https://doi.org/10.1007/s12686-011-9548-7>

500 **Evanno, G., Regnaut, S., Goudet, J. 2005.** Detecting the number of clusters of individuals using the  
501 software STRUCTURE: a simulation study. Mol. Ecol. 14, 2611-2620. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-294X.2005.02553.x)  
502 [294X.2005.02553.x](https://doi.org/10.1111/j.1365-294X.2005.02553.x)



503 **Gepts, P. and Bliss, F.A., 1985.** F1 hybrid weakness in the common bean: differential geographic  
504 origin suggests two gene pools in cultivated bean germplasm. *J. Hered.*76, 447-450.  
505 <https://doi.org/10.1093/oxfordjournals.jhered.a110142>

506 **Gepts, P., Osborn, T.C., Rashka, K., Bliss, F.A., 1986.** Phaseolin protein variability in wild forms  
507 and landraces of the common bean (*Phaseolus vulgaris*): evidence for multiple centers of domestication.  
508 *Econ. Bot.* 40, 451-468. <https://doi.org/10.1007/BF02859659>

509 **Gepts, P. and Bliss, F.A. 1988.** Dissemination pathways of common bean (*Phaseolus vulgaris*,  
510 Fabaceae) deduced from phaseolin electrophoretic variability. II. Europe and Africa. *Econ. Bot.* 42(1),  
511 86-104. <https://doi.org/10.1007/BF02859038>

512 **Gepts, P. 1999.** Development of an integrated genetic linkage map in common bean (*Phaseolus*  
513 *vulgaris* L.) and its use. In: Singh, S. Bean breeding for for the 21st century. Dordrecht: Kluwer  
514 Publishing House, 389-391.

515 **Geravandi, M., Cheghamirza, K., Farshadfar, E., Gepts, P. 2020.** QTL analysis of seed size and  
516 yield-related traits in an inter-gene pool population of common bean (*Phaseolus vulgaris* L.). *Sci. Hort.*  
517 274, 109678. <https://doi.org/10.1016/j.scienta.2020.109678>

518 **Gioia, T., Logozzo, G., Attene, G., Bellucci, E., Benedettelli, S., Negri, V., Papa, R., Spagnoletti**  
519 **Zeuli, P. 2013.** Evidence for introduction bottleneck and extensive inter-gene pool (Mesoamerican x  
520 Andean) hybridization in the European common bean (*Phaseolus vulgaris* L.) germplasm. *PLoS ONE*  
521 8(10), e75974. <https://doi.org/10.1371/journal.pone.0075974>

522 **Gioia, T., Logozzo, G., Marzario, S., Spagnoletti Zeuli, P., Gepts, P. 2019.** Evolution of SSR  
523 diversity from wild types to U.S. advanced cultivars in the Andean and Mesoamerican domestications  
524 of common bean (*Phaseolus vulgaris*). *PLoS ONE* 14(1), e0211342.  
525 <https://doi.org/10.1371/journal.pone.0211342>

526 **Hegay, S., Geleta, M., Bryngelsson, T., Asanaliev, A., Garkava-Gustavsson, L., Persson**  
527 **Hovmalm, H., Ortiz, R. 2013.** Genetic diversity analysis in *Phaseolus vulgaris* L. using morphological  
528 traits. *Genet. Resour. Crop Evol.* 61, 555-566. <https://doi.org/10.1007/s10722-013-0056-3>

529 **Kwak, M. and Gepts, P. 2009.** Structure of genetic diversity in the two major gene pools of common  
530 bean (*Phaseolus vulgaris* L., Fabaceae). *Theor. Appl. Genet.* 118, 979-992.  
531 <https://doi.org/10.1007/s00122-008-0955-4>

532 **Leitao, S.T., Dinis, M., Veloso, M.M., Šatović, Z., Vaz Patto, M.C. 2017.** Establishing the bases for  
533 introducing the unexplored Portuguese common bean germplasm into the breeding world. *Front. Plant.*  
534 *Sci.* 8: 1296. <https://doi.org/10.3389/fpls.2017.01296>

535 **Logozzo, G., Donnoli, R., Macaluso, L., Papa, R., Knupffer, H., Spagnoletti Zeuli, P. 2007.**  
536 Analysis of the contribution of Mesoamerican and Andean gene pools to European common bean  
537 (*Phaseolus vulgaris* L.) germplasm and strategies to establish a core collection. *Genet. Resour. Crop*  
538 *Evol.* 54, 1763-1779. <https://doi.org/10.1007/s10722-006-9185-2>

539 **Kelly, D.K., Kolkman, J.M., Schneider, K. 1998.** Breeding for yield in dry bean (*Phaseolus vulgaris*  
540 L.). *Euphytica* 102, 343-356. <https://doi.org/10.1023/A:1018392901978>

541 **Mallor, C., Barberan, M., Albar, J. 2018.** Recovery of common bean landrace (*Phaseolus vulgaris*  
542 L.) for commercial purposes. *Front. Plant Sci.* 9: 1440. <https://doi.org/10.3389/fpls.2018.01440>

543 **Maras, M., Pipan, B., Šuštar-Vozlič, J., Todorović, V., Đurić, G., Vasić, M., Kratovalieva, S.,**  
544 **Ibusoska, A., Agić, R., Matotan, Z., Čupić, T., Meglič, V. 2015.** Examination of genetic diversity of  
545 common bean from Western Balkans. *J. Am. Soc. Hortic. Sci.* 140(4), 308-316.  
546 <https://doi.org/10.21273/JASHS.140.4.308>

547 **Maras, M., Ibusoska, A., Kratovalieva, S., Agić, R., Šuštar-Vozlič, J., Meglič, V. 2016.** Genetic  
548 diversity of common bean accessions from Former Yugoslav Republic of Macedonia as revealed by  
549 molecular and morphological markers. *Genetika* 48 (2), 729-742.  
550 <https://doi.org/10.2298/GENSR1602729M>

551 **Masi, P., Logozzo, G., Donini, P., Spagnoletti Zeuli, P. 2009.** Analysis of genetic structure in widely  
552 distributed common bean landraces with different plant growth habits using SSR and AFLP markers.  
553 *Crop Sci.* 49, 187-199. <https://doi.org/10.2135/cropsci2008.05.0265>

554 **McClellan, P. and Lee, R. 2007.** Genetic architecture of chalcone isomerase non-coding regions in  
555 common bean (*Phaseolus vulgaris* L.). *Genome* 50 (2), 203-214. <https://doi.org/10.1139/g07-001>

556 **Mhlaba, Z.B., Mashilo, J., Shimelis, H., Assefa, A.B., Modi, A.T. 2018.** Progress in genetic analysis  
557 and breeding of tepary bean (*Phaseolus acutifolius* A. Gray): A review. *Sci. Hort.* 237, 112-119.  
558 <https://doi.org/10.1016/j.scienta.2018.04.012>

559 **Nanni, L., Bitocchi, E., Bellucci, E., Rossi, M., Rau, D., Attene, G., Gepts, P., Papa, R. 2011.**  
560 Nucleotide diversity of a genomic sequence similar to SHATTERPROOF (PvSHP1) in domesticated  
561 and wild common bean (*Phaseolus vulgaris* L.). *Theor. Appl. Genet.* 123(8), 1341-1357.  
562 <https://doi.org/10.1007/s00122-011-1671-z>

563 **Park, S. 2001.** Microsatellite Toolkit. Department of Genetics, Trinity College, Dublin.

564 **Peakall, R. and Smouse, P.E. 2006.** GenAlex 6: genetic analysis in excel. Population genetic software  
565 for teaching and research. *Mol. Ecol. Notes* 6, 288-295. [https://doi.org/10.1111/j.1471-](https://doi.org/10.1111/j.1471-8286.2005.01155.x)  
566 [8286.2005.01155.x](https://doi.org/10.1111/j.1471-8286.2005.01155.x)

567 **Pipan, B., Šuštar-Vozlič, J., Meglič, V. 2013.** Genetic differentiation among sexually compatible  
568 relatives of *Brassica napus* L. *Genetika* 45 (2), 309-327. <https://doi.org/10.2298/GENSR1302309P>

569 **Pipan, B. and Meglič, V. 2019.** Diversification and genetic structure of the western-to-eastern  
570 progression of European *Phaseolus vulgaris* L. germplasm. *BMC Plant Biol.* 19, 442.  
571 <https://doi.org/10.1186/s12870-019-2051-0>

572 **Pritchard, J.K., Wen, X., Falush, D. 2009.** Documentation for STRUCTURE software: version 2.3.  
573 USA: Department of Human Genetics, University of Chicago and Department of Statistics, University  
574 of Oxford.

575 **Raggi, L., Tiranti, B., Negri, V. 2013.** Italian common bean landraces: diversity and population  
576 structure. *Genet. Resour. Crop Evol.* 60, 1515-1530. <https://doi.org/10.1007/s10722-012-9939-y>

577 **Rodino, P., Santalla, M., Gonzalez, A.M., De Ron, A.M., Singh, S.P. 2006.** Novel genetic variation  
578 in common bean from the Iberian Peninsula. *Crop Sci.* 46, 2540-2546.  
579 <https://doi.org/10.2135/cropsci2006.02.0104>

580 **Savić, A., Zorić, M., Brdar-Jokanović, M., Zdravković, M., Dimitrijević, M., Petrović, S.,**  
581 **Živanov, D., Vasić, M. 2020.** Origin and diversity study of local common bean (*Phaseolus vulgaris*  
582 L.) germplasm from Serbia: phaseolin and phenotyping approach. *Genet. Resour. Crop Evol.*  
583 <https://doi.org/10.1007/s10722-020-00974-9>

584 **Santalla, M., Rodino, P., De Ron, A.M. 2020.** Allozyme evidence supporting southwestern Europe as  
585 a secondary centre of genetic diversity for the common bean. *Theor. Appl. Genet.* 104, 934-944.  
586 <https://doi.org/10.1007/s00122-001-0844-6>

587 **Scarano, D., Rubio, F., Ruiz, J.J., Rao, R., Corrado, G. 2014.** Morphological and genetic diversity  
588 among and within common bean (*Phaseolus vulgaris* L.) landraces from the Campania region (Southern  
589 Italy). *Sci. Hortic.* 180, 72-78. <https://doi.org/10.1016/j.scienta.2014.10.013>

590 **Sicard, D., Nanni, L., Porfiri, O., Bulfon, D., Papa, R. 2005.** Genetic diversity of *Phaseolus vulgaris*  
591 L. and *P. coccineus* L. landraces in central Italy. *Plant Breed.* 124, 464-472.  
592 <https://doi.org/10.1111/j.1439-0523.2005.01137.x>

593 **Singh, S.P., Gutierrez, J.A., Molina, A., Urrea, C., Gepts, P., 1991.** Genetic diversity in cultivated  
594 common bean. II. Marker based analysis of morphological and agronomic traits. *Crop Sci.* 31, 23-29.  
595 <https://doi.org/10.2135/cropsci1991.0011183X003100010005x>

596 **Sinkovič, L., Pipan, B., Sinkovič, E., Meglič, V. 2019.** Morphological seed characterization of  
597 common (*Phaseolus vulgaris* L.) and runner (*Phaseolus coccineus* L.) bean germplasm: A Slovenian  
598 gene bank example. *Biomed Res. Int.*, 6376948. <https://doi.org/10.1155/2019/6376948>

599 **Sitohy, M., Deskoy, E.M., Osman, A., Rady, M.R. 2020.** Pumpkin seed protein hydrolysate treatment  
600 alleviates salt stress effects on *Phaseolus vulgaris* by elevating antioxidant capacity and recovering ion  
601 homeostasis. *Sci. Hort.* 271, 109495. <https://doi.org/10.1016/j.scienta.2020.109495>

602 **Štajner, N. 2010.** Mikrosatelitski markerji uporabni za identifikacijo kultivarjev vinske trte (*Vitis*  
603 *vinifera* L.). *Acta Agric. Slov* 95.2, 183-192. <http://aas.bf.uni-lj.si/julij2010/08stajner.pdf>

604 **Šuštar-Vozlič, J., Maras, M., Javornik, B., Meglič, V. 2006.** Genetic diversity and origin of Slovene  
605 common bean (*Phaseolus vulgaris* L.) germplasm as revealed by AFLP markers and phaseolin analysis.  
606 *J. Am.Soc. Hortic. Sci.* 131(2), 242-249. <https://doi.org/10.21273/JASHS.131.2.242>

607 **Vasić, M., Vujičić, B., Tepić, A., Gvozdanović-Varga, J., Šumić, Z. 2009.** Dietary fibre content in  
608 some dry beans. *Acta Periodica Technologica* 40, 103-110. <https://doi.org/10.2298/APT0940103V>

609 **Yu, K.F., Park, S.J., Poysa, V. 1999.** Abundance and variation of microsatellite DNA sequences in  
610 beans (*Phaseolus* and *Vigna*). *Genome* 42, 27-34. <https://doi.org/10.1139/g98-100>

611 **Zeven, A.C. 1997.** The introduction of the common bean (*Phaseolus vulgaris* L.) into western Europe  
612 and the phenotypic variation of dry beans collected in the Netherlands in 1946. *Euphytica* 94, 319-328.  
613 <https://doi.org/10.1023/A:1002940220241>  
614 **Zhang, X., Blair, M.W., Wang, S. 2008.** Genetic diversity of Chinese common bean (*Phaseolus*  
615 *vulgaris* L.) landraces assessed with simple sequence repeat markers. *Theor. Appl. Genet.* 117, 629-  
616 640. <https://doi.org/10.1007/s00122-008-0807-2>