



# EUFORIA

European Forest Research  
and Innovation Area

## EUFORINNO

*European Forest Research and Innovation*

The final EUFORINNO conference  
31<sup>st</sup> August - 4<sup>th</sup> September 2015,  
Rogla, Slovenia

*Programme and Book of Abstracts*

### PROGRAMME COMMITTEE

**CHAIR:**

Hojka Kraigher (SFI)

**EUFORINNO project partners:**

Reinhart Ceulemans (PLECO, B), Monika Konnert (ASP, D), Dieter Ernst (Helmholtz, D), Zachary Kayler (ZALF, D), Danny McCarroll (Uni. SWANSEA, UK), Giorgio Matteucci (IBAF-CNR, I), Heljä-Sisko Helmisaari (Uni. HELSINKI, SF), Håkan Wallander (Uni. LUND, S)

**EUFORINNO team at SFI:**

Primož Simončič (Director & Co-Chair), Robert Robek (WP1 leader), Tine Grebenc (WP2 leader), Tom Levanič (WP3 leader), Jožica Gričar (Chair of SFI Scientific Board)

**EUFORINNO scientific recruits at SFI:**

Nate McDowell, Tanja Mrak, Saša Zavadlav, Ivan Kreft

### LOCAL ORGANISING COMMITTEE

**Chair:** Tine Grebenc, **Secretary:** Tina Drolc, **Treasurer and sponsors:** Tjaša Baloh, **Website:** Žiga Lipar



Electronic version available from:  
<http://eprints.gozdis.si/1440/>



This project is funded by  
the European Union

**Publisher:**

Slovenian Forestry Institute, *Silva Slovenica* Publishing Centre,  
Ljubljana 2015

**Editorial Board of The *Silva Slovenica* Publishing Centre:**

prof. dr. Tom Levanič, Chief; dr. Andreja Ferreira, dr. Barbara Piškur,  
prof. dr. Dušan Jurc, dr. Gregor Božič, prof. dr. Hojka Kraigher, doc. dr.  
Jožica Gričar, dr. Lado Kutnar, dr. Marko Kovač, doc. dr. Matjaž Čater,  
dr. Mitja Ferlan, dr. Nike Kranjc, dr. Nikica Ogris, dr. Primož Simončič,  
dr. Robert Robek, dr. Tine Grebenc, dr. Urša Vilhar

**Title:**

Programme and Book of Abstracts: EUFORIA: European Forest  
Research and Innovation Area, The final EUFORINNO Scientific  
Conference, September 2015, Rogla, Slovenia

**Editor-in-Chief:**

Prof. dr. Hojka Kraigher

**Editorial Board of Programme and Book of Abstracts:**

Prof. dr. Hojka Kraigher, dr. Tine Grebenc

**Technical editors:**

Natalija Senčar, Magda Špenko, Tine Grebenc

**Design:**

Sonja Rutar

**Cover page photos:**

Domen Finžgar - LIFE GENMON forest genetic monitoring plot for Silver fir

**Printed by:**

DMB projekt d.o.o., Trbovlje

**Circulation:**

250 copies

**Price:**

free

**Co-financing:**

The publication was financed by EUFORINNO, European Forest  
Research and Innovation, 7<sup>th</sup> FP EU Capacities RegPot No. 315982

**Electronic issue:**

<http://eprints.gozdis.si/1440/>

## FOREWORD

European forests are critical to carbon storage, biodiversity conservation, water quality and a number of ecosystem services, socio-demographic stability and economic revenue. EUFORINNO (<http://euforinno.gozdis.si/presentation/>) is a means for the Slovenian Forestry Institute (SFI) to raise its scientific excellence and infrastructure, and better exploit its innovative outputs.

The project has brought so far an unprecedented increase in the scientific infrastructure, excellence and visibility of SFI, and contributed to awareness on the role of forests and forestry for nature and people. It has contributed to a deeper integration within the European Research Area (ERA), and with its final conference it contributes to the European Forest Research and Innovation Area – EUFORIA - an increase of forestry research excellence and awareness throughout Europe.

EUFORINNO relies on 8 excellent research partners who support SFI in building its long-term strategy, welcome researchers from SFI during their secondments abroad, and provide on-site training on new methodologies, data interpretation and publishing. In the organization of the final conference the partner leaders, and members of the EUFORINNO Advisory Board, have agreed to take an active role in the conference programme committee and are contributing their own keynote presentations in the different conference sections.

Without our great partners, Reinhart Ceulemans, Monika Konnert, Dieter Ernst, Zachary Kayler, Danny McCarroll, Giorgio Matteucci, Heljä-Sisko Helmisaari, and Håkan Wallander, there would be no project, and no EUFORIA!

Together with our most distinguished invited keynote speakers, Petr Baldrian, Cristina Cruz, John Dighton, Arthur Gessler, Jožica Gričar, John King, Antoine Kremer, Hrvoje Marjanović, Rainer Matyssek, Nate McDowell, Emilia Gutiérrez Merino, Hans Göransson, and Mike Starr, they are a guarantee for an outstanding scientific conference presenting the latest research and expanding horizons in forest science and innovation worldwide.

Furthermore, the project has relied on permanent collaboration, exchange of students and developing common strategies within the South-East European region; we have organized a number of regional workshops on the main EUFORINNO topics with the help of our friends and experts from the region, the National Contact Points (NCP), especially Prof. Saša Orlović, Prof. Dalibor Ballian, Doz. Elvis Toromani, Prof. Phil Aravanopoulos, and Prof. Ibrahim Ortaş, while other NCPs, together with our partners, have contributed with their lectures, students, and topics. We are happy that we can welcome almost all our NCPs and their colleagues also at the final EUFORINNO conference.

Yet the essence of the project is the EUFORINNO management and working group leaders, and all the colleagues who have built up an effective project team, with its core group of early stage researchers, both employed within the project and contributing to it through active participation at secondments, trainings, workshops, conferences, through guardianship of new equipment, developing and producing standardized operation procedures, building up the new open access repository, contributing to the publishing centre, and preparing the IP management strategy. It was a pleasure to collaborate with all of you!

Finally, thanks to the local organizing committee, and all collaborators from different departments of SFI, and co-organizers of the Round table, adding their hands to the organization of all events during the conference!

I am looking forward to the lectures, presentation of EUFORINNO impacts, and posters, and hope to finalize the conference with an effective contribution to further development and recognition of forests, forest science, and forestry in the European forest research and innovation area.

Hojka Kraigher  
Chair of the Programme Committee & EUFORINNO Project Coordinator

## Foreword

## List of Abstracts

<b>Kremer A.:</b> ADAPTIVE RESPONSES OF TREES TO CLIMATE CHANGE	1
<b>Konnert M. et al.:</b> GENETIC MONITORING IN FORESTS – EFFORTS FOR A EUROPEAN IMPLEMENTATION	2
<b>Alexandrov A. H.:</b> ENDEMIC FOREST TREE SPECIES IN SOUTHEAST EUROPE, THEIR GENETIC RESOURCES AND USE	3
<b>Kandemir G. et al.:</b> DETERMINATION OF GENETIC VARIATION EXISTING IN ANATOLIAN BLACK PINE NATURAL STAND AND PLANTATION AREA IN KASATURA BAY	4
<b>Westergren M. et al.:</b> FOREST GENETIC MONITORING – DELINEATION OF MONITORING REGIONS ON A TRANSECT FROM BAVARIA TO GREECE	5
<b>Čortan D. et al.:</b> INTER AND INTRA-POPULATION VARIATION OF LEAF STOMATAL TRAITS OF <i>Populus nigra</i> L. IN VOJVODINA, NORTHERN SERBIA	6
<b>De Boeck H. et al.:</b> BIODIVERSITY-STABILITY RELATIONSHIPS UNDER CLIMATE EXTREMES	7
<b>Ernst D. et al.:</b> EUROPEAN WHITE BIRCH AND RAGWEED: IMPACT OF AIR POLLUTIONS ON THE ALLERGENIC POLLEN, AN OVERVIEW	8
<b>Ceulemans R. et al.:</b> BIO-ENERGY FROM BIOMASS: FULL GREENHOUSE GAS BALANCE, ENERGY BALANCE AND ENVIRONMENTAL LIFE CYCLE ASSESSMENT	9
<b>King J. S. et al.:</b> QUANTIFICATION OF LINKED CARBON AND WATER CYCLES IN A LOBLOLLY PINE-SWITCH GRASS INTERCROPPING SYSTEM FOR BIOENERGY AND WOOD PRODUCTS	10
<b>Eler K. et al.:</b> SUCCESSION INDUCED CHANGES IN ECOSYSTEM FUNCTIONING WITH EMPHASIS ON CARBON AND WATER EXCHANGE	11
<b>Gutiérrez E.:</b> EFFECTS OF WARMING ON <i>Pinus sylvestris</i> IBERIAN FORESTS	12
<b>McCarroll D.:</b> STABLE ISOTOPES IN TREE RINGS	13
<b>Gričar J.:</b> WHAT INFORMATION IS STORED IN PHLOEM STRUCTURE AND RELATIONSHIP BETWEEN PHLOEM AND XYLEM INCREMENTS?	14
<b>Ionel P. et al.:</b> STAND DYNAMICS AND DISTURBANCE HISTORY IN MIXED FOREST OF NORWAY SPRUCE AND CEMBRA PINE FROM EASTERN CARPATHIANS	15
<b>Hafner P. et al.:</b> ENVIRONMENTAL SIGNALS IN <i>Quercus robur</i> L. TREE-RING WIDTHS AND $\delta^{13}\text{C}$	16
<b>Levanič T. et al.:</b> CLIMATE SIGNAL AND POTENTIAL OF BOSNIAN PINE ( <i>Pinus heldreichii</i> CHRIST) FOR CLIMATE RECONSTRUCTION IN CENTRAL W BALKAN REGION	17
<b>Gessler A.:</b> THE IMPACT OF ENVIRONMENTAL FACTORS ON BIOGEOCHEMICAL CYCLES IN TREES AND FOREST ECOSYSTEMS - FROM MOLECULAR MECHANISMS TO ECOSYSTEM FUNCTIONING	18
<b>Kayler Z. et al.:</b> DROUGHT AND HEAT STRESS EFFECTS ON THE PLANT-SOIL CARBON CONTINUUM AND BELOWGROUND MICROBIAL COMMUNITIES	19

<b>Matteucci G. et al.:</b> RESPONDING TO CHALLENGES IN FOREST RESEARCH AND MONITORING: FROM PAST LEGACY TO INTEGRATION OF RESEARCH INFRASTRUCTURES	20
<b>Baldrian P.:</b> THE EFFECT OF FOREST TREES ON MICROBIAL PROCESSES INVOLVED IN CARBON CYCLING	21
<b>Göransson H. et al.:</b> TREE BIODIVERSITY AND ECOSYSTEM FUNCTION	22
<b>Helmisaari H-S. et al.:</b> FINE-ROOT TRAITS AND SOIL PROCESSES IN BOREAL FORESTS	23
<b>Leppälammki-Kujansuu J. et al.:</b> CARBON INPUT INTO SOIL ORIGINATING FROM FINE ROOT AND FOLIAGE LITTER	24
<b>Makita N. et al.:</b> FINE ROOT PHYSIOLOGICAL AND MORPHOLOGICAL TRAITS OF FINNISH BOREAL TREE SPECIES	25
<b>Dighton J. et al.:</b> DISTURBANCE AND RESILIENCE IN FORESTS: EXAMPLES FROM NEW JERSEY PINE BARRENS	26
<del>Cristina C.</del> <b>et al.:</b> IMPACTS OF INCREASED NITROGEN AVAILABILITY ON THE STRUCTURE AND FUNCTIONING OF A MEDITERRANEAN BASIN ECOSYSTEM	27
<b>Kreft I.:</b> FOREST – AGRICULTURE INTERFACE: A CASE OF BUCKWHEAT	28
<b>Kovič Dine M.:</b> PROTECTION OF FORESTS UNDER INTERNATIONAL LAW	29
<b>Starr M. et al.:</b> IMPACT OF FOREST HARVESTING PRACTICES ON BASE CATION BUDGETS – A SUSTAINABILITY STUDY USING MEASURED AND MODELLED DATA	30
<b>Horemans J. A. et al.:</b> VARIANCE DECOMPOSITION OF STEM BIOMASS INCREMENT PREDICTIONS FOR EUROPEAN BEECH	31
<b>Vieira J. et al.:</b> ADJUSTMENT CAPACITY OF MARITIME PINE CAMBIAL ACTIVITY IN DROUGHT-PRONE ENVIRONMENTS	32
<b>Ahmed I. U. et al.:</b> ABOVE AND BELOWGROUND TREE BIOMASS ALLOCATIONS- AS INFLUENCED BY SPECIES COMPOSITION IN TEMPERATE DECIDUOUS PLANTATION FOREST	33
<b>Marjanović H. et al.:</b> CARBON CYCLING IN LOWLAND OAK FOREST ECOSYSTEMS	34
<b>McDowell N.:</b> ACCELERATING FOREST MORTALITY	35
<b>Matyssek R.:</b> OZONE RESEARCH, <i>QUO VADIS?</i> LESSONS FROM THE FREE-AIR CANOPY FUMIGATION EXPERIMENT AT KRANZBERG FOREST	36
<b>Kraigher H. et al.:</b> OVERVIEW OF EUFORINNO PROJECT	37
<b>Bajc M. et al.:</b> MODERN LABORATORIES AND STANDARDISATION OF PROTOCOLS	38
<b>Žlindra D. et al.:</b> EUFORINNO METHODS TOWARDS FORMAL ACCREDITATION	39
<b>Mrak T.:</b> PROGRESS OF MICROSCOPY FACILITIES WITHIN EUFORINNO PROJECT	40
<b>Zavadlav S. et al.:</b> USE OF STABLE ISOTOPES IN FOREST ECOSYSTEM STUDIES: ON THE PROGRESS OF STABLE ISOTOPE LABORATORY AT THE SFI	41
<b>Ferlan M.:</b> MEASURING ISOTOPIC COMPOSITION OF CARBON DIOXIDE IN THE AIR	42

<b>Peteh M. et al.:</b> THE PUBLISHING CENTER <i>SILVA SLOVENICA</i> AND MONOGRAPH SERIES <i>STUDIA FORESTALIA SLOVENICA</i>	43
<b>Grebenc T. et al.:</b> <i>ACTA SILVAE ET LIGNI</i>	44
<b>Peteh M. et al.:</b> <i>SCIVIE – SCIENCES DE LA VIE</i> FROM THE PAST AND FOR THE FUTURE	45
<b>Kajba D. et al.:</b> LEAF UNFOLDING VARIABILITY IN CLONAL SEED ORCHARD OF <i>Fraxinus angustifolia</i> Vahl	47
<b>Ballian D. et al.:</b> PHENOLOGICAL VARIABILITY OF EUROPEAN BEECH ( <i>Fagus sylvatica</i> L.) IN THE INTERNATIONAL PROVENANCE TRIAL	48
<b>Bashkim T. et al.:</b> THE IMPACTS OF FOREST FIRES IN THE CURRENT STATE OF FOREST RESOURCES IN KOSOVO	49
<b>Rasztovits E. et al.:</b> RETREATING SESSILE OAK FOREST WITH IMPROVING VITALITY – INCLUDING TREE MORTALITY IN VITALITY ASSESSMENT	50
<b>Božič G. et al.:</b> GENETIC ANALYSIS OF NATIVE EUROPEAN BLACK POPLAR ( <i>populus nigra</i> L.) POPULATIONS IN SLOVENIA AND CROATIA BY USING SSR MARKERS	51
<b>Finžgar D. et al.:</b> IMPLEMENTING FOREST GENETIC MONITORING PLOTS	52
<b>Galović V. et al.:</b> TESTING OF SALT TOLERANCE ASSOCIATED GENES IN SERBIAN POPLAR CLONES	53
<b>Galović V. et al.:</b> MOLECULAR ANALYSES OF ITS rDNA OF THREE AUTOCHTHONOUS FUNGAL SPECIES FROM MOUNTAIN FORESTS IN SERBIA	54
<b>Grebenc T. et al.:</b> HYPOGEOUS FUNGI DIVERSITY AND ECOLOGY IN SE EUROPE	55
<b>Hrenko M. et al.:</b> TYPES OF ECTOMYCORRHIZA IN THE 34 YEARS OLD <i>Pinus sylvestris</i> L. SEED PLANTATION IN THE LOWLAND FOREST SITE “MURSKA ŠUMA”	56
<b>Hukić E. et al.:</b> SOIL PROPERTIES IN RELATION TO SELECTIVE LOGGING IN BEECH AND FIR FOREST STANDS - CASE STUDY AT BJELAŠNICA MOUNTAIN	57
<b>Katanić M. et al.:</b> SEASONAL VARIATION OF ECTOMYCORRHIZAL COMMUNITY FROM MATURE POPLAR PLANTATION	58
<b>Kavčič A.:</b> POSSIBILITIES FOR THE USE OF LASER VIBROMETRY IN THE MANAGEMENT OF INSECT PESTS	59
<b>Kern A. et al.:</b> FOREST PHENOLOGY IN CENTRAL-EUROPE BASED ON MODIS DATA	61
<b>Kiourtsis F. et al.:</b> THE IMPORTANCE OF SEEDLINGS QUALITY IN TIMBER AND BIO-ENERGY PRODUCTION	62
<b>Kobal M.:</b> ROCKFALL MONITORING BY THE COMBINATION OF LIDAR AND UNMANNED AERIAL VEHICLE TECHNOLOGY	63
<b>Kühndorf K.:</b> <i>Leotia cf. lubrica</i> FORMS ARBUTOID MYCORRHIZA WITH <i>Comarostaphylis arbutoides</i> (ERICACEAE)	64
<b>Kutnar L.:</b> EFFECTS OF DIFFERENT FOREST MANAGEMENT INTENSITIES ON BIODIVERSITY IN NATURA 2000 FOREST HABITATS	65

<b>Lavrič M. et al.:</b> XYLEM SAP FLOW AND RADIAL GROWTH IN <i>QUERCUS PUBESCENS</i> WILLD. FROM ABANDONED GRASSLANDS IN SLOVENIAN KARST REGION	66
<b>Lazarević J. et al.:</b> WILD EDIBLE MUSHROOMS AND MEDICAL HERBS ON MONTENEGRIN MOUNTAINS AND KATUNS	68
<b>Lazarević J.:</b> APPLICATION OF AUTOCHTHONOUS FUNGI FOR MYCORRHIZATION OF CONIFERS –TRIALS IN MONTENEGRO	69
<b>Luthar Z. et al.:</b> STUDENTS' ATTITUDE TOWARDS COMMERCIALIZATION OF TRANSGENIC FOREST TREES IN SLOVENIA	70
<b>Martinović T. et al.:</b> FUNGAL AND BACTERIAL COMMUNITIES IN SPRUCE WOOD SAMPLES EXPOSED TO OUTDOORS ENVIRONMENTAL CONDITIONS	71
<b>Matović B. et al.:</b> COMPARISON OF STAND STRUCTURE IN MANAGED AND VIRGIN EUROPEAN BEECH FORESTS IN SERBIA	72
<b>Mrak T. et al.:</b> ECTOMYCORRHIZA BETWEEN <i>Scleroderma Areolatum</i> EHRENB. AND <i>Fagus sylvatica</i> L.	73
<b>Ortas I. et al.:</b> TOTAL AMOUNT OF CARBON HOLD UNDER DIFFERENT AGES EUCALYPTUS TREES, PLANTS WITH DIFFERENT SOIL AND CROP MANAGEMENT SYSTEMS IN COST OF THE MEDITERRANEAN CONDITIONS	74
<b>Pinuela Samaniego Y.:</b> INFLUENCE OF HEAVY FOREST MACHINERY ON FINE ROOTS: OBSERVATIONS ON MINIRHIZOTRON DATA	75
<b>Rantaša B.:</b> VISIBILITY AND DISSEMINATION IN FOREST GENETIC MONITORING	76
<b>Rupel M. et al.:</b> OZONE IN URBAN FORESTS	77
<b>Sinjur I. et al.:</b> SOIL RESPIRATION VARIABILITY IN BOREAL PINE FOREST IN VÄRRÖ, FINLAND	78
<b>Toromani E. et al.:</b> RADIAL GROWTH RESPONSES OF <i>PINUS HALEPENSIS</i> MILL. AND <i>Pinus pinea</i> L. FORESTS TO CLIMATE VARIABILITY IN WESTERN ALBANIA	79
<b>Unuk T. et al.:</b> LINKAGES OF PLANT MORPHOLOGICAL-FUNCTIONAL TRAITS TO SOIL PROPERTIES ON EXTENSIVELY USED GRASSLANDS OF POHORJE MOUNTAINS	80
<b>Železnik P. et al.:</b> FINE ROOT DYNAMICS IN SLOVENIAN BEECH FORESTS IN RELATION TO SOIL TEMPERATURE AND WATER AVAILABILITY	82
<b>Železnik P. et al.:</b> ROOT GROWTH DYNAMICS OF THREE BEECH ( <i>Fagus sylvatica</i> L.) PROVENANCES	83

## Programme overview

	Monday, August 31 <sup>st</sup>	Tuesday, September 1 <sup>st</sup>	Wednesday, September 2 <sup>nd</sup>	Thursday, September 3 <sup>rd</sup>	Friday, September 4 <sup>th</sup>
08:00 - 09:00		Arrival and registration	Biodiversity & ecosystem processes	Forest disturbance, management and modelling	Round table: Forest science and innovation in Europe
09:00 - 09:30		Opening of the conference			
09:30 - 11:00		Forest genetic diversity and climate change			
11:00 - 12:00					Press conference
12:00 - 12:30					
12:30 - 13:00					Closure of the conference
13:00 - 14:00		Lunch			Lunch
14:00 - 15:00	Arrival and registration	Carbon dynamics in time and space	Excursion to a forest reserve/ research plot	Future forest research and innovation in Europe	Departure or participants
15:00 - 18:00				Impacts of EUFORINNO	
18:00 - 19:00	Welcome mixer	Posters with pizza and beverages	Local tourism with wines of the region		
19:00 - 19:30					
19:30 - 21:00				Conference dinner and awards ceremony	
21:00 - 22:00					

## Scientific programme

<b>Tuesday September 1<sup>st</sup></b>		
09:00-09:30	<i>Opening of the conference</i>	
<b>Session 1</b>	<b><i>Forest genetic diversity and climate change</i></b>	<b><i>Moderators: Monika Konnert &amp; Marjana Westergren</i></b>
09:30-10:00	<b>Antoine KREMER (KN1)</b>	<b>Adaptive responses of trees to climate change</b>
10:00-10:30	<b>Monika KONNERT (KN2)</b>	<b>Genetic monitoring in forests - LIFE GENMON</b>
10:30-10:45	Alexander ALEXANDROV	Endemic tree species in SEE, their genetic resources and use ...
10:45-11:00	Gaye KANDEMIR	Determination of genetic variation in Anatolian black pine...
11:00-11:30	<i>Coffee and tea</i>	
11:30-11:45	Marjana WESTERGREN	Forest genetic monitoring – delineation of monitoring ...
11:45-12:00	Dijana ČORTAN	Inter and intra-population variation of leaf stomatal traits...
12:00-12:15	Hans de BOECK	Biodiversity-stability relationships under climate extremes
12:15-12:45	<b>Dieter ERNST (KN3)</b>	<b>European white birch and ragweed: ...allergenic pollen...</b>
12:45-14:00	<i>Lunch</i>	
<b>Session 2</b>	<b><i>Carbon dynamics in time and space</i></b>	<b><i>Moderators: Danny McCarroll and Tom Levanič</i></b>
14:00-14:30	<b>Reinhart CEULEMANS (KN4)</b>	<b>Bio-energy from biomass: Full GHG, energy balance, LCA...</b>
14:30-15:00	<b>John KING (KN5)</b>	<b>Quantification of linked carbon and water cycles ....</b>
15:00-15:15	Klemen ELER	Succession induced changes in ecosystem functioning...
15:15-15:45	<b>Emilia GUTIÉRREZ (KN6)</b>	<b>Effects of warming on <i>Pinus sylvestris</i> Iberian forests</b>
15:45-16:15	<i>Coffee and tea</i>	
16:15-16:45	<b>Danny McCARROLL (KN7)</b>	<b>Stable isotopes in tree rings</b>
16:45-17:15	<b>Jožica GRIČAR (KN8)</b>	<b>What information is stored in phloem structure...</b>
17:15-17:30	Ionel POPA	Stand dynamics and disturbance history in mixed forests...
17:30-17:45	HAFNER Polona	Environmental signals in <i>Quercus robur</i> tree ring widths...
17:45-18:00	Tom LEVANIČ	Climate signal and potential of Bosnian pine <i>P. heldreichii</i> ...
18:00-21:00	<b><i>Posters with pizza and beverages</i></b>	

## Scientific programme

<b>Wednesday September 2<sup>nd</sup></b>		
<b>Session 3</b>	<b><i>Biodiversity &amp; ecosystem processes</i></b>	<b><i>Moderators: Arthur Gessler and Tine Grebenc</i></b>
<b>09:00-09:30</b>	<b>Arthur GESSLER (KN9)</b>	<b>The impact of environmental factors on BGC cycles...</b>
<b>09:30-10:00</b>	<b>Zachary KAYLER (KN10)</b>	<b>Drought and heat stress effects on the plant-soil carbon ...</b>
<b>10:00-10:30</b>	<b>Giorgio MATTEUCCI (KN11)</b>	<b>Responding to challenges of forest ecosystem research ...</b>
<b>10:30-11:00</b>	<b><i>Coffee and tea</i></b>	
<b>11:00-11:30</b>	<b>Petr BALDRIAN (KN12)</b>	<b>The effect of forest trees on microbial processes ... in C ...</b>
<b>11:30-12:00</b>	<b>Hans GÖRANSSON (KN13)</b>	<b>Tree biodiversity and ecosystem function</b>
<b>12:00-12:30</b>	<b>Heljä-Sisko HELMISAARI (KN14)</b>	<b>Fine root traits and soil processes in boreal forests</b>
<b>12:30-12:45</b>	<b>Jaana LEPPÄLAMMI-KUJANSUU</b>	<b>Carbon input into soil originating from FR and foliage litter</b>
<b>12:45-13:00</b>	<b>Naoki MAKITA</b>	<b>Fine root physiological and morphological traits of ...</b>
<b>13:00-14:00</b>	<b><i>Lunch</i></b>	
<b>14:00-22:00</b>	<b><i>Field excursion and local tourism</i></b>	

## Scientific programme

<b>Thursday September 3<sup>rd</sup></b>		
<b>Session 4</b>	<b><i>Forest disturbance, management &amp; modelling</i></b>	<b><i>Moderators: Michael Starr and Urša Vilhar</i></b>
09:00-09:30	John DIGHTON (KN15)	Disturbance and resilience in forests...
09:30-10:00	Cristina CRUZ (KN16)	Impacts of increased nitrogen availability...
10:00-10:15	Ivan KREFT	Forest-agriculture interface: a case of buckwheat
10:15-10:30	Maša KOVIČ DINE	Protection of forests under international law
10:30-11:00	<i>Coffee and tea</i>	
11:00-11:30	Michael STARR (KN17)	Impact of forest harvesting on base cation budgets ...
11:30-11:45	Joanna HOREMANS	Variance decomposition of stem biomass increment ...
11:45-12:00	Joana VIEIRA	Adjustment capability of Maritime Pine cambial activity in ...
12:00-12:15	Iftekhar U. AHMED	Above and belowground tree biomass allocations ...
12:15-12:45	Hrvoje MARJANOVIĆ (KN18)	Carbon cycling in lowland oak ecosystems
12:45-14:00	<i>Lunch</i>	
<b>Session 5</b>	<b><i>Future forest research and innovation</i></b>	<b><i>Moderators: Reinhart Ceulemans and Hojka Kraigher</i></b>
14:00-14:30	Nate McDOWELL (KN19)	Consistent predictions of future forest mortality
14:30-15:00	Rainer MATYSSEK (KN20)	Ozone research, <i>QUO VADIS?</i> ...
15:00-15:15	Discussion	
15:15-15:45	<i>Coffee and tea</i>	
	<b><i>EUFORINNO impacts</i></b>	<b><i>Moderators: Reinhart Ceulemans and Hojka Kraigher</i></b>
15:45-16:00	Hojka KRAIGHER & WP leaders	Overview of EUFORINNO project & deliverables
16:00-16:15	Marko BAJC	Modern laboratories and standardization of protocols
16:15-16:30	Daniel ŽLINDRA	Towards accreditation
16:30-16:45	Tanja MRAK	Progress of microscopy facilities within EUFORINNO project
16:45-17:00	Saša ZAVADLAV	Use of stable isotopes in forest ecosystem: on the progress of SIL at the SFI
17:00-17:15	Mitja FERLAN	Measuring isotopic composition of carbon dioxide in the air
17:15-17:25	Maja PETEH	The publishing center <i>Silva Slovenica</i> and monograph series <i>Studia Forestalia Slovenica</i>
17:25-17:35	Tine GREBENC	The scientific journal <i>Acta Silvae et Ligni</i>
17:35-17:45	Maja PETEH	<i>SciVie - Sciences de la vie</i> - from the past and for the future
17:45-18:00	Ralitsa ATANASOVA	EUFORINNO pre-evaluation
19:30-22:00	<i>Conference dinner and awards ceremony</i>	

## Scientific programme

Friday, September 4 <sup>th</sup>		
<b>Round table</b>	<b>Forest science and innovation in Europe</b>	<b>Moderators: Reinhart Ceulemans and Primož Simončič</b>
<b>09:00-11:00</b>	<p><b>Hojka KRAIGHER</b>, SFI, Head of Programme group P4-0107 "Forest Biology, Ecology and Technology", Scientific Board for Biotechnical Sciences, SRA, member of LEGS Committee, Science Europe</p> <p><b>Tadej BAJD</b>, President, Slovenian Academy of Sciences and Arts</p> <p><b>Miha HUMAR</b>, Vice-Dean, Biotechnical Faculty, University of Ljubljana, member of the Scientific Board for Biotechnical Sciences, SRA</p> <p><b>Rainer MATYSSEK</b>, Chair for Ecophysiology of Plants, Center of Life and Food Sciences Weihenstephan, Department of Ecology, TUM</p> <p><b>Primož SIMONČIČ</b>, Director, SFI, Slovenian representative in LULUCF in 2007</p> <p><b>Representatives</b> of the ministries, responsible for forestry, science, technology, environment, Slovenian Research Agency, Biotechnical Faculty – Uni. Ljubljana, Slovenia Forest Service, and of international organizations</p>	<p><b>Contributions (5 minutes each):</b></p> <ul style="list-style-type: none"> <li>Overview of forestry scopes and challenges</li> <li>Forestry and wood science in Slovenian evaluation schemes</li> <li>Forest research and monitoring and its financing in Slovenia</li> <li>Climate change and research infrastructures in forestry</li> <li>The Slovenian reality and prosperity of science</li> <li>Smart specialization and natural resources</li> <li>The global perspectives in forest science and innovation</li> <li>The Slovenian and international horizons</li> <li>Needs &amp; expectations from forests, forestry &amp; wood industry</li> </ul> <p><b>Discussion (1 hour)</b></p>
<b>11:00-12:00</b>	<b>Press conference</b>	
<b>12:00-12:30</b>	<b>Closure of the conference</b>	
<b>12:30-14:00</b>	<b>Lunch</b>	

## ADAPTIVE RESPONSES OF TREES TO CLIMATE CHANGE

*Antoine Kremer*<sup>1</sup>

<sup>1</sup> INRA, Functional ecology and genomics UMR 1202 Biodiversité Gènes & Communautés INRA, 69 route d'Arcachon, 33612 CESTAS Cedex, France

E-mail: antoine.kremer@pierroton.inra.fr

**Keywords:** paleobotany, adaptation, migration, hybridization, oaks

Adaptation of long lived tree species to ongoing rapid climate change is a pivotal question in ecology and forest management. There are indeed concerns that trees may not be able to cope with future environmental change, due to their long generation time. However, there is a growing body of evidence stemming from different sources of information that were recently documented (Quaternary evolutionary history; observations from population and species transfers; provenance experiments) that trees may have resources and mechanisms to cope with climate change. I will review the genetic and ecological mechanisms that have facilitated adaptation of trees during historical “natural” warming periods taking as an example European oaks. Assembling lessons from phylogeography, paleobotany and simulations, I will show how oaks have responded quite rapidly to environmental change, despite their low evolutionary rate at the gene level. I will further examine how these mechanisms may be stimulated during the ongoing climatic changes.

### **Acknowledgements**

EU projects TREEPEACE, MOTIVE and FORGER

## GENETIC MONITORING IN FORESTS – EFFORTS FOR A EUROPEAN IMPLEMENTATION

*Monika Konnert<sup>1</sup>, Barbara Fussi<sup>1</sup>, Marjana Westergren<sup>2</sup>, Filipos Aravanopoulos<sup>3</sup>, Hojka Kraigher<sup>2</sup>*

<sup>1</sup> Bayerisches Amt für forstliche Saat- und Pflanzenzucht, Forstamtsplatz 1, 83317 Teisendorf, Germany

<sup>2</sup> Slovenian Forestry Institute, Večna pot 2, 1000 Ljubljana, Slovenia

<sup>3</sup> Aristotele University of Thessaloniki, University Campus, GR54124 Greece

E-mail: monika.konnert@asp.bayern.de

**Keywords:** genetic monitoring, indicators, verifiers, LIFEENMON

Sustainable forest management is based on a long-term adaptability of forest ecosystems and starts at the lowest, the gene level. Forest genetic monitoring (FGM) implies the long-term observation of the status and the temporal developments (changes) in the genetic system of forest tree populations on the basis of criteria, indicators and verifiers. As genetic variation is the basis of adaptability, forest genetic monitoring is a crucial component of any sustainable forest management because it detects changes of forest adaptability before they are seen on higher levels. At the same time genetic monitoring is assumed to contribute essentially to the estimation and evaluation of the effect of influences on the genetic system in forests, thus making it an early warning and controlling system for ecosystem changes. In the present contribution suitable criteria, indicators and verifiers for a FGM system are presented and the objectives of genetic monitoring are described. The recently started European LIFE project (LIFEENMON) aims to define optimal indicators and verifiers for monitoring of genetic diversity changes in time across a transect from Bavaria to Greece for two target species, *Abies alba* and *Fagus sylvatica*. In the same time it intends to prepare guidelines for forest genetic monitoring for these two and additional five forest tree species at a national, regional and European scale, to elaborate a manual for implementation of FGM and to disseminate knowledge on the necessity for conservation of forest genetic resources in general and observation of changes through genetic monitoring in detail. The project objectives and the expected results will be presented

Konnert M, Maurer W, Degen B, Kätzel R (2011): Genetic monitoring in forests – early warning and controlling system for ecosystemic changes. *iForest* 4: 77-81.

Graudal L, Aravanopoulos F, Bennadji Z, Changtragoon S, Fady B, Kjær DE, Loo J, Ramamonjisoa L, Vendramin GG (2014): Global to local genetic diversity indicators of evolutionary potential in tree species within and outside forests. *Forest Ecology and Management* <http://dx.doi.org/10.1016/j.foreco.2014.05.002>.

## ENDEMIC FOREST TREE SPECIES IN SOUTHEAST EUROPE, THEIR GENETIC RESOURCES AND USE

Alexander H. Alexandrov<sup>1</sup>

<sup>1</sup> Forest Research Institute, 132 Kliment Ohridski Blvd., 1756 Sofia, Bulgaria

E-mail: alexandrov\_38@abv.bg

**Keywords:** endemism, relict, in situ and ex situ conservation

The biggest biodiversity in the Old continent is established in the Iberian and the Balkan peninsulas due to wide variety of relief and climate. One of the indicators for biodiversity resources is the presence of endemic species. The endemic rate of Southeast European flora reaches up to 8-9 %, while in West, Central and North Europe it hardly gets near 1-2 % or even none. Most important endemic forest tree species from ecological and economic point of view in Southeast Europe are: Macedonian pine (*Pinus peuce* Grisb.), Bosnian pine (*Pinus heldreichii* Christ.), Serbian spruce (*Pinus omorica* (Panc.) Purc.), Greek fir (*Abies cephalonica* Loud.), Bulgarian fir (*Abies borisii regis* Mattf.), Balkan maple (*Acer heldreichii* Orph. ex Boiss.), Horse chestnut (*Aesculus hippocastanum* L.) and Rila oak (*Quercus protoroburoides* Don. et Bouz.). The valuable timber of Macedonian pine, ornamental characteristics of Serbian spruce and horse chestnut, as well as anti-erosion properties of Bosnian and Macedonian pine, make them very useful forest tree species. Most of these endemics including relicts, i.e. paleoendemics, require additional measures and concerns for their protection. The conservation of forest genetic resources of endemic species is performed by both in situ and ex situ methods.

## DETERMINATION OF GENETIC VARIATION EXISTING IN ANATOLIAN BLACK PINE NATURAL STAND AND PLANTATION AREA IN KASATURA BAY

*Gaye Kandemir<sup>1</sup>, Burcu Çengel<sup>1</sup>, Yasemin Tayanç<sup>1</sup>, Ercan Veliöglu<sup>2</sup>, Zeki Kaya<sup>3</sup>*

<sup>1</sup> General Directorate of Forestry, Forest Tree Seeds and Tree Breeding Research Institute Directorate, Ankara, Turkey

<sup>2</sup> General Directorate of Forestry, Poplar and Fast Growing Forst TRees Research Institute, Izmit, Turkey

<sup>3</sup> Middle East Technical University, Ankara, Turkey

E-mail: gayekandemir@ogm.gov.tr

**Keywords:** nature conservation area, Black Pine (*Pinus nigra*), genetic variation, RAPD, isozyme

The aim of this study is to determine the magnitude and pattern of genetic variation existing in Kasatura Bay Nature Conservation Area Anatolian black pine (*Pinus nigra* Arnold) natural stand (designated as a Nature Conservation Area) and plantation area nearby. In addition, to estimate whether there is genetic contamination from plantation to natural stand. Cones were collected from 97 mother trees (39 trees from natural stand and 58 trees from plantation) to obtain seeds which are produced by half-sib families. Genetic diversity parameters were estimated for both populations by using RAPD (Randomly Amplified Polymorphic DNA) markers. For detection of pollen contamination, isozymes were used by using 30 trees from each population. Eight RAPD primers generated 74 polymorphic loci. Genetic diversity parameters of the studied populations were generally similar. Percentage of polymorphic loci was 99 % for natural stand and 96 % for plantation. Mean number of observed alleles was 1.98 in natural stand and 1.96 in plantation. The mean number of effective alleles was about 1.62 in both populations. Observed heterozygosity values were estimated to be 0.42 and 0.39 for plantation and natural stand, respectively. Negative FIT (-0.11) and FIS (-0.15) values implied excess heterozygosity in the studied populations. Mean FST value (0.04) indicates that there was little differentiation between natural stand and plantation and vast majority (96 %) of genetic diversity was contained within populations. Studied 5 enzyme systems revealed 11 isozyme loci and 20 polymorphic alleles. Genetic (pollen) contamination was found to be 80.4 % in natural population.

## FOREST GENETIC MONITORING – DELINEATION OF MONITORING REGIONS ON A TRANSECT FROM BAVARIA TO GREECE

*Marjana Westergren<sup>1</sup>, Barbara Fussi<sup>2</sup>, Filippos Aravanopoulos<sup>3</sup>, Evangelia V. Avramidou<sup>3</sup>, Roland Baier<sup>2</sup>, Gregor Božič<sup>1</sup>, Ioannis V. Ganopoulos<sup>3</sup>, Darius Kavaliuskas<sup>2</sup>, Monika Konnert<sup>2</sup>, Ermioni S. Malliarou<sup>3</sup>, Vladko Andonovski<sup>4</sup>, Dalibor Ballian<sup>5</sup>, Davorin Kajba<sup>6</sup>, Heino Konrad<sup>7</sup>, Saša Orlović<sup>8</sup>, Fotis Kiourtsis<sup>9</sup>, Živan Veselić<sup>10</sup>, Hojka Kraigher<sup>1</sup>*

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI-1000 Ljubljana, Slovenia

<sup>2</sup> Bavarian Office for Forest seeding and planting, Forstamtsplatz 1, DE-83317 Teisendorf, Germany

<sup>3</sup> Aristotle University of Thessaloniki, University Campus, GR-54124 Thessaloniki, Greece

<sup>4</sup> University Ss. Cyril and Methodius, Faculty of Forestry, MK-1000 Skopje, Republic of Macedonia

<sup>5</sup> University of Sarajevo, Faculty of Forestry, Zagrebačka 20, BA-71000 Sarajevo, Bosnia and Herzegovina

<sup>6</sup> University of Zagreb, Faculty of Forestry, Svetošimunska 25, HR-10000 Zagreb, Croatia

<sup>7</sup> Federal Research and Training Centre for Forests, Natural Hazards and Landscape, Hauptstraße 7, A-1140 Vienna, Austria

<sup>8</sup> Institute of Lowland Forestry and Environment, Antona Čehova 13, RS-21000 Novi Sad, Republic of Serbia

<sup>9</sup> The Decentralized Administration of Macedonia – Thrace, GR-54008 Thessaloniki, Greece

<sup>10</sup> Slovenia Forest service, Večna pot 2, SI-10000 Ljubljana, Slovenia

E-mail: marjana.westergren@gozdis.si

**Keywords:** forest genetic monitoring, monitoring regions, forest genetic resources, LIFEGENMON, LIFE

Conservation and management of forest genetic resources for sustainable use is a crucial, but not an easy task and special tools, such as forest genetic monitoring are needed to recognise the state and changes in their composition in a timely manner. The six-year LIFE + implementation project LIFEGENMON, led by the Slovenian Forestry Institute, and supported by six partners from Germany, Greece and Slovenia, is intended to design, test and implement forest genetic monitoring on the transect from Germany to Greece. However, one of the elementary requirements needed for implementation of forest genetic monitoring is delineation of monitoring regions. Within the project, this has been done for seven tree species or species complexes differing in their biology and distribution (*Fagus sylvatica*, *Abies alba* / *A. borisii regis* complex, *Fraxinus excelsior*, *Populus nigra*, *Pinus nigra*, *Prunus avium*, *Quercus robur* / *Q. petraea* complex). Criteria for selecting and delineating monitoring regions were: (i) equal coverage of environmental zones, (ii) coverage of distinguished races or ecotypes as well as inclusion of marginal and peripheral populations at the (latitudinal and altitudinal) leading and rear edge of the species distribution range, (iii) each region should preferably include already defined dynamic gene conservation units (EUFGIS), (iv) known levels of standing genetic variation, (v) equal coverage of standing genetic structure and recolonisation routes, (vi) local expert knowledge regarding forest types, vitality and value (biodiversity, economic) of populations. Six to nine monitoring regions per species / species complex were recognised.

### Acknowledgements

This work was financially supported by the European Union's LIFE financial mechanism (LIFEGENMON project, LIFE13 ENV/SI/000148) and national co-financers, in Slovenia MKGP, MOP, SFI, in Germany StMELF, and in Greece the Green Fund.

## INTER AND INTRA-POPULATION VARIATION OF LEAF STOMATAL TRAITS OF *Populus nigra* L. IN VOJVODINA, NORTHERN SERBIA

*Dijana Čortan*<sup>1</sup>, *Dragica Vilotić*<sup>2</sup>, *Mirjana Šijačić-Nikolić*<sup>2</sup>

<sup>1</sup> Faculty of Education, University of Novi Sad, Podgorička 4, 25000 Sombor, Serbia

<sup>2</sup> Forest Faculty, University of Belgrade, Kneza Višeslava 1, 11000 Belgrade, Serbia

E-mail: dijanacortan@yahoo.com

**Keywords:** *Populus nigra* L., stomatal traits, Vojvodina, interpopulation and intrapopulation variability

Black poplar (*Populus nigra* L.) as one of the most important riparian species is currently threatened by extinction across whole Europe. Since the middle of last century it has been faced with severe threats pertaining to its survival, mostly because of the serious loss of its natural habitat and the lack of natural regeneration.

The research was conducted in four *Populus nigra* L. natural populations located in the basin of three river valleys of Vojvodina region, northern Serbia. The research involved the examination of inter- and intra-population variation of stomatal traits: stomatal density, ratio adaxial / abaxial stomatal densities, stomatal dimensions (length and width of stomatal guard cells and stomatal aperture), potential conductance index and stomatal shape coefficient, in *Populus nigra* L. leaves. The stomatal characteristics were examined in fully expanded leaves, from two leaf positions - the sun-exposed and shaded side of the tree, and from both sides of leaf considering fact that this species is amphistomatous.

Results show that within and between studied populations exists considerable variability, with the variability much more pronounced within than between populations. According to CDA analysis traits that most contribute to the differences between these populations are: adaxial stomatal width, abaxial stomatal aperture width and shape coefficient of abaxial stomatal aperture.

## BIODIVERSITY-STABILITY RELATIONSHIPS UNDER CLIMATE EXTREMES

*Hans De Boeck<sup>1</sup>, Juliette Bloor<sup>2</sup>, Michaela Zeiter<sup>3</sup>, Jürgen Kreyling<sup>4</sup>, Johannes Ransijn<sup>5</sup>, Ivan Nijs<sup>1</sup>, Anke Jentsch<sup>5</sup>*

<sup>1</sup> Centre of Excellence PLECO (Plant and Vegetation Ecology), Universiteit Antwerpen, Universiteitsplein 1, B-2610 Wilrijk, Belgium

<sup>2</sup> INRA, UR 874, Unité de Recherche sur l'Ecosystème Prairial, 5 chemin de Beaulieu, F-63100 Clermont-Ferrand, France

<sup>3</sup> School of Agricultural, Forest and Food Sciences, Bern University of Applied Sciences, Länggasse 85, CH-3052 Zollikofen, Switzerland

<sup>4</sup> Experimental Plant Ecology, Institute of Botany and Landscape Ecology, Greifswald University, Soldmannstrasse 15, D-17487 Greifswald, Germany

<sup>5</sup> Disturbance Ecology, Bayreuth Center of Ecology and Environmental Research, University of Bayreuth 95440 Bayreuth, Germany

E-mail: hans.deboeck@uantwerp.be

Theoretical and empirical studies both suggest that high species diversity within ecosystems tends to increase plant community stability because the sensitivity to fluctuations in environmental conditions varies so that more diverse communities have a wider range of sensitivities. The ultimate result is that community functioning tends to be more stable under a range of conditions. While most evidence gathered to support this theory originates from studies focusing on year-to-year stability and moderate fluctuations in environmental conditions, we assessed whether the biodiversity-stability concept applies also to more extreme, sudden events such as severe droughts and heat waves. In literature, we found that studies simultaneously considering climate extremes and species richness differed in their conclusions. Some studies were supportive of the diversity-stability hypothesis, while others came to opposite conclusions or found no or inconclusive effects. We propose several reasons why the biodiversity-stability concept may not always apply under extreme event conditions. This relates to ambiguities in what exactly constitutes an “extreme” and how stability is measured, but also to community traits changing with changes in biodiversity. We subsequently suggest ways to improve mechanistic understanding of biodiversity-stability effects in order to facilitate comparison between studies and avoid making the right predictions for the wrong reasons.

### Acknowledgements

The SIGNAL pan-European research project, funded by the BiodivERSA call of the ERA-Net and for Belgium specifically BELSPO (Belgian Science Policy).

## EUROPEAN WHITE BIRCH AND RAGWEED: IMPACT OF AIR POLLUTIONS ON THE ALLERGENIC POLLEN, AN OVERVIEW

Ulrike Frank<sup>1</sup>, Christine von Thörne<sup>1</sup>, Jeroen T. M. Buters<sup>2</sup>, Sebastian Öder<sup>2</sup>, Claudia Traidl-Hoffmann<sup>3</sup>, Dieter Ernst<sup>1</sup>

<sup>1</sup> Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany

<sup>2</sup> ZAUM – Center for Allergy and Environment, München, Germany

<sup>3</sup> UNIKA-T, Technische Universität München, Augsburg, Germany

E-mail: ernst@helmholtz-muenchen.de

**Keywords:** air pollution, allergenicity, birch, pollen, ragweed

Birch pollen is mainly responsible for allergic diseases in Central and Northern Europe, whereas ragweed pollen is the main elicitor for such diseases in Northern America and the weed has become a spreading neophyte in Europe. Climate change and air pollution will affect the allergenic potential of pollen, either by changes of the pollen season, the pollen amount, or by directly increasing the transcripts and allergenic proteins.

Catkins of birch were collected in the area of Munich. High environmental O<sub>3</sub> concentrations showed a positive correlation to the allergen Bet v 1 content and a negative correlation was observed for NO<sub>2</sub> [1]. Skin prick tests showed larger wheals indicating an enhanced allergenicity in an O<sub>3</sub>-enriched environment [1]. However, regional and year-to-year variations in Bet v 1 release were also observed [2]. Fumigation of birch pollen with O<sub>3</sub>/NO<sub>2</sub> led to a pronounced increase of protein nitration, which might increase the allergenicity [3].

Regarding ragweed transcriptomic analyses showed changes in allergen-encoding ESTs upon elevated O<sub>3</sub>/CO<sub>2</sub>/NO<sub>2</sub> and drought stress [4, 5]. Increased amounts of allergenic proteins were found upon CO<sub>2</sub>/NO<sub>2</sub>-treatment [6], in addition nitrosylation of Amb a isoforms were also observed. CO<sub>2</sub>-treated pollen or drought stress lead to higher IgE-levels in B cells of mice, and an increased pollen allergenicity was shown by immunoblotting with ragweed antisera.

### Acknowledgements

This work was supported by the grant 3/09 CK-CARE, Christine-Kühne Center for Allergy Research and Education.

### References

1. Beck et al (2013) *Plos One* 8:e80147.
2. Buters et al (2008) *Int Arch Allergy Imm* 145: 122-130.
3. Franze et al (2005) *Environ Sci Technol* 39: 1673-1678.
4. Kanter et al (2013) *Plos One* 8: e61518.
5. El-Kelish et al (2014) *BMC Plant Biol* 14: 176.
6. Frank et al (2014) *Ambrosie* 29: 32-43.

## BIO-ENERGY FROM BIOMASS: FULL GREENHOUSE GAS BALANCE, ENERGY BALANCE AND ENVIRONMENTAL LIFE CYCLE ASSESSMENT

*Reinhart Ceulemans<sup>1</sup> and POPFULL team (N. Arriga, G. Berhongaray, L. S. Broeckx, T. De Grootte, O. El Kasmoui, C. Görres, S. Njakou Djomo J. Segers, S. Vanbeveren, M. S. Verlinden and T. Zenone)<sup>1</sup>*

<sup>1</sup> University of Antwerp, Research Centre of Excellence on Plant & Vegetation Ecology, Department of Biology, Universiteitsplein 1, BE-2610 Wilrijk, Belgium

E-mail: reinhart.ceulemans@uantwerp.be

**Keywords:** POPFULL, yield, ecological balance, economic assessment, energy efficiency

With regard to efficacy of bioenergy as a viable renewable energy source three questions need to be answered. Q1: Is bioenergy from woody biomass energy efficient? Q2: Does bioenergy present a favourable greenhouse gas balance? Q3: Is bioenergy from woody biomass economically profitable? Answers to these three questions are provided in this lecture based on the first complete cradle-to-grave analysis of a renewable bioenergy source. Over five years (2009-2014) an operational plantation of fast-growing poplar (*Populus*) trees was managed as a short-rotation coppice in two-year rotation cycles. After each rotation the plantation was harvested and the coppice left to regrow as a multi-stem culture. The plantation was intensively monitored from establishment through to the production of bioenergy in combined heat-and-power plants. In answer to Q1 the bioenergy production was highly energy efficient, yielding nine times more energy than was put in. To answer Q2, the net fluxes of the five principal greenhouse gases between the plantation and the atmosphere were continuously monitored from a meteorological mast in the field. The plantation was a net carbon sink, i.e., absorbed more carbon from the atmosphere than was produced. But the non-CO<sub>2</sub> greenhouse gases (methane and nitrous oxide) represented an overall net emission to the atmosphere. In answer to Q3 the bioenergy culture was not economically viable. The total cost of producing bioenergy was five times higher than the benefit of the renewable energy produced. The unique results of POPFULL were presented by National Geographic (<http://greenenergycleanenergy.com/behind-the-science-renewable-energy-national-geographic>).

### Acknowledgements

Funding from the European Research Council under the EC Seventh Framework Program as Advanced Grant n° 233366 (POPFULL), Flemish Hercules Foundation (contract ZW09-06), and the Flemish Methusalem Program.

## QUANTIFICATION OF LINKED CARBON AND WATER CYCLES IN A LOBLOLLY PINE-SWITCH GRASS INTERCROPPING SYSTEM FOR BIOENERGY AND WOOD PRODUCTS

*J. S. King<sup>1</sup>, M. Fischer<sup>1,2</sup>, J-C. Domec<sup>1,3</sup>, J. Williamson<sup>1</sup>, E. Sucre<sup>4</sup>*

<sup>1</sup> Department of Forestry and Environmental Resources, North Carolina State University, Raleigh, NC, USA,

<sup>2</sup> Department of Agriculture Systems and Bioclimatology, Mendel University, Brno, Czech Republic

<sup>3</sup> Bordeaux Sciences Agro UMR INRA-TCEM, University of Bordeaux, Bordeaux, France

<sup>4</sup> Southern Timberlands R&D, Weyerhaeuser NR Company, Vanceboro, NC, USA

E-mail: john\_king@ncsu.edu

**Keywords:** bioenergy, climate change, forest intercropping, US Southeast, sustainability

Dependence on fossil fuel energy exposes the U.S. to potentially catastrophic climate change, economic instability, and risks to national security. The goal of the project is to assess sustainability of a novel pine-switch grass intercropping system for the production of bioenergy and traditional wood products based on regionally appropriate crops and indigenous biomass production practices. We hypothesize the asynchronous physiology and growth of the C3 trees and C4 grass along with the different soil horizons exploited by the respective root systems will allow for greater nutrient retention and more efficient utilization of site water. We also hypothesize that the additional soil volume exploited by switchgrass roots and associated fine root turnover will increase soil organic C, thus increasing C sequestration. The work is being done at a 32 ha replicated field experiment installed in the Lower Coastal Plain of North Carolina being implemented by Weyerhaeuser NR Company. The experiment consists of a randomized complete block design of traditional pine silviculture and switch grass intercropping treatments, deployed individually and together, in 0.8 ha plots sufficient for ecosystem analysis. Over the course of the five-year project we are carefully monitoring ecosystem productivity, soil carbon and nutrient dynamics, and water utilization. Measurements are being used to parameterize ecosystem models to scale results to the region and assess the effects of a changing climate.

## SUCCESSION INDUCED CHANGES IN ECOSYSTEM FUNCTIONING WITH EMPHASIS ON CARBON AND WATER EXCHANGE

*Klemen Eler<sup>1,2</sup>, Mitja Ferlan<sup>2</sup>, Giorgio Alberti<sup>3</sup>, Alessandro Peresotti<sup>3</sup>, Primož Simončič<sup>2</sup>, Dominik Vodnik<sup>1</sup>*

<sup>1</sup> University of Ljubljana, Biotechnical Faculty, Jamnikarjeva 101, Ljubljana, Slovenia

<sup>2</sup> Slovenian Forestry Institute, Večna pot 2, Ljubljana, Slovenia

<sup>3</sup> Department of Agricultural and Environmental Sciences, University of Udine, Udine, Italy

E-mail: klemen.eler@bf.uni-lj.si

**Keywords:** drought effects, secondary forest, calcareous grassland, eddy-covariance, submediterranean Slovenia

Many marginal agricultural areas in Europe and elsewhere are being abandoned and subjected to spontaneous afforestation with shrub and tree species. Within ecosystem research this process is rarely studied (but see Stoy et al. 2008). Here, the functioning in terms of carbon and water vapour exchanges of a secondary forest developed on a former semi-dry calcareous grassland in the sub-Mediterranean region of Slovenia was investigated and compared with the nearby pasture still in use. Eddy covariance and other ancillary data was used on both plots to estimate fluxes of CO<sub>2</sub> and H<sub>2</sub>O. Yearly and seasonal net ecosystem exchange and evapotranspiration of both ecosystems were examined together with their environmental controls (soil water content, air temperature, light availability, air humidity) during the period of 6 years. There were significant differences in yearly cumulatives, seasonal phenology of C assimilation and drought resistance between the investigated ecosystems. The occurrence of drought seemed to largely govern ecosystem functioning and productivity. By identifying the drought periods using the evapotranspiration response to soil water availability it was shown that more than the severity of drought the duration of the longest yearly drought affected the estimated yearly gross primary productivity. The functioning of ecosystems was also compared with respect to ecosystem-level water use efficiency, rain use efficiency and light use efficiency in different environmental conditions and phenological phases. The uncertainty in derived fluxes will also be discussed particularly with respect to spatial heterogeneity of the investigated ecosystems and possible abiotic (geological) sources of CO<sub>2</sub> stemming from carbonate dissolution / precipitation and subterranean storage in karst cavities (Kowalski et al. 2008; Serrano-Ortiz et al. 2010).

### Acknowledgements

This work was supported by grants from the Slovenian Research Agency and the Ministry of Agriculture and Environment of Slovenia (research projects J4-1009, V4-0536, J4-4224, and L4-4318 and research programs P4-0085 and P4-0107). Part of the work was performed within the EUFORINNO project (RegPot No. 315982).

Kowalski AS, Serrano-Ortiz P, Janssens IA, Sánchez-Moral S, Cuezva S, Domingo F, Were A, Alados-Arboledas L (2008) Can flux tower research neglect geochemical CO<sub>2</sub> exchange? *Agric. For. Meteorol.* 148: 1045–1054.

Serrano-Ortiz P, Roland M, Sanchez-Moral S, Janssens IA, Domingo F, Goddérés Y, Kowalski AS (2010) Hidden, abiotic CO<sub>2</sub> flows and gaseous reservoirs in the terrestrial carbon cycle: Review and perspectives. *Agric. For. Meteorol.* 150: 321–329.

Stoy PC, Katul GG, Siqueira MBS, Juang J-Y, Novick KA, McCARTHY HR, Oishi AC, Oren R (2008) Role of vegetation in determining carbon sequestration along ecological succession in the southeastern United States. *Glob. Chang. Biol.* 14: 1409–1427.

## EFFECTS OF WARMING ON *Pinus sylvestris* IBERIAN FORESTS

Emilia Gutiérrez<sup>1</sup>

<sup>1</sup> Dept. of Ecology, Faculty of Biology, University of Barcelona, Av. Diagonal 643, 08028 Barcelona, Spain

E-mail: emgutierrez@ub.edu

**Keywords:** Scots pine forests, Iberian Peninsula, warming, growth trends

Scots pine (*Pinus sylvestris* L.) is the conifer tree species with the widest distribution in the world reaching its southern distribution limit in Spain; it has a wide niche breadth, forming forests across a variety of climates and soil types in Europe and Asia. The Spanish Iberian Scots pine forests occupy a discontinuous area of 700,000 ha (Alía et al. 2001) and Studies on Scots pine responses to climate change (warming, climate variability and drought) have become abundant in the last 2 decades in Spain. Warming and drought have been causing an increased physiological stress (Poyatos et al. 2013) and tree growth decline and occasionally tree death in those more prone areas to drought stress (Martínez-Vilalta and Piñol 2002, Galiano et al. 2010, Heres et al. 2012). (e.g. Andreu-Hayles et al. 2011, Sanchez-Salguero et al. 2015). In this study, I review the main results of Scots pine responses to global climate change in Spain, and second, I will present results on Scots pine growth trends and distribution of resources at different temporal scales from locations with contrasting climate conditions.

Andreu-Hayles L, Planells O, Gutiérrez E, Muntan E, Helle G, Anchukaitis KJ, Schleser GH (2011) Long tree-ring chronologies reveal 20th century increases in water-use efficiency but no enhancement of tree growth at five Iberian pine forests. *Global Change Biology* 17: 2095-2112.

Alía R, Moro-Serrano J, Notivol E (2001) Genetic variability of Scots pine (*Pinus sylvestris*) provenances in Spain: growth traits and survival. *Silva Fennica* 35(1): 27-38.

Galiano L, Martínez-Vilalta J, Lloret F (2010) Drought-induced multifactor decline of Scots pine in the Pyrenees and potential vegetation change by the expansion of co-occurring oak species. *Ecosystems* 13: 978-991.

Heres AM, Martínez-Vilalta J, López BC (2012) Growth patterns in relation to drought-induced mortality at two Scots pine (*Pinus sylvestris* L.) sites in NE Iberian Peninsula. *Trees* 26(2): 621-630.

Martínez-Vilalta J, Piñol J (2002) Drought-induced mortality and hydraulic architecture in pine populations of the NE Iberian Peninsula. *For Ecol Man* 161: 247-256.

## STABLE ISOTOPES IN TREE RINGS

*Danny McCarroll*<sup>1</sup>

<sup>1</sup> Department of Geography, Swansea University, Singleton Park, Swansea UK

E-mail: d.mccarroll@swansea.ac.uk

**Keywords:** carbon, oxygen, climate change, dendrochronology

Tree rings provide perhaps the best natural archive of information on past climate change. Many trees produce a ring every year, so careful cross dating provides perfect annual resolution with no uncertainty. However, the measures of tree growth that are used as proxies for past climate, such as ring widths and densities, only respond strongly to climate when trees are growing under stress. The best records are obtained from high latitude or high altitudes or from places that are very dry. The result is that there is a strong spatial bias in tree-ring based climate reconstructions towards places where very few people live. Stable isotope ratios in tree rings are not measures of net tree growth but act rather as passive indicators of changing climate. They can thus provide reliable palaeoclimate proxies for trees that are not growing under strong climatic or environmental stress. I will present results of carbon and oxygen isotope analysis from British oak trees that are not growing under stress. The carbon isotope ratios record the balance between stomatal conductance and photosynthetic rate. In Britain the dominant control is photosynthetic rate and the isotope ratios correlate very strongly with summer sunshine and summer temperature. The oxygen isotope ratios record both the isotopic ratios of summer precipitation and evaporative enrichment. However, in Britain it seems that the isotopic ratios of summer rainfall are strongly dominant, providing a record of past changes in summer circulation and therefore of the amount of summer rainfall. Calibration and verification results suggest that these palaeoclimate proxies are very strong and temporally stable and that they record positive and negative extremes with equal skill. There would seem to be great potential for using stable isotope ratios in tree rings to redress the geographical bias in available palaeoclimate proxies.

## WHAT INFORMATION IS STORED IN PHLOEM STRUCTURE AND RELATIONSHIP BETWEEN PHLOEM AND XYLEM INCREMENTS?

*Jožica Gričar*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Dept. of Yield and Silviculture, Večna pot 2, SI-1000 Ljubljana, Slovenia

E-mail: jozica.gricar@gozdis.si

Structure and formation of phloem have generally earned less attention than the xylem. This could be partly explained by relatively low economic value of phloem. However, unlike xylem formation, annual formation of phloem is crucial for tree survival as it maintains the translocation pathways for photosynthates and signalling molecules.

Seasonal dynamics of xylem and phloem formation are not synchronous processes thus capturing different environmental information. Xylem formation has already proven to be influenced by environmental factors, whereas phloem formation is more stable and presumably less affected by changes in growing conditions. In addition to secondary changes in older phloem, this might be among the main reasons for excluding phloem from dendroecological studies so far.

This presentation seeks to address the following questions: Which signals influence the structure of phloem? Do phloem structure and the relationship between phloem and xylem increments have potential to be used as sensitive stress indicators? Would information on phloem contribute to a more thorough understanding of radial growth of trees and their response to environmental changes? Examples of phloem in different tree species will be shown to discuss these questions.

### **Acknowledgements**

The work was supported by the Slovenian Research Agency, program P4-0107 and project Z4-9662 and by EUFORINNO (RegPot No. 315982) of the FP7 Infrastructures programme.

## STAND DYNAMICS AND DISTURBANCE HISTORY IN MIXED FOREST OF NORWAY SPRUCE AND CEMBRA PINE FROM EASTERN CARPATHIANS

*Popa Ione<sup>1</sup>, Nechita Constantin<sup>1</sup>, Sidor Cristian<sup>1</sup>*

<sup>1</sup> National Research and Development Institute for Silviculture, Calea Bucovinei 73 bis, Campulung Moldovenesc, Romania

E-mail: popaicas@gmail.com

**Keywords:** dendroecology, boundary line, Carpathians, disturbance

Old-growth forest structure is the results of the interaction between internal competition and external natural disturbance regime. Scale and frequency of the disturbance strongly affect forest structure and functions. This study presents the main results concerning the stand structure dynamics and the disturbance history in a mixed forest of *Picea abies* and *Pinus cembra* from Calimani National Park, Eastern Carpathians (Romania). Three permanent plots (surface between 0.7 and 2.2 ha) were established in an altitudinal transect. The disturbance dynamics and stand regeneration history highlights an important advance of spruce limits in the last 50 years in case of timberline plot. The second plot targeted a forest out in late optimal development phase where the actual spatial structure is the effect of multiple wind disturbances. The first wave of regeneration is observed on 1850 with a maximum on 1890, following a high intensity wind damage dated by dendroecological methods around 1840. The second regeneration wave, dominated by spruce, start after 1940 with maximum in 1960-1970 and the intensity is 2-3 times higher that precedent period. The last plot was established in a forest regenerated after catastrophic wind damage (dated in 1840). The regeneration history, analyzed using dendroecological methods, highlights the stand establishing starting with 1850 until 1910 with a maximum in 1880. This type of stand developments is specific to a regeneration of open-field and the following stand structure, on dimensional level, is conditioned by competition processes specific to forest ecosystem. Based on millennium tree ring chronology of cembra pine from Calimani Mts. was possible to reconstruct the disturbance dynamics in the last eight century from this region. A significant increase of frequency of disturbance is observed the last century and the largest one been that from 1840 when over 70 % from the trees included in the datasets show a significant growth release.

## ENVIRONMENTAL SIGNALS IN *Quercus robur* L. TREE-RING WIDTHS AND $\delta^{13}\text{C}$

*Polona Hafner*<sup>1</sup>, *Jožica Gričar*<sup>1</sup>, *Mitja Skudnik*<sup>1</sup>, *Tom Levanič*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana

E-mail: polona.hafner@gozdis.si

**Keywords:** Pedunculated oak, earlywood, latewood, dendrohydrology, dendroecology

We analysed two groups of *Quercus robur*, growing at nearby plots with different micro-location condition (W-wet and D-dry) in the floodplain Krakovo forest, Slovenia. In the study we compared growth response of two different tree groups to environmental variables, the potential signal stored in earlywood structure and the potential difference of the information stored in carbon discrimination of earlywood and latewood. For that purpose earlywood and latewood widths and carbon discrimination have been measured. We found out that W oaks were growing significantly better over the whole analysed period. The difference between D and W oaks was significant in all analysed variables with the exception of stable carbon discrimination in latewood. In W oaks, latewood widths correlated with summer (June to August) climatic variables, while carbon discrimination was more connected to Krka River flow during the summer. Earlywood discrimination correlated with summer and autumn River Krka flow of the previous year, while latewood discrimination correlated with flow during the current year. In the case of D oaks, environmental signal appears to be vague most probably due to less favourable growth conditions resulting in markedly reduced increments. Our study reveals important differences in responses to environmental factors between the two oak groups of different physiological conditions that are preconditioned by environmental stress. Environmental information stored in tree-ring features may vary, even within the same forest stand, and largely depends on the micro-environment. Our analysis confirmed our assumptions that separate EW and LW analysis of widths and carbon isotope discrimination provides complementary information in *Quercus robur* dendroecology.

### Acknowledgements

The work was supported by the Slovenian Research agency, young researchers' program (PH), program P4-0107, project L4-9653, and by EUFORINNO (RegPot No. 315982) of the FP7 Infrastructures programme. We acknowledge the help of Špela Jagodic and Robert Krajnc for help in the field and laboratory.

McCarroll D, Loader NJ (2004) Stable isotopes in tree rings. *Quaternary Sci Rev* 23: 771-801.

Gričar J, de Luis M, Hafner P, Levanič T (2013) Anatomical characteristics and hydrologic signals in tree-rings of oaks (*Quercus robur* L.). *Trees* 27: 1669–1680.

## CLIMATE SIGNAL AND POTENTIAL OF BOSNIAN PINE (*Pinus heldreichii* CHRIST) FOR CLIMATE RECONSTRUCTION IN CENTRAL W BALKAN REGION

*T. Levanič<sup>1</sup>, S. Poljanšek<sup>1</sup>, D. Ballian<sup>2</sup>*

<sup>1</sup> Department of Yield and Silviculture, Slovenian Forestry Institute, Slovenia

<sup>2</sup> Forestry faculty, University of Sarajevo, Bosnia and Herzegovina

E-mail: tom.levanic@gozdis.si

**Keywords:** long chronologies, Bosnia and Herzegovina, temperature, precipitation

Bosnian pine (*Pinus heldreichii* Christ) - PIHE, is a species native to mountainous areas of the Balkans and southern Italy. PIHE is a tree species of the upper tree line in the Dinaric mountains and as such a promising source of valuable climate information. Due to its longevity it might have a good potential for long climate reconstructions based solely on living trees. Aim of our study was to investigate potential of PIHE (1) for construction of long PIHE chronologies for the region of the central W Balkan and (2) for long climate reconstructions based mostly on cores from living trees.

Preliminary PIHE chronology is 437 (1573-present) years long and has a sufficient sample depth for EPS>0.85 from 1659 onwards. Comparison with climate data show specific response of PIHE to climate – trees respond positively to above average temperature in March (Mt) and negatively to above average temperature in June and July (JJt). We found weak positive response to June and July precipitation (JJp). This is somehow expected since precipitation are abundant along the Dinaric mountain ridge and are not factor in minima. Running window statistics reveal a significant temporal change in the identified signals – JJt and JJp signals were significant before 1965, and Mt after 1965, both temperature and precipitation loose signal after 1965.

Although already quiet long, PIHE chronology can be extended by adding additional old trees from the studied region. Climate signal in tree rings is somehow specific comparing to other trees species in the region (e.g. *Pinus nigra*). JJt and JJp seems to play an important role in the formation of tree rings (see also Seim et al. 2012), however a clear positive March temperature signal in PIHE tree-rings after 1965 and almost complete disappearance of the JJt and JJp signal after 1965 deserves attention as well. The reasons for that are yet to be investigated.

### Acknowledgment

Research reported in this work was supported by the Slovenian Research Agency under grant numbers P4-0107 - Programme and research group “Forest biology, ecology and technology” and J4-5519 basic research project “Paleoclimate data enhances drought prediction in the W Balkan regionf” as well as by the European Union 7<sup>th</sup> frame programme grant RegPot Nr.315982 - EUFORINNO.

Seim A, Büntgen U, Fonti P, Haska H, Herzig F, Tegel W, Trouet V, Treydte K (2012) Climate sensitivity of a millennium-long pine chronology from Albania. *Climate Research* 51 (3): 217-228. doi:10.3354/cr01076.

## **THE IMPACT OF ENVIRONMENTAL FACTORS ON BIOGEOCHEMICAL CYCLES IN TREES AND FOREST ECOSYSTEMS - FROM MOLECULAR MECHANISMS TO ECOSYSTEM FUNCTIONING**

*Arthur Gessler<sup>1</sup>*

<sup>1</sup> Swiss Federal Research Institute WSL, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland

E-mail: arthur.gessler@wsl.ch

Trees are large global stores of carbon (C) that will be impacted by increased carbon dioxide levels and climate change. However, at present we cannot properly predict the carbon balance of forests in future as we lack knowledge on how plant physiological processes, the transfer of carbon within the plant, carbon storage, and remobilization in the plant tissues as well as the release of carbon from the roots to the soil interact with environmental drivers and ecosystem-scale processes.

Moreover, precipitation variability will increase in future and drought and rainfall periods will alternate. As a consequence, the ability of roots to utilise intermittent rainfall events will be crucial for tree performance and survival.

This talk will summarise how stable isotope techniques can give new insights in the fate of newly assimilated C in plants and ecosystems on the one hand and water uptake and use on the other hand.

The talk will span from the molecular to the ecosystem scale and will also highlight how physiological information can be transferred on larger spatio-temporal scale levels in order to predict the effects of environmental drivers and biotic interactions under future climatic conditions.

## DROUGHT AND HEAT STRESS EFFECTS ON THE PLANT-SOIL CARBON CONTINUUM AND BELOWGROUND MICROBIAL COMMUNITIES

Isabell von Rein<sup>1</sup>, Arthur Gessler<sup>1,2</sup>, Katrin Premke<sup>1</sup>, Andreas Ulrich<sup>1</sup>, Zachary Kayler<sup>1</sup>

<sup>1</sup> Institute for Landscape Biogeochemistry, Leibniz Centre for Agricultural Landscape Research, Eberswalder Straße 84, D-15374 Müncheberg, Germany

<sup>2</sup> Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), CH-8903 Birmensdorf, Switzerland

E-mail: kayler@zalf.de

Important questions remain about the relevance of the plant-soil continuum in extreme climate event scenarios, such as, how might the plant-soil continuum benefit belowground systems? And, at which point is the connectivity severed? Furthermore, what do we stand to learn about microbial communities when pushed to their niche limits? We excavated soil monoliths, thus keeping the understory plant-microbe communities intact, from a Beech (*Fagus sylvatica* L.) forest and imposed an extreme climate event, consisting of drought and a single heat-pulse event, we then followed microbial community dynamics over a short time period of 28 days. During the treatment we labeled the canopy with <sup>13</sup>CO<sub>2</sub> with the goal (1) of quantifying the different time delays of assimilate arrival to belowground plant tissues and microbial phospholipid fatty acids (PLFA). From this assessment we determined the strength of plant-microbe carbon linkages under control, drought, heat, and heat-drought treatments. Based on the <sup>13</sup>C labeled isotopic PLFAs we (2) characterized microbial groups that are tightly linked to the plant-soil carbon continuum during environmental stress. And, we used 16s rRNA sequencing of bacteria from the soil mineral horizon to determine (3) the short-term changes in the active microbial community structure.

In all treatments, root tissue tended to increase in <sup>13</sup>C, indicating that labeled carbon had been allocated belowground. None of the treatments were too severe to disrupt within-plant transport over the experiment, and the depletion of the aboveground signal with the concurrent enrichment of the belowground root tissue suggested that carbon sinks belowground were still dominant. If we consider only the general PLFA marker (c16: 0), which was the most enriched across all treatments, we observed that the heat-drought monoliths were the most severely affected. We also observed that the label increased over time in the PLFA markers under drought. We infer two pathways of carbon flow to belowground microbial communities based on the time delay of labeled carbon in the PLFAs. The first pathway is evidenced from the strong increase in <sup>13</sup>C excess directly after labeling and the second pathway delivers assimilates more slowly over time. Based on our high-throughput sequencing of the 16S rRNA, we observed multiple trajectories in the community shifts due to the stress treatments.

Heat in combination with drought had a clear negative effect on diversity (only in heat-drought we saw a significant decrease in diversity) and resulted in a distinct shift in relative abundances of various phylotypes.

## RESPONDING TO CHALLENGES IN FOREST RESEARCH AND MONITORING: FROM PAST LEGACY TO INTEGRATION OF RESEARCH INFRASTRUCTURES

*Giorgio Matteucci<sup>1,2</sup>, Bruno De Cinti<sup>1</sup>*

<sup>1</sup>National Research Council of Italy, Inst. of Agroenvironmental and Forest Biology (CNR – IBAF), Italy

<sup>2</sup>National Research Council of Italy, Inst. for Agriculture and Forestry Systems in the Mediterranean (CNR-ISAFOM), Italy

E-mail: giorgio.matteucci@cnr.it

**Keywords:** Forest ecosystems, Long Term, Climate change, Biodiversity, Sustainable Forest Management

Forests have been a “study and observation” object by humans since centuries.

In this long story, the way in which humans approached forests started to change significantly in the 19<sup>th</sup> century, when it started to become clearer the need for a long-term management view aimed at achieving a permanent and sustainable production. Forest inventories and growth and yield trials started to be implemented.

In the second half on the 20<sup>th</sup> century, concerns were on forest health and the impact of pollution and long-term forest monitoring programmes were initiated. In the last part of 20<sup>th</sup> and the beginning of 21<sup>st</sup> centuries, the overarching challenge is global change effects on forest ecosystems that were addressed through impact studies (Open-top-chamber and FACE experiments) and modelling. Furthermore, ecosystem scale fluxes of carbon and other gases started to be measured (currently on more than 400 sites on different ecosystems, many of those forests). In the same period, the development of remote sensing and modelling approaches opened the way in scaling up results in spatial (remote sensing) and temporal (modelling) scales.

This long history of monitoring and research has improved a lot our understanding of the functioning and long-term trends of forest ecosystems: however, it is now time to push for an integration in large-scale European infrastructure to provide renovated objectives for the future of forest monitoring and research schemes.

“Supersites (or Level III)” research / monitoring plots, measuring process-related parameters are needed to provide information on the effects of the driving forces on forest ecosystem dynamics and functioning (e.g. C and N cycles) and on the long term responses to global changes. As an example, considering carbon sequestration, the changes of carbon storage in European forests (including soils) and their response to climate impacts can only be assessed, if their variability, uncertainty and complete element cycles are known.

Other important issues for the future of forest research and monitoring is to expand the aims to the assessment of “ecosystem services” provided and granted by forests and to use data and results from research infrastructures to inform (and drive, if possible) forest management.

The presentation will address how basic and intensive monitoring networks should be coupled to research sites where process-based studies may provide deeper process understanding that can be used for addressing challenges for 21<sup>st</sup> century forest ecosystem research.

### Acknowledgements

Ideas and concepts presented have been developed within EU (CarboEurope, EUFORINNO, eLTER, ExpeER, FutMon, EnvEurope, ManFor, Smart4Action) and national-funded projects (CarboItaly, PRIN2012, NextData, I-Amica, ALForLab).

## THE EFFECT OF FOREST TREES ON MICROBIAL PROCESSES INVOLVED IN CARBON CYCLING

Petr Baldrian<sup>1</sup>

<sup>1</sup>Institute of Microbiology of the ASCR, Videnska 1083, 14220 Praha 4, Czech Republic

E-mail: baldrian@biomed.cas.cz

**Keywords:** decomposition, forest soil, litter, microbial ecology, plant-microbe interactions

Trees largely shape ecosystem processes in forest ecosystems, being the dominant primary producers responsible for the bulk of C fixation and its allocation belowground. The production of litter as well as root exudates affects the community of microbial decomposers in forest soils, most importantly the decomposers and root-associated symbionts. This is reflected by the facts that the seasonality of the composition of soil microbial communities and their activity in decomposition correspond to the seasonal changes in C allocation belowground and litter production (Voříšková et al. 2014) and demonstrated by the complex restructuring of microbial community and pronounced alterations in decomposition patterns in the case of tree dieback, associated with the transition from soil fungal communities dominated by ectomycorrhizal root symbionts to those mainly composed of saprotrophic taxa (Štursová et al. 2014). Microbial community analysis and shotgun metatranscriptomics can help to explore the differences in microbial communities among various trees and their activity throughout the year. Fungi highly contribute to the total microbial transcription in litter and soil while the Proteobacteria, Acidobacteria and Actinobacteria are most transcriptionally active among bacteria in an acidic *Picea abies*-dominated forest. Notably, the share of fungal transcription decreases dramatically between summer and winter and the transcription of genes involved in mycorrhiza formation is most affected. In contrast, the relative activity of Acidobacteria increases in winter. The results show that both tree species composition and the seasonal changes of their physiology largely affect both community composition and processes involved in decomposition and C cycling.

Štursová M, Šnajdr J, Cajthaml T, Bárta J, Šantrůčková H, Baldrian P (2014) When the forest dies: the response of forest soil fungi to a bark beetle-induced tree dieback. *ISME Journal* 8: 1920-1931.

Voříšková J, Brabcová V, Cajthaml T, Baldrian P (2014) Seasonal dynamics of fungal communities in a temperate oak forest soil. *New Phytol* 201: 269-278.

## TREE BIODIVERSITY AND ECOSYSTEM FUNCTION

*Hans Göransson<sup>1</sup>, Andrew Smith<sup>2</sup>, Iftekhar Ahmed<sup>1</sup>, Douglas Godbold<sup>1</sup>*

<sup>1</sup> Institute of Forest Ecology, Universität für Bodenkultur, 1190 Vienna, Austria

<sup>2</sup> School of Environment, Bangor University, Bangor, UK

E-mail: hans.goeransson@boku.ac.at

**Keywords:** mixed stands, elevated CO<sub>2</sub>, drought

As a consequence of land-use change and the burning of fossil fuels, atmospheric concentrations of CO<sub>2</sub> are increasing which directly affects the carbon cycling in forests but also affects the climate. One of the consequences of climate change in Europe is an increase in long drought periods. Both the effect of elevated levels of CO<sub>2</sub> and drought has been extensively tested on monocultures. When modelling the effect of elevated CO<sub>2</sub> and climate change on mixed stands one often has to assume that the different tree species respond as they would have grown in monocultures due to the lack of data from mixed stands.

To investigate how mixed stands were affected by elevated CO<sub>2</sub>, *Alnus glutinosa*, *Betula pendula* and *Fagus sylvatica* were planted in areas of single species and a three species polyculture. The trees were exposed to ambient or elevated CO<sub>2</sub> (580 μmol mol<sup>-1</sup>) for 4 years. Plots at the same site and planting scheme were also used for a throughfall reduction experiment using sub canopy roofs, which removed 70 % of the throughfall. Hence soil water potential was reduced over two consecutive years. As expected biomass increase due to elevated CO<sub>2</sub> and decreased due to drought. But the response of the mixed stand and the individual tree species within the mixture could not be foreseen by the response in the monocultures.

During elevated CO<sub>2</sub> the overyielding in the mixed stands was lower than the control, whereas it remained the same under drought. In the controls *Alnus glutinosa* benefited the most from growing in mixture and overyielded with over 37 %. But under elevated CO<sub>2</sub> the over yielding was only 14 % whereas *Betula pendula* had the same overyielding (8 %) independent of CO<sub>2</sub> concentration. During drought experiment *Betula pendula* was over yielding the most in the control. Under drought however, the dominance of *Betula pendula* decreased and *Alnus glutinosa* started to contribute to the overyielding in the mixed stand.

## FINE-ROOT TRAITS AND SOIL PROCESSES IN BOREAL FORESTS

*Heljä-Sisko Helmisaari<sup>1</sup>, Jaana Leppälampi-Kujansuu<sup>1</sup>*

<sup>1</sup> University of Helsinki, Dept. Forest Sciences, Helsinki, Finland

E-mail: helja-sisko.helmisaari@helsinki.fi

Fine root traits and processes were studied during the last decade in Norway spruce and Scots pine forests on a climatic and soil fertility gradient in Finland. The overall aim was to understand the forest above- and belowground relationships and quantify carbon (C) allocation in relation to environmental factors, including site fertility. C allocation is one way for trees to cope with nutrient deficiencies. Both tree species responded to nitrogen (N) deficiency resulting from low fertility or a short growing season, or both, by maintaining more EcM root tips (Helmisaari et al., 2009; Ostonen et al. 2011) per unit foliage and basal area which may be associated with an increased share of N uptake in an organic form. Stand basal area predicted fine root biomass better than any other variable alone (Lehtonen et al. 2015). Fine root turnover rate was slower in the north, and the less productive sites produced relatively more litter belowground than aboveground (Leppälampi-Kujansuu et al. 2014). Increased partitioning to belowground in nutrient deficiency was recently indirectly shown in a pine stand in Sweden where N-fertilization-induced increase of aboveground net primary production was associated with a decreased partitioning to belowground (Lim et al., 2015). Needles and roots in the north vary in chemistry, containing more water-extractable compounds, especially phenols, than in the south (Hilli et al., in prep.). Polyphenols have been shown to regulate the formation of mor-humus by inhibiting N mineralisation. We conclude that coniferous roots may have a greater impact to soil humus formation in the north and / or less fertile sites.

Helmisaari H-S, Ostonen I, Löhmus K, Derome J, Lindroos A-J, Merilä P, Nöjd P (2009) Ectomycorrhizal root tips in relation to site and stand characteristics in Norway spruce and Scots pine stands in boreal forests. *Tree Physiol.* 29(3): 445-456.

Lehtonen A, Palviainen M, Ojanen P, Kalliokoski T, Nöjd P, Kukkola M, Penttilä T, Mäkipää R, Leppälampi-Kujansuu J, Helmisaari H-S (2015) Modelling fine root biomass of boreal tree stands by using site and stand variables. *For. Ecol. Manage.* (accepted)

Leppälampi-Kujansuu J, Aro L, Salemaa M, Hansson K, Berggren Kleja D, & Helmisaari H-S (2014) Fine root longevity and carbon input into soil from below- and aboveground litter in climatically contrasting forests. *For. Ecol. Manage.* 326: 79-90.

Lim H, Oren R, Palmroth S, Tor-Ilngern P, Mörling T, Näsholm T, Lundmark T, Helmisaari H-S, Leppälampi-Kujansuu J. & Linder S (2015) Inter-annual variability of precipitation constrains the production response of boreal *Pinus sylvestris* to nitrogen fertilization. *For. Ecol. Manage.* 348: 31-45

Ostonen I, Helmisaari H-S, Borken W, Tedersoo L, Kukumägi M, Bahram M, Lindroos A-J, Nöjd P, Uri V, Merilä P, Asi E, Löhmus K (2011) Fine root foraging strategies in Norway spruce forests across a European climate gradient. *Global Change Biol.* 17: 3620-3632.

## CARBON INPUT INTO SOIL ORIGINATING FROM FINE ROOT AND FOLIAGE LITTER

*Jaana Leppälammii-Kujansuu<sup>1</sup>, Lasse Aro<sup>2</sup>, Maija Salemaa<sup>2</sup>, Karna Hansson<sup>3</sup>, Sune Linder<sup>4</sup>, Dan Berggren Kleja<sup>4</sup>, Heljä-Sisko Helmisaari<sup>1</sup>*

<sup>1</sup> University of Helsinki, Dept. Forest Sciences, Helsinki, Finland

<sup>2</sup> Natural Resources Institute, Parkano / Vantaa, Finland

<sup>3</sup> Centre INRA de Nancy, Nancy, French

<sup>4</sup> Swedish Univ. Agricultural Sciences, Uppsala / Alnarp, Sweden

E-mail: jaana.leppalammi-kujansuu@helsinki.fi

**Keywords:** fine root litter, foliage litter, root turnover, minirhizotron, carbon input

Carbon models require knowledge of litter inputs. Contrary to the foliage litter, the data quantifying the major belowground carbon input, fine root production, is very scarce, especially using comparable methods. The minirhizotron method, i.e. belowground digital filming of roots, is a method which is applied worldwide to study fine root turnover. An estimate of annually produced fine root litter can be calculated by multiplying the root standing crop (from soil cores) by the root turnover rate. In Finland the minirhizotron results have hitherto been reported only from three Norway spruce stands, of which in only two the foliage litter fall was also measured. Here we present the results from these two sites from Finland and one site from Sweden. Altogether in Sweden the respective results have been published from five Norway spruce sites, and one birch and pine sites. In a changing climate there is an urgent need, not least for the modellers, to have more data on root turnover, especially covering the whole ecosystem. Soil fertility or temperature changes will affect not only the aboveground tree growth but also the belowground parts which are tightly interconnected. Therefore, quantification of carbon input into the soil should always include both fine root and foliage litter.

### Acknowledgements

The Academy of Finland, project 260708

Leppälammii-Kujansuu J et al. (2014) Fine root longevity and carbon input into soil from below- and aboveground litter in climatically contrasting forests. *For Ecol and Management* 326: 79-90.

Leppälammii-Kujansuu J et al. (2014) Fine root turnover and litter production of Norway spruce in a long-term temperature and nutrient manipulation experiment. *Plant Soil* 374: 73-88.

## FINE ROOT PHYSIOLOGICAL AND MORPHOLOGICAL TRAITS OF FINNISH BOREAL TREE SPECIES

*Naoki Makita*<sup>1,2</sup>, *Heljä-Sisko Helmisaari*<sup>1</sup>, *Vesa Nousiainen*<sup>1</sup>, *Jaana Leppälampi-Kujansuu*<sup>1</sup>

<sup>1</sup> Department of Forest Sciences, University of Helsinki, Helsinki 00014, Finland

<sup>2</sup> Forestry and Forest Products Research Institute, Kyoto 612-0855, Japan

E-mail: makita701@gmail.com

**Keywords:** boreal forest, Norway spruce, root respiration, Scots pine, Silver birch

Tree species have an impact on growth and survival of woody roots, with potential consequences for source and sink activities of root organ. However, the effects of different tree species on specific traits of fine roots are still unclear. We elucidated the magnitude of physiological and morphological traits of fine roots of different tree species and the relationships between specific respiration rates and morphological traits in a boreal forest in Kivalo, northern Finland. In this study, we used fine roots of Norway spruce (*Picea abies* (L.) Karst.), Scots pine (*Pinus sylvestris* L.), and Silver birch (*Betula pendula* Roth.), which are dominant tree species in Fennoscandia. The specific respiration rate of fine root segments was measured in the forest in July 2014. The N concentration and morphology, including root tissue density (RTD; g cm<sup>-3</sup>) and specific root length (SRL; m g<sup>-1</sup>) were then observed in the laboratory. Mass-based specific root respiration rates, the N concentration, and morphology differed among tree species. In all three species, the respiration rates were significantly positively correlated with root N and SRL, and negatively correlated with RTD. The regression slopes of respiration with root N and RTD were significantly higher in birch than in spruce and pine. Although no differences in slopes of root respiration with SRL were found across the species, there were significant shifts in intercept along the common slope. These results suggest that a contrasting pattern in interspecific relationships between root respiration and N, RTD, and SRL exists. Therefore, tree species influenced the magnitude of fine root physiological and morphological traits and their relationships. Boreal tree species also affected the trait-based root performance by covarying the root respiration, root N and morphology for optimizing potential nutrient and water uptake.

## DISTURBANCE AND RESILIENCE IN FORESTS: EXAMPLES FROM NEW JERSEY PINE BARRENS

*John Dighton<sup>1</sup>, Dennis M. Gray<sup>1</sup>*

<sup>1</sup> Rutgers University Pinelands Field Station, PO Box 206, 501 Four Mile Road, New Lisbon, NJ 08064 USA

E-mail: dighton@camden.rutgers.edu

**Keywords:** pine barrens, prescribed burning, N-deposition, harvesting

The NJ pine barrens consists of pitch pine and oaks with an understory of ericaceous shrub species on marine sediment oligotrophic sandy soils. Historically, the forest has been cut for charcoal production. Although a protected area, disturbance comes from natural wildfires, prescribed burning, logging, recreational vehicle use and non-point source pollutants. Alternate stable states of graminoid and lichen understory vegetation are thought to be derived from disturbance factors. Here we will examine the impacts of experimental tree harvesting with soil disturbance, prescribed fire and nitrogen deposition, which appear to have only transitory effects on soil factors and understory vegetation. In a forest protected for its vulnerability to disturbance, we show that it has considerable resilience.

### Acknowledgements

Funds provided by National Science Foundation, United States Department of Agriculture, New Jersey Department of Environmental Protection.

Adams-Krumins, Dighton J, Gray D, Franklin RB, Morin P & Roberts MS (2009) Soil microbial community response to nitrogen enrichment in two scrub oak forests. *For. Ecol. Manage.* 258: 1383-1390.

Dighton J, Tuininga AR, Gray DM, Huskins RE & Belton T (2004) Impacts of atmospheric deposition on New Jersey pine barrens forest soils and communities of ectomycorrhizae. *For. Ecol. Manage.* 201: 131-144.

Gray DM, Swanson J, Dighton J (2012) The influence of contrasting ground cover vegetation on soil properties. *Appl. Soil Ecol.* 60: 41-48.

Geng Y, Dighton J, Gray DM (2012) The effects of thinning and soil disturbance on enzyme activities under pitch pine soil in New Jersey Pinelands. *Appl. Soil Ecol.* 62: 1-7.

Tuininga AR, Dighton J (2004) Changes in ectomycorrhizal communities and nutrient availability following prescribed burning in two upland pine-oak forests in the New Jersey Pine Barrens. *Can. J. For. Res.* 43: 1755-1765.

## IMPACTS OF INCREASED NITROGEN AVAILABILITY ON THE STRUCTURE AND FUNCTIONING OF A MEDITERRANEAN BASIN ECOSYSTEM

*Teresa Dias<sup>1</sup>, Maria Amélia Martins-Loução<sup>1</sup>, Lucy Sheppard<sup>2</sup>, Cristina Cruz<sup>1</sup>*

<sup>1</sup> Universidade de Lisboa, Faculdade de Ciências, cE3c, Portugal

<sup>2</sup> Centre of Ecology and Hydrology (CEH), Bush Estate, Penicuik, EH26 OQB, United Kingdom

E-mail: ccruz@fc.ul.pt

**Keywords:** ammonium, biodiversity, impacts, nitrate, N cycling, N retention, plants, soil protection

Increased nitrogen (N) availability, resulting from agro-industrial activities, affects ecosystems' health and stability. Mediterranean Basin ecosystems are biodiversity hotspots threatened by N deposition, however, little or no information is available on their responses to increased N. We develop an integrated system-level approach to study the responses of an N limited Mediterranean Basin maquis to increased N availability. Since 2007, the form and dose of N available at the experimental site (ambient N deposition 5.2 kg ha<sup>-1</sup> yr<sup>-1</sup> and soil N 0.1 %) was modified by the addition of 40 kg NH<sub>4</sub><sup>+</sup>-N ha<sup>-1</sup> yr<sup>-1</sup>, 40 and 80 kg NH<sub>4</sub>NO<sub>3</sub>-N ha<sup>-1</sup> yr<sup>-1</sup>. Control plots were not fertilized. Over the following years, the effects on plant and soil microbial diversity and ecosystem services were assessed: soil N retention, below and aboveground C sequestration, and soil protection.

Over the first year, the added N was retained by the system and detected in the autumn as soil inorganic N. The increased N promoted plant and soil microbial diversity.

On the fifth spring of N additions, plant richness increased with enhanced N and was more related to the cumulative ammonium than with the cumulative nitrate. Exposure to 40 kg NH<sub>4</sub><sup>+</sup>-N ha<sup>-1</sup> yr<sup>-1</sup> (either alone or with nitrate) enhanced plant richness, but did not increase aboveground C sequestration or soil protection; soil N retention even decreased under 80 kg N ha<sup>-1</sup> yr<sup>-1</sup>. The treatment containing less ammonium, 40 kg NH<sub>4</sub>NO<sub>3</sub>-N ha<sup>-1</sup> yr<sup>-1</sup>, did not enhance plant diversity but promoted aboveground C sequestration and soil protection. The dataset permitted the first estimation of an N critical load for this European habitat (20-30 kg N ha<sup>-1</sup> yr<sup>-1</sup>).

Data suggest that agriculture, the main source of NH<sub>4</sub><sup>+</sup>, may affect the structure of the neighboring maquis, promoting soil erosion and N leakage. In contrast, industrial and urban activities that increase N availability as NO<sub>x</sub> may increase fire risk. This is of significant importance for land-use management in biodiverse and fragmented ecosystems such as the Mediterranean ones.

### Acknowledgements

FCT for financial support, Parque Natural da Arrábida

## FOREST – AGRICULTURE INTERFACE: A CASE OF BUCKWHEAT

*Ivan Kreft*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: ivan.kreft@guest.arnes.si

**Keywords:** buckwheat, environment, sustainability, project EUFORINNO, final conference

Buckwheat is in Bhutan, Japan, China and Korea used mostly to prepare noodles and other pasta products. In Italy, buckwheat flour is used to prepare a typical pasta and in Austria and Slovenia traditional dishes such as kasha and bread are made (Vombergar et al., 2014). In Asia (Japan, Korea, China, Nepal) buckwheat is cultivated in high altitudes, sometimes above 3000 m. Humans used fire as an effective management tool to transform the evergreen forests into secondary vegetation, such as buckwheat cultivation (Shu et al., 2013). Buckwheat fields are often mixed with areas covered by forest. Such systems are traditional in Northern Italy, Slovenia and Bosnia and Herzegovina. Forest trees give to buckwheat fields protection from strong winds and are a shelter for propagation of pollinators. In Japan, the abundance of the important pollinator wild bee *Apis cerana* in the buckwheat fields show a stronger positive correlation with the total area of surrounding natural forests compared to that of surrounding plantation forests (Taki et al., 2011). Buckwheat fields are source not only for nectar for wild insects, but as well a food source for birds and wild animals. It is important that in buckwheat no spraying is needed, so buckwheat is a source of »organic« food for wildlife.

### Acknowledgements

This research was financed by the ARRS (Slovenian Research Agency), through the program P3-0395 "Nutrition and Public Health" and projects J4-4224, J4-5524, supported by EUFORINNO 7<sup>th</sup> FP EU Infrastructure Program (RegPot No. 315982).

Shu JW, Sasaki N, Takahara H, Hase Y (2013) Vegetation and fire history with their implication for climatic change and fire events since the last deglacial in the Aso Valley, central Kyushu, southwestern Japan: new pollen and charcoal data. *Veget Hist Archaeobot* 22: 285–298.  
 Vombergar B, Kreft I, Horvat M, Vorih S (2014) *Ajda – Buckwheat*. Ljubljana: Založba kmečki glas, pp. 131.  
 Taki H, Yamaura Y, Okabe K, Maeto K (2011) Plantation vs. natural forest: Matrix quality determines pollinator abundance in crop fields. *Scientific Reports* 1, Art. No. 132: 1-4.

## PROTECTION OF FORESTS UNDER INTERNATIONAL LAW

*Maša Kovič Dine*<sup>1</sup>

<sup>1</sup> Faculty of Law, University of Ljubljana, Poljanski nasip 2, SI – 1000 Ljubljana, Slovenia

E-mail: masa.kovic-dine@pf.uni-lj.si

**Keywords:** international law, forest protection, climate change, common concern of humanity

The presentation will address the lack of legally binding international regulation on forest protection and present the recognition of forests as a common concern of humanity as a possible solution for adoption of an international convention on forests. For centuries forests have been recognized as one of the most important natural resources. The fear of overexploitation of forest and loss of their ecosystem benefits has led the process of negotiations for a binding international forest convention. However, after nearly forty years of negotiations, States have still not been able to reach an agreement on a legally binding international document. The majority of developing States oppose such a document out of fear that it would deprive them of their sovereign right to exploit their own natural resources in accordance with their development policies. On the contrary, developed States see such a document as the only means to prevent major deforestation in the fight against climate change. A solution to overcome this difference of opinion among States is the recognition of forests as a common concern of humanity. Legally, both climate change and biodiversity have already been recognized as common concern of humanity. However, international environmental law presents no definition of this concept. After the analysis of the common denominators of climate change and biodiversity, the author has developed a working definition of this concept that also enables its application to forests.

Kovič Dine M (2014) Regulation of forest ecosystem services: Slovenian Act on Forests as an example of good practice. In: Sancin V et al. (Eds.). *International environmental law: Contemporary concerns and challenges in 2014*. Ljubljana: IUS Software, GV založba, pp. 401-408.

Kovič Dine M (2013) Can the world heritage regime aid the adoption of an international framework for forest protection. In: Yankov A (Ed.). *Studies on international law*. Sofija: Bolgarska asociacija po međunarodno pravo, pp. 137-149.

Kovič Dine M (2013) Mednarodne obveznosti držav za zagotavljanje trajnostnega razvoja gozdov. *Zbornik znanstvenih razprav* 73 (1): 161-190.

Kovič Dine M (2012) Forests: does state sovereignty hinder their protection at the international level? In: Sancin V (Ed.). *International environmental law: Contemporary concerns and challenges*. Ljubljana: GV založba, pp. 109-128.

## IMPACT OF FOREST HARVESTING PRACTICES ON BASE CATION BUDGETS – A SUSTAINABILITY STUDY USING MEASURED AND MODELLED DATA

*Mike Starr<sup>1</sup>, Päivi Merilä<sup>2</sup>, Brandon Stephens<sup>2,3</sup>, Antti-Jussi Lindroos<sup>2</sup>, Tiina M. Nieminen<sup>2</sup>, Pekka Nöjd<sup>2</sup>, Kirsti Derome<sup>2</sup>, Liisa Ukonmaanaho<sup>2</sup>*

<sup>1</sup> University of Helsinki, Finland

<sup>2</sup> Natural Resources Institute Finland (Luke), Finland

<sup>3</sup> US Forest Service, US

E-mail: [mike.starr@helsinki.fi](mailto:mike.starr@helsinki.fi)

We simulated the impacts of harvestings on base cation (Ca, Mg, and K) budgets for Finnish forests under three harvesting practices: stem-only (SOH), whole-tree (WTH), and WTH + stump and coarse root harvesting (WTSR). The study included five Scots pine and five Norway spruce stands belonging to the UN-ECE ICP Forests Level II programme in Finland. A mass balance approach was used in which the base cation (BC) input fluxes total deposition (TD) and weathering (W) were balanced against the output fluxes leaching (L) and harvesting removals (H). The mean annual fluxes were estimated using a combination of measured and modelled data. The impact of harvesting practice evaluated in relation to soil BC stocks, stand tree species, and climate (latitude). Positive values of  $TD + W - L - H$  were considered to indicate sustainability while negative values were considered to indicate unsustainability. Spruce stand TD of BC correlated negatively with latitude, and was significantly higher than that for the pine stands. BC amounts in the hypothetically harvested removals of all three harvesting practices were larger for the spruce than for the pine stands. Harvesting removals of BC were significantly higher with WTH and WTSR than with SOH, but BC removals with WTH and WTSR did not significantly differ from each other. Harvesting intensity negatively impacted the sustainability of BC. WTH in the spruce stands had a more negative impact on soil BC stocks than in the pine stands. The mass balance for K in all stands and for Ca in the spruce stands were, on average, negative under WTH and WTSR scenarios. For Mg, the mass balances for both spruce and pine were, on average, positive under all harvesting scenarios.

## VARIANCE DECOMPOSITION OF STEM BIOMASS INCREMENT PREDICTIONS FOR EUROPEAN BEECH

*J. A. Horemans<sup>1</sup>, M. Bosela<sup>2,3</sup>, L. Dobor<sup>4</sup>, M. Barna<sup>5</sup>, G. Deckmyn<sup>1</sup>, R. Ceulemans<sup>1</sup>*

<sup>1</sup> Centre of Excellence PLECO, Department of Biology, University of Antwerp, Universiteitsplein 1, B-2610 Wilrijk, Belgium

<sup>2</sup> National Forest Centre - Forest Research Institute Zvolen, Slovak Republic

<sup>3</sup> Czech University of Life Sciences, Praha, Czech Republic

<sup>4</sup> Eötvös Loránd University, Budapest, Hungary

<sup>5</sup> Slovak Academy of Sciences, Zvolen, Slovakia

E-mail: Joanna.Horemans@uantwerpen.be

**Keywords:** empirical model, process-based model, climate scenario, forest management, forest modeling

Different sources of uncertainty in the results of forest model simulations were identified and quantified. The responses of annual stem biomass increment of European beech (*Fagus sylvatica* L.) to changes in climate and in forest management were simulated using two forest models, i.e. the empirical SIBYLA model and the process-based ANAFORE model. Both models were calibrated using experimental tree growth data from four plots in central Slovakia between 1989 and 2003. Three of these plots were subjected to different levels of thinning while one was left untouched as a control. Ten regional climate models (RCMs) based on the IPCC scenario A1B provided an ensemble of climate projections up to 2100. Simulations were made: (i) with the two forest models; (ii) for four thinning intensities; (iii) using the 10 RCMs; and (iv) for three time windows, each spanning 15 years. The contribution of these four components to the total variance in the simulated stem biomass increment was quantified. The forest model type explained most of the total variance in the simulation results, followed by forest management (i.e. thinning) and the time window. The effect of the RCMs on model uncertainty was limited. Stem biomass increment predictions obtained from both models used were different in absolute terms, but the models agreed well in their relative response to RCM, to forest management and to time window. The total variance of the predictions was 10 times higher for the process-based ANAFORE model than for the empirical SIBYLA model.

### Acknowledgements

Research supported by the Belgian Science Policy (MASC project), by the Slovak Research and Development Agency, the Slovak Grant Agency and the Hungarian Scientific Research Fund.

## ADJUSTMENT CAPACITY OF MARITIME PINE CAMBIAL ACTIVITY IN DROUGHT-PRONE ENVIRONMENTS

*Joana Vieira<sup>1</sup>, Filipe Campelo<sup>1</sup>, Sergio Rossi<sup>2</sup>, Ana Carvalho<sup>1</sup>, Helena Freitas<sup>1</sup>, Cristina Nabais<sup>1</sup>*

<sup>1</sup> CFE – Centre for Functional Ecology, Department of Life Sciences, University of Coimbra, Coimbra, Portugal

<sup>2</sup> Département des Sciences Fondamentales, Université du Québec à Chicoutimi, Chicoutimi, Canada

E-mail: joana.vieira@uc.pt

**Keywords:** cambial activity, anatomy, intra-annual density fluctuations, drought, Mediterranean

Intra-annual density fluctuations (IADFs) are anatomical features formed in response to changes in the environmental conditions within the growing season. These anatomical features are commonly observed in Mediterranean pines, being more frequent in younger and wider tree rings. However, the process behind IADF formation is still unknown. Weekly monitoring of cambial activity and wood formation would fill this void. Although studies describing cambial activity and wood formation have become frequent, this knowledge is still fragmentary in the Mediterranean region. Here we present data from the monitoring of cambial activity and wood formation in two diameter classes of maritime pine (*Pinus pinaster* Ait.) over two years in order to test: (i) whether the differences in stem diameter in an even-aged stand were due to timings and/or rates of xylogenesis; (ii) if IADFs were more common in large trees; and (iii) if their formation is triggered by cambial resumption after the summer drought. Larger trees showed higher rates of cell production and longer growing seasons, due to an earlier start and later end of xylogenesis. When a drier winter occurs, larger trees were more affected, probably limiting xylogenesis in the summer months. In both diameter classes a latewood IADF was formed in response to late-September precipitation, confirming that the timing of the precipitation event after the summer drought is crucial in determining the resumption of cambial activity and whether or not an IADF is formed. It was the first time that the formation of a latewood IADF was monitored at a weekly time scale in maritime pine. The capacity of maritime pine to adjust cambial activity to the current environmental conditions represents a valuable strategy under the future climate change conditions.

## ABOVE AND BELOWGROUND TREE BIOMASS ALLOCATIONS- AS INFLUENCED BY SPECIES COMPOSITION IN TEMPERATE DECIDUOUS PLANTATION FOREST

*Iftekhhar U. Ahmed<sup>1</sup>, Andy R. Smith<sup>2</sup>, Hans Göransson<sup>1</sup> and Douglas L. Godbold<sup>1</sup>*

<sup>1</sup> Institute of Forest Ecology, Universität für Bodenkultur, 1190 Vienna, Austria

<sup>2</sup> School of Environment, Bangor University, Bangor, UK

E-mail: iftekhar.ahmed@boku.ac.at

**Keywords:** allometry, woody biomass, root biomass, polyculture

Afforestation and management of first growing tree species enhance the terrestrial carbon stocks by offsetting anthropogenic emissions of CO<sub>2</sub>, and thus can be the potential tools for mitigation of climate change impacts. Effects of species polyculture on ecosystem C balance in plantation forest has been assessed by some authors. However, the mechanistic understanding of how stand composition affect the tree biomass in polyculture compare to monoculture is still unclear. We estimated above and belowground biomass by developing species specific allometric equations and sequential coring and subsequent litter flux and fine root turnover in mono and polyculture stands of *Betula pendula*, *Alnus glutinosa* and *Fagus sylvatica* in the temperate forest ecosystems of North Wales, UK. Significantly higher woody biomass was observed in the single stand of *B. pendula* and *A. glutinosa* then *F. sylvatica*. Fine root (<2 mm) biomass was higher in *A. glutinosa* whilst the root turnover rate was higher in *B. pendula*. At the stand level, clear additive mixture effects on above ground biomass was observed, however at the tree level, *B. pendula* tended to lower biomass in mixture presumably due to suppression by the faster growing *A. glutinosa*. Our results indicated that selection of tree species in polyculture can affect the biomass allocation through species interactions. The concentration of SOC at the top 20 cm of plantation plots increased following the order: *F. sylvatica* < species mixture < *A. glutinosa* < *B. pendula*, reflecting the impacts of species specific functional traits such as biomass inputs and litter quality. Over all, with judicious selection of species, tree polyculture might be the best option in plantation design.

## CARBON CYCLING IN LOWLAND OAK FOREST ECOSYSTEMS

*Hrvoje Marjanović<sup>1</sup>, Maša Zorana Ostrogović Sever<sup>1</sup>, Mislav Anić<sup>1</sup>, Anikó Kern<sup>2</sup>*

<sup>1</sup> Croatian Forest Research Institute, Cvjetno naselje 41, 10450 Jastrebarsko, Croatia

<sup>2</sup> Department of Geophysics and Space Science, Eötvös Loránd University, Budapest, Hungary

E-mail: hrvojem@sumins.hr

**Keywords:** forest productivity, eddy covariance, dendrometers, NPP, MODIS

Lowland oak forests are among the most productive forest ecosystems in the region of SE Europe, with high ecological and commercial value. Monitoring and quantifying carbon fluxes in them is necessary to meet the reporting requirements of UNFCCC and Kyoto protocol, but even more so it is needed in our efforts to understand carbon dynamics and underlying processes and key drivers.

Our study is conducted in Jastrebarsko forest, which is a part of 12 kHa of lowland forests of Kupa river basin. Forests within the basin are managed with continuous cover, even-aged management system. Eight stands dominated by Pedunculate oak (*Quercus robur* L.), aged 5 to 168 years, were selected in spring 2010 to represent an oak chronosequence, and in one of them (the 37 year old) an eddy covariance flux tower was already installed in September 2007.

Carbon stocks and fluxes are monitored using classical dendrometric techniques, eddy covariance (EC), and soil respiration measurements (SR). Decomposition fluxes from fine woody debris are assessed with decomposition experiment, while fine root and soil carbon stocks are measured with soil core sampling. Depth to the ground water was also recorded during vegetation season.

Carbon pool stocks change during stand development with soil as a dominant carbon pool in young age, while live biomass carbon pool dominates in older age. Carbon stocks in soil organic layer increase with stand age, but in mineral soil layer no statistically significant age-dependent trend was observed. When successfully regenerated, as in our case, oak stand becomes carbon sink very early in a development phase, between the age of 5 and 13 years and remains a carbon sink even after the age of 160 years, or until final harvest.

Weekly and annual productivity (GPP and NPP) estimates from EC were compared with MODIS NPP and GPP (MOD17, col. 5.5) products from NASA's TERRA satellite, and validated with weekly NPP estimated with dendrometer measurements and SR for years 2008-2012. Results indicate good agreement in "average" years, but poor agreement with MODIS after prolonged dry period when MODIS underestimates the productivity. Ground water, i.e. soil moisture seems to be the key element, as MODIS routines do not consider soil water availability in calculations of GPP and NPP.

### Acknowledgments

The research has been supported in part by the Croatian Science Foundation (HRZZ UIP-11-2013-2492) and Hungarian Scientific Research Fund (OTKA PD-111920).

Marjanović H, Ostrogović MZ, Alberti G, Balenović I, Paladinić E, Indir K, Peressotti A, Vuletić D (2011) Carbon dynamics in younger stands of Pedunculate oak during two vegetation periods. *Šumar. list* 135 (special issue): 59–73.

Ostrogović MZ (2013) Carbon stocks and carbon balance of an even-aged Pedunculate Oak (*Quercus robur* L.) forest in Kupa river basin. *Dissertation*. Croatian Forest Research Institute, 130 pp.

Running SW, Nemani RR, Heinsch FA, Zhao M, Reeves M, Hashimoto H (2004) A Continuous Satellite-Derived Measure of Global Terrestrial Primary Production. *Bioscience* 54(6): 547-560.

## ACCELERATING FOREST MORTALITY

*Nate McDowell*<sup>1,2</sup>

<sup>1</sup> Los Alamos National Lab, Los Alamos, NM 87545, USA

<sup>2</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: mcdowell@lanl.gov

**Keywords:** project EUFORINNO, final conference, 7FP, programme, topics

Forest mortality is rising throughout the northern hemisphere and may be driven in part by rising temperature and increasing frequency or intensity of droughts. The EUFORINO effort has enabled improved technical, computational, and theoretical abilities to investigate the patterns, physiology, and predictions of forest mortality in temperate forests. Evidence thus far suggests that rising temperature is the primary culprit driving rising mortality via increasing vapor pressure deficit (VPD), which forces increasing vulnerability to carbon starvation and hydraulic failure. Predictions suggest that rising VPD will cause a shift in forest structure from tall trees to shorter shrubs and weeds, and will particularly impact gymnosperms more than angiosperms. Global evidence suggests that increasing droughts will also impact angiosperms as well, however, and the bulk of evidence points to increasing vulnerability of forests to climate change throughout the Earth. We review the evidence underlying these predictions.

## OZONE RESEARCH, *QUO VADIS?* LESSONS FROM THE FREE-AIR CANOPY FUMIGATION EXPERIMENT AT KRANZBERG FOREST

*R. Matyssek*<sup>1</sup>

<sup>1</sup> Technische Universität München, Ecophysiology of Plants, D-85354 Freising-Weißenstephan, Germany

E-Mail: matyssek@wzw.tum.de

**Keywords:** ozone, climate change, Supersite network, integrative research, hemispheric scope

Enhanced ground-level ozone (O<sub>3</sub>) has gained awareness worldwide as an agent of climate change. Forest ecosystem response to O<sub>3</sub> has remained uncertain, though, aside from scarcely validated modelling predictions. The 8-year free-air O<sub>3</sub> canopy fumigation experiment at Kranzberg Forest (Germany) was the first to examine adult forest trees of climax species (evergreen *Picea abies*, deciduous *Fagus sylvatica*). What was learnt, and what is to be strived for?

Key findings will be highlighted from the tree-soil system as a starting point for defining upcoming conceptual and methodological challenges. Amongst such, perception of O<sub>3</sub> as part of multi-factorial, abiotic and biotic interaction networks and effective dose-related tree and stand-level risk assessment are ultimate requirements. Rationales must be both mechanistic and holistic, and ecosystem-level O<sub>3</sub> research must reach out for hemispheric scopes.

Integrated empirical and modelling approaches are to be conducted within a global “Supersites for Forest Research” network, for which a concept is outlined that unifies mechanistic clarification, process scaling and modelling-guided risk assessment. Will O<sub>3</sub> studies remain a “non-topic” within climate change research?

Matyssek R, Wieser G, Fleischmann F, Grünhage L (2013) Ozone Research, quo vadis? Lessons from the free-air canopy fumigation experiment at Kranzberg Forest. In: Matyssek R, Clarke N, Cudlin P, Mikkelsen TN, Tuovinen J-P, Wieser G, Paoletti E (eds) *Climate Change, Air Pollution and Global Challenges: Understanding and Perspectives from Forest Research*. *Developments in Environmental Science* 13, Elsevier, pp. 103-129.

## OVERVIEW OF EUFORINNO PROJECT

*Hojka Kraigher<sup>1</sup>, Tjaša Baloh<sup>1</sup>, Robert Robek<sup>1</sup>, Tine Grebenc<sup>1</sup>, Tom Levanič<sup>1</sup>, Primož Simončič<sup>1</sup>*

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: [hojka.kraigher@gozdis.si](mailto:hojka.kraigher@gozdis.si)

**Keywords:** Capacities, support action, scientific excellence, research infrastructure, visibility

EUFORINNO: European Forest Research and Innovation is an FP7 CAPACITIES project (RegPot No. 315982) coordinated by the Slovenian Forestry Institute (SFI). It provides a mean for SFI to raise its scientific excellence and infrastructure, and better exploit its innovative outputs. The project focuses on filling the gaps in four research areas: i) genetic monitoring; ii) biodiversity and functional diversity; iii) belowground complexity; and iv) ecosystem carbon exchange in time and space; and in transversal activities: publishing, innovation and IP management, and networking, contributing to the exploitation of innovation.

EUFORINNO relies on 8 excellent research partners who support SFI in building its long-term strategy, welcome researchers from SFI during their secondments abroad, provide on-site training on new methodologies, data interpretation and publishing.

The project has so far proven to be of unprecedented importance for SFI, as well as for the role of forest science in the national, regional and in the European research and innovation area. The number of international projects coordinated or participated by SFI has accelerated, and a number of common project proposals have been submitted to the H2020 calls. The purchase and operationalization of research equipment, realizations of secondments, trainings at SFI and workshops abroad have mostly been finalized, in which 75 % of all SFI researchers participated and benefited from. However, we have also faced several problems, including lower financing rates of recruits due to internal systemization regulations and due to a restrictive law on salaries accepted after the project negotiation has been finalized. Therefore a prolongation of time for financing the core project activities was accepted, finishing in January 2016, while the project ends on March 31<sup>st</sup> 2016.

Selected EUFORINNO impacts shall be presented in the session of the conference dedicated to the procedures aimed at accreditation of selected equipment applications, and publishing.

### **Acknowledgements**

The project EUFORINNO (RegPot No. 315982) is funded by the European Union, 7<sup>th</sup> Framework programme. Co-financing of taxes for equipment by the Slovenian Forestry Institute research programme (P4-0107) and projects, financed by the Slovenian Research Agency and the ministry, responsible for forestry, as well co-financing of the laboratory reconstruction prior to the EUFORINNO project by the ministry, responsible for science, is appreciated.

## MODERN LABORATORIES AND STANDARDISATION OF PROTOCOLS

*Marko Bajc<sup>1</sup>, Robert Robek<sup>1</sup>*

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: marko.bajc@gozdis.si

**Keywords:** project EUFORINNO, laboratory equipment, laboratory protocols, modernisation, standardisation, accreditation, scientific excellence

EUFORINNO Work Package 1 (WP1) activities focused on procurement of nineteen different types of modern laboratory equipment. The acquired equipment included both state-of-the-art core research instruments as well as general laboratory equipment. Initial list of equipment was formulated during the preparation of project application and was later refined and updated by incorporation of project partner expert suggestions. According to Public procurement regulations eighteen out of nineteen pieces of equipment were acquired through public tendering by 'bidding process after prior publication' and one by 'the best of three offers' selection process. Six public tenders were issued from December 2012 to July 2013 and successfully finalized with all equipment delivered and installed between May and December 2013.

Following the delivery, installation, basic user training and familiarization with equipment use and maintenance, SFI operators began preparing Installation and Maintenance Protocols (IMP), Operational Manuals (OM) and Standard Operational Protocols (SOP) for the new equipment. Major contributing factor for development of high quality SOPs and their standardization were trainings and secondments at project partner institutions (EUFORINNO WP2).

Based on their state-of-the-art nature, operator expertise and interest of the research community it was established that SFI would benefit the most if (1) Isotope Ratio Mass Spectrometer (EA/TC-IRMS) and (2) PALM Laser Microdissection Microscope System were selected for bringing analytical procedures performed on these two systems to accreditation-ready level. Efforts are currently underway to make all procedures and documents pertaining to the use of these two instruments compliant with SIST EN ISO/IEC 17025:2005 standard which forms the basis for accreditation. During project prolongation at least one of the selected SOPs will be subjected to audit by an external accreditation authority.

The newly acquired research equipment, combined with implementation of new and optimized SOPs and know-how acquired during trainings into research practice, has had an immensely positive impact on SFI research capabilities and an overall increase in the level of scientific excellence. New state-of-the-art core research equipment enables SFI to perform cutting edge analyses and produce highly relevant and publishable results as well as to venture into new areas of forest research. New general laboratory equipment has also contributed to a general increase in quality of laboratory work by enabling separation of processes where cross contamination is a major concern and also increased the processing capacity (throughput). The modernization of SFI laboratories and implementation/standardization of new SOPs have made SFI more competent, recognizable and attractive as a potential project partner and as a provider of analytical services for outside customers. During EUFORINNO WP1 activities SFI personnel have also gained invaluable experience in successful preparation and issuing of complex public tenders, which will undoubtedly have a beneficial impact on future public procurement activities at SFI.

## EUFORINNO METHODS TOWARDS FORMAL ACCREDITATION

*Daniel Žlindra<sup>1</sup>, Marko Bajc<sup>1</sup>, Robert Robek<sup>1</sup>*

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: daniel.zlindra@gozdis.si

**Keywords:** project EUFORINNO, final conference, accreditation, infrastructure

The value of the testing laboratory depends very much on its working procedures, structure, reliability and results. With all these words we are describing the quality of the laboratory. Laboratory quality systems were established with the idea that when testing the same sample all laboratories following such a quality system should produce the same results, within the defined precision interval. The accreditation is the process in which certification of competency, authority, or credibility is presented. The procedure for achieving accreditation is very comprehensive, labour-intensive, involves all organisational levels of the laboratory and has to be accepted and recognized by the laboratory personnel and the entire institution as a means for constantly achieving improvement and maintaining a high level of scientific excellence. During the audit an accreditation body checks the laboratory if it meets all requirements of the corresponding standard, in our case SIST EN ISO/IEC 17025. At the Slovenian Forestry Institute we formally started with the processes in January 2015 with an education seminar for a wider audience focusing on most important aspects of the standard SIST EN ISO/IEC 17025 and its implementation, followed by a workshop for a smaller group of laboratory personnel with the aim to obtain education and certification for internal auditor. All participants had successfully absolved the internal audit workshop and received an official certificate of competence to perform internal audit according to SIST EN ISO/IEC 17025. Standard operational procedures (SOP's) for EUFORINNO equipment were built upon the 16 operational manuals and know-how transfer form CP (collaborative partners). Altogether 7 SOP drafts were upgraded to proposals and four of them were harmonised following the principles of SIST EN ISO/IEC 17025 laboratory standard. Steps of internal audit and external audit by an independent body will follow.

### **Acknowledgements**

The activities were fully supported by EUFORINNO project (RegPot No. 315982).

SIST EN ISO/IEC 17025:2005: General requirements for the competence of testing and calibration laboratories. p. 28.  
International Laboratory Accreditation Cooperation: How does using an Accredited Laboratory benefit Government and Regulators?, ILAC B3:05/2011, p. 8.

## PROGRESS OF MICROSCOPY FACILITIES WITHIN EUFORINNO PROJECT

Tanja Mrak<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: tanja.mrak@gozdis.si

**Keywords:** laboratory equipment, research infrastructure, operating manual, installation and maintenance protocol, standard operating procedures, applications of light microscopy in forestry

Within Euforinno project, the outdated equipment of microscopy room was upgraded with modern and high quality microscopes Zeiss Stereo Lumar motorized dissecting microscope, Zeiss Axio Imager Z2 semi-motorized upright research microscope and PALM laser microdissection system in conjunction with Zeiss Axio Observer Z1 motorized inverted research microscope. All the instruments are operated on Zeiss ZEN software platform, while for laser microdissection special user-friendly PALM Robo software is available. For all the microscopes, operating manuals, as well as installation and maintenance protocols were prepared, and selected standard operating procedures for house rules in microscopy room, laser microdissection service, paraffin embedding of root samples, preparation of permanent semi-thin sections of mycorrhiza and measurements with ZEN software developed. The main study subjects of the facility are a) conservation biology and biotic diversity of mycorrhizal fungi, b) identification of ectomycorrhizal morphotypes, tree fine roots and lichens, c) identification and taxonomy of mycorrhizal and saprotrophic fungi, d) bioindication with ectomycorrhizal fungi, e) responses of root symbioses to environmental factors (ozone, temperature, water availability,...), f) dynamics and extent of lignification of cell walls, g) growth rings formation (on microscopy level) in dependence on environmental factors. During the project, microscopy facilities were extensively used for development of identification key for tree roots (atlas of tree roots) that encompass twelve common European temperate tree species, for which morphological and anatomical features of roots, measuring 5, 3 and 1 mm in diameter, and the most distal fine roots were studied. Based on the findings of this pilot study we will be able to expand the identification key on further tree and shrub species.

Mrak T (2015) SOP MILAB 01. Standard operating procedure. Microscopy rules. Version 1.1. Ljubljana: Slovenian Forestry Institute.

Mrak T (2015) SOP MILAB 02. Standard operating procedure. Paraffin embedding of root samples. Version 1.1. Ljubljana, Slovenian Forestry Institute.

Mrak T (2015) SOP MILAB 03. Standard operating procedure. Laser microdissection service. Version 1.1. Ljubljana, Slovenian Forestry Institute.

Mrak T (2015) SOP MILAB 04. Standard operating procedure. Preparation of permanent slides of mycorrhiza for light microscopy. Version 1.2. Ljubljana, Slovenian Forestry Institute.

Mrak T (2015) SOP MILAB 05. Standard operating procedure. Measurements of lengths on photographs taken with Zeiss microscopes. Version 1.1. (*in Slovene*) Ljubljana, Slovenian Forestry Institute.

Mrak T, Gričar J, Kraigher H (2015) Development of an atlas of fine roots of European tree species. *Rhizosphere* 4, Maastricht, the Netherlands, June 21-25. Available: <http://www.rhizo4.org/sites/default/files/All%20Posters%20Abstract%20-%20V2.pdf>, p. 233.

Mrak T, Gričar J, Kraigher H (2015) Identification of mycorrhizal partners in heterogeneous samples. In: ICOM 8 Paper and Poster Abstracts. *Eighth International Conference on Mycorrhiza, Flagstaff, Arizona, USA, August 3rd-7th, 2015*, Flagstaff: Northern Arizona University, p. 122.

Mrak T, Gričar J, Železnik P, Kraigher H (2014) Anatomical-morphological identification of fine roots of the common European tree species. In: The 1st Annual Meeting of COST Action FP1305 Biolink, Reading, 5th-6th January 2014. *What are we linking?: COST Action FP1305 Biolink: Linking belowground biodiversity and ecosystem function in European forests*. Book of abstracts. Reading: University of Reading, p. 35.

Mrak T, Kraigher H (2014) Laser microdissection: the bridge linking microscopy with molecular methods and methods of physical chemistry. In: Kraigher H, Humar M (Eds.). *Smart specialization in forestry and wood product chain: book of abstract of the Scientific Meeting Forest and Wood, Ljubljana, May 27th, 2014*. Ljubljana: Slovenian Forestry Institute, The Silva Slovenica Publishing Centre, 2014, pp. 23-24.

Mrak T, Železnik P, Gričar J, Kraigher H (2014). Towards fine root identification key of common tree species. In: Dolenc Koče J et al. (Eds.). *Book of abstracts*. Ljubljana: Slovenian Society of Plant Biology, p. 49.

## USE OF STABLE ISOTOPES IN FOREST ECOSYSTEM STUDIES: ON THE PROGRESS OF STABLE ISOTOPE LABORATORY AT THE SFI

*Saša Zavadlav<sup>1</sup>, Tom Levanič<sup>1</sup>*

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: sasa.zavadlav@gozdis.si

**Keywords:** stable isotopes, forest ecosystem, TC/EA-IRMS, IsoPrime100, vario Pyro cube

In the past three decades, the use of stable isotope techniques in forest research has greatly contributed to the understanding of biochemical, physiological and ecological processes on a spatial and time scale. Stable isotopes of light elements (i.e. hydrogen, carbon, nitrogen oxygen and sulphur) are often used as integrative tools in different fields related to forestry; from forest ecology to managements through genetics and palaeoclimatology (Ferrio et al., 2005). Natural biogeochemical processes which regulate the uptake and flow of matter cause differences in the isotopic signature of nutrient compounds due to isotopic fractionation. Thus, tracing these processes in ecosystems and determining the amount of isotopic fractionation can help identify transfer pathways and capacities (Bauer et al., 2000) important for the understanding of the dynamics of forest ecosystems responding to ever-changing environmental conditions.

The abundance of stable isotopes in samples is typically measured by an (stable) isotope ratio mass spectrometry technique (IRMS) coupled to a peripheral unit, e.g. elemental analyser, that converts a samples to a gaseous compound (CO<sub>2</sub>, N<sub>2</sub>, CO or H<sub>2</sub>). In 2013, in the frame of the EUFORINNO project (RegPot No. 315982), a continuous flow *IsoPrime100* IRMS (Isoprime, UK) coupled with a *vario Pyro cube* high-temperature elemental analyser (Elementar, Germany) was installed in the newly founded Stable Isotope Laboratory at the Slovenian Forestry Institute. The analysing system enables simultaneous elemental and stable isotope ratio determination of H, C, N, O and S in solid and liquid samples. The elemental analyser can be operated in a CNS mode (for C, N and S analyses) or in high-temperature pyrolysis mode (for H and O analyses) in a temperature range between 950 and 1450 °C. The gaseous compounds are then analysed with the IRMS for the isotope ratio determination.

Following successful installation of the analysing instruments, necessary equipment (weighing balance, reference and He gas tanks etc.), personnel training and purchase of selected certified reference materials, a variety of internal research projects were initiated to (1) calibrate the analysing instruments and test their performance, and (2) to introduce the usefulness of isotope tracer techniques to the interested research community within the institute and others. Around 3500 analyses of samples were performed so far, mainly of  $\alpha$ -cellulose isolated from tree rings, plant leaves, mosses, fungi and soil materials. The stable C and O isotopes of tree rings of oaks, spruce, chestnut tree, catalpa and wisteria helped us constrain the influence of anthropogenic pollution by burning fossil fuels, urban impacts on tree growth and the ability to survive drought period, diminished groundwater levels, and to reconstruct past climate changes. To understand the dynamics of Slovenian forests more thoroughly, isotope ratio analyses of C and N in soil profiles, roots and fungi of mixed forest stands growing in different climate areas (as an upgrade of data collected within the ManFor C.BD project) were used to reveal impacts of forest management on the soil biogeochemical processes. Additionally, along with actual elemental and isotope ratio measurements in samples, a good laboratory practice is constantly being implemented by developing standard operating procedures on the sample preparation and TC/EA-IRMS use and maintenance.

Ferrio JP, Resco V, Williams DG, Serrano L, Voltas J (2005) Stable isotopes in arid and semi-arid forest systems. *Invest. Agrar: Sist Recur For* 14 (3): 371-382.

Bauer GA, Gebauer G, Harrison AF, Högbom P, Högbom L, Schinkel H, Taylor AFS, Novak M, Buzek F, Harkness D, Persson T, Schulze ED (2000) Biotic and abiotic controls over ecosystem cycling of stable natural nitrogen, carbon and sulphur isotopes. In: Schulze ED (Ed). *The carbon and nitrogen cycling in European Forest Ecosystems*. Berlin Heidelberg: Springer, pp. 189-216.

## MEASURING ISOTOPIC COMPOSITION OF CARBON DIOXIDE IN THE AIR

*Mitja Ferlan*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

The stable carbon dioxide isotopologues analyzer – model CCIA-46 – is the latest product of Los Gatos Research, Inc. The instrument outputs  $^{13}\text{CO}_2$ ,  $^{12}\text{CO}_2$ ,  $^{12}\text{CO}^{17}\text{O}$ ,  $^{12}\text{CO}^{18}\text{O}$  in ppm and thus the isotopic ratios of  $\delta^{13}\text{C}$ ,  $\delta^{17}\text{O}$ ,  $\delta^{18}\text{O}$  in real time. The main unit for performing accurate and reliable measurements is a midinfrared quantum cascade laser using. All LGR analyzers utilize a unique laser absorption technology called Off-Axis Integrated Cavity Output Spectroscopy (OA-ICOS). This LGR-patented technique offers superior performance, value and reliability compared to Laser Absorption Spectroscopy (LAS) or cavity ringdown spectroscopy (CRDS). Features of the instrument are fast response (5Hz), portability for field measurements and insensitivity to methane and other hydrocarbons. The instrument also allow high concentration measurements via dynamic dilution system and manual sample injection via syringe. Carbon dioxide isotopologues measurements allow determination of uptake, residence time and sequestration of carbon dioxide in it's cycle in the ecosystems. The instrument could be used in net ecosystem exchange measurements via Eddy covariance method or connected to an automated soil respiration system. Instrument could also be used in labeling experiments with CO<sub>2</sub> isotopologues.

## THE PUBLISHING CENTER *SILVA SLOVENICA* AND MONOGRAPH SERIES *STUDIA FORESTALIA SLOVENICA*

*Maja Peteh*<sup>1</sup>, *Peter Železnik*<sup>2</sup>, *Tina Drolc*<sup>2</sup>, *Tom Levanič*<sup>2</sup>, *Hojka Kraigher*<sup>2</sup>

<sup>1</sup> Forestry Library - Slovenian Forestry Institute & University of Ljubljana, Biotechnical faculty, Department of Forestry and Forest Resources, Večna pot 2 and 83, SI – 1000 Ljubljana, Slovenia

<sup>2</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: maja.peteh@gozdis.si

**Keywords:** Project EUFORINNO, scientific series, publishing centre, scientific communication

*Silva Slovenica* is the publishing centre of the Slovenian Forestry Institute (SFI), founded in 1949. The publishing centre concentrates its efforts in publishing scientific monographs and co-publishes the main Slovenian scientific journal in the field of forestry, wood science and paper, but it also publishes professional publications, leaflets and brochures for promotion of certain applied and implementation results important for the civil forest service, forests and forest ecosystems.

Through the project EUFORINNO, SFI gained resources, needed for the internationalization and professionalization of its publishing centre and the monographs series. Its activities have been intensified in the last three years in several activities:

- development of a publishing system from author through technical preparation of texts with proofreading and print, which rationalises all costs of the publishing process;
- preparation of documents according to the Slovenian and international copyright law (Declaration of authorship and publication);
- set-up of the open access repository *SciVie*, which authors can use as their home repository;
- activities concerning obtaining DOI for published e-material (e.g. membership in a specific DOI organization).

Since its establishment the scientific outputs of the Slovenian forest and wood science were published in a scientific and professional monograph series *Strokovna in znanstvena dela*. With no. 122 in the year 2005 the series was renamed into *Studia Forestalia Slovenica* (ISSN 0353-6025). It is published in cooperation with the Department of Forestry and Renewable Forest Resources and Department of Wood Science and Technology at Biotechnical Faculty of the University of Ljubljana. By now 146 titles have been published, and among most prominent monographs the photo-monograph *The Virgin Forest* was published in 2014.

As a small publishing centre we are flexible, cost effective and adaptable to needs of authors, and expect to proliferate as such also after the project EUFORINNO reaches its end.

### **Acknowledgment**

The contribution was financed by the EUFORINNO project (RegPot No. 315982).

**ACTA SILVAE ET LIGNI**

*Tine Grebenc<sup>1</sup>, Maja Peteh<sup>2</sup>, Tina Drolc<sup>1</sup>, Tom Levanič<sup>1</sup>, Hojka Kraigher<sup>1</sup>*

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

<sup>2</sup> Forestry Library - Slovenian Forestry Institute & University of Ljubljana, Biotechnical faculty, Department of Forestry and Forest Resources, Večna pot 2 and 83, SI – 1000 Ljubljana, Slovenia

E-mail: tine.grebenc@gozdis.si

**Keywords:** project EUFORINNO, *Acta Silvae et Ligni*, scientific publishing, academic journals, digitalisation, open access

The scientific publications were always the key way of disseminating the research outputs and also an important stage in their evaluation and putting in frame of international importance. Recently changes in scientific communication due to European Commission activities and worldwide are encouraging researches to open science, in particular publications and available data to all communities. These shifts are also reflected in publishing activities of the Slovenian Forestry Institute (SFI). The European project EUFORINNO has offered a financial support to publishers of *Acta Silvae et Ligni* to focus and intensify activities for improvement of the journal.

*Acta Silvae et Ligni* is the only Slovenian scientific journal dedicated to publish original or review scientific and professional contributions from the fields of forests, forestry, wood science and technology, forested landscape, nature and the environment. Journal in particular supports the development of scientific and professional terminology in Slovenian language. The journal is published in cooperation of three institutions: the Slovenian Forestry Institute, Department of Forestry and Renewable Forest Resources and Department of Wood Science and Technology at Biotechnical Faculty of the University of Ljubljana. In 2013, with no. 100, its current title has been introduced with the purpose of boosting the internationalization of the journal. Several other changes have also been done on graphical design, home page, policy strategy, open access to bring the journal closer to all levels of readers, and a fully electronic submission system. The journal is under evaluation within two major databases, Web of Science and Scopus. Also the process of obtaining DOI for all articles has been lounged (e.g. membership in a specific DOI organization).

In addition the digitalization process of 643 articles from the Zbornik gozdarstva in lesarstva was conducted and entered into the repository *SciVie*.

With all these activities we try to enable use and re-use of new and old materials that might otherwise not be readily accessible, and we thus support the advancement of forest research and development.

**Acknowledgment**

The contribution was financed by the EUFORINNO project (RegPot No. 315982). Publishing of journal *Acta Silvae et Ligni* is co-financed by the Slovenian Research Agency.

## **SCIVIE – SCIENCES DE LA VIE FROM THE PAST AND FOR THE FUTURE**

*Maja Peteh<sup>1</sup>, Irena Rebov<sup>2</sup>, Tina Drolc<sup>1</sup>, Žiga Lipar<sup>1</sup>, Tom Levanič<sup>2</sup>, Hojka Kraigher<sup>1</sup>*

<sup>1</sup> Forestry Library - Slovenian Forestry Institute & University of Ljubljana, Biotechnical faculty, Department of Forestry and Forest Resources, Večna pot 2 and 83, SI – 1000 Ljubljana, Slovenia

<sup>2</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: maja.peteh@gozdis.si

**Keywords:** project EUFORINNO, repository, *SciVie*, digitalisation, open access, life sciences

Changes in scientific communication due to European Commission activities and worldwide are facing directions to open science, in particular publications and available data to all communities. These shifts are reflecting also in publishing activities of the Slovenian Forestry Institute (SFI). With the financial support of the project EUFORINNO we have started the open access repository *SciVie*.

The repository *SciVie* (deriving from ‘Science de la Vie’) with its open access materials provides more efficient re-use of past research, promotion of research results, researchers and the institution itself including the possibility of increased citations and recognition of scientific works, training future scientists for open access, increasing the number of potential users and giving the general public the opportunity of improving their lives resulting from the research findings. The repository *SciVie* was launched on a base of the EPrints 3 software as multi-institutional repository in the fields of natural and life sciences. It is intended for storage of articles, monographs, databases and other works, with the aim of meeting the requirements of EU co-financers. Therefore it has been entered in the databases OpenDOAR, RAOR and OpenAir. Currently several institutions already contribute to *SciVie*, including the Slovenian Forestry Institute, Agricultural Institute of Slovenia, National Institute of Biology and Biotechnical Faculty of the University of Ljubljana, Department of Wood Science and Technology. Big efforts were put into developing the policy strategy and copyright requirements that meet the Slovenian and EU policy, into its graphical design and home page.

Activities concerning promotion of open access have been organized at all participating institutions as well as in the southeast Europe.

In addition the three stage digitalization process for already published works (dating from the establishing SFI in 1950’) was set up. Firstly 643 articles from the Zbornik gozdarstva in lesarstva (now *Acta Silvae et Ligni*) were digitalized and entered into *SciVie*. In the second stage 324 study reports were digitalised which were considered as grey literature and the only copy (usually) only exists in the Forestry library. That task has been a challenge even for the chosen digitalizing company, since materials were on a different paper formats, old or damaged paper and printed or typed in a diverse quality. Special challenge has been the digitalization of supplementary maps and other outsized material and finding a way to disseminate digitalized copies to public (via *SciVie*) as not too large files. The third stage, the digitalization of a set of study reports and other similar material published by SFI, is under way.

With all these activities we try to enable use and re-use of new and old materials that might otherwise not be readily accessible, and we thus support the advancement of forest research and development.

### **Acknowledgment**

The contribution was financed by the EUFORINNO project (RegPot No. 315982).

# POSTERS

in alphabetical order of first authors

## LEAF UNFOLDING VARIABILITY IN CLONAL SEED ORCHARD OF *Fraxinus angustifolia* Vahl

Ivan Andrić<sup>1</sup>, Davorin Kajba<sup>1</sup>

<sup>1</sup> Faculty of Forestry University of Zagreb, Svetošimunska 25, 10 000 Zagreb, Croatia

E-mail: davorin.kajba@zg.t-com.hr

**Keywords:** narrow leaf ash, flushing phases, phenoform group

In Croatia, narrow-leaved ash (*Fraxinus angustifolia*) is distributed in the Pannonian lowland area, along the rivers with pedunculate oak (*Quercus robur*) in mixed stands. Greatest portion of its genetic variability is localized in the region of Sava river (area of 30 000 ha) where in 2005 established clonal seed orchard with 56 clones (Forest Range Office Nova Gradiška). The grafts were planted with 4 × 4 m spacing in the area of 3.50 ha. Clonal seed orchard have been regularly maintained by pruning and with other agrotechnical treatments from the moment of establishment. Pomotechnical treatments are used and successful pomology techniques is maintaining the balance between the vegetative and generative activity.

Phenological seasonal rhythm has recently receives importance and with better understanding of leaf phenophases we will be able to more precisely define response to climate change. Through the period of three years (2012, 2014, 2015) it was observed phenological clonal differences in monitoring of flushing phases on four ramets per clone on which we based our strategy to determine interclonal and intraclonal variability. The start date of the leaf unfolding stage is defined as the point at which the entire leaf blade and leaf stalk were visible (Phase 2). Each year, observation began before any buds began to break (10<sup>th</sup> of March), and was made every seven days until the 1st of June.

Analysis of variance showed statistically significant difference between the studied genotypes for each year of investigation for the parameters period of leaf development and beginning of leafing ( $F = 6.89$ ,  $F = 7.15$ ,  $F = 4.91$ ,  $Pr < 0.0001$ ). There is a statistically significant difference over the years ( $F = 539.21$ ) and for the interaction clone × year ( $F = 2.44$ ,  $Pr < 0.0001$ ). The results of k-means clustering of the clones according to their leaf unfolding clearly classified clones into three groups (early, intermediate and late flushing). Each of the phenoform groups are also an important characteristic related to the pests and insects and to the occurrence of the late spring frost. For the early flushing group, the mean value for the period of leaf development were in average from 24 days in 2012, 25 days in 2014 and 20 days in 2015. The intermediate group required an average of 27 days in 2012, 26 days in 2014 and 19 days in 2015. For the late flushing group, the mean value were in average from 31 days in 2012, 26 days in 2014 and 21 days in 2015. Among early, intermediate and late flushing group of clones due to the time to complete the leaf development for each year of observation there was no statistically significant difference ( $p = 0.238$ ) but it was between the years ( $p = 0.017$ ). In this period seed orchard was also continuously monitored and checked for the presence of *Chalara fraxinea*. The presence of *Chalara fraxinea* was not confirmed, and on the clones there were no symptoms that would indicate it.

Statistically significant differences were obtained between genotypes in clonal seed orchards for period of leaf development and leaf unfolding. There is a statistically significant difference over the years and significant differences in the interaction between the clones × years, which means that analyse variable for phenoform group over the year do not behave the same. Investigated 54 clones were divided into three groups depending on the beginning of leafing. With the results gathered from this monitoring we are able to determine and classify clones into early, intermediate and late flushing groups, which has a high practical value not only for clonal seed orchard managing and maintaining, but also for the production of quality forest reproductive material and their distribution as very important adaptive trait for the future.

### Acknowledgments

This research has been financially supported by the Croatian Science Foundation through the research project FRAXINPRO (IP-11-2013).

## PHENOLOGICAL VARIABILITY OF EUROPEAN BEECH (*Fagus sylvatica* L.) IN THE INTERNATIONAL PROVENANCE TRIAL

*Dalibor Ballian*<sup>1</sup>, *Benjamin Jukić*<sup>1</sup>, *Davorin Kajba*<sup>2</sup>, *Georg von Wüehlich*<sup>3</sup>

<sup>1</sup> Šumarski fakultet Univerziteta u Sarajevu, Bosnia and Herzegovina

<sup>2</sup> Šumarski fakultet Sveučilišta u Zagrebu, Croatia

<sup>3</sup> Federal Research Institute for Rural Areas, Forestry and Fisheries (vTI), Germany

E-mail: balliandalibor9@gmail.com

**Keywords:** European Beech, provenance trial, phenology, coloration and winter leaf retention

Common beech is the most important species of forest trees in Bosnia and Herzegovina. This research provides the results of investigation into the phenological variability of 22 common beech provenances in the international trial near Kakanj, both for the properties of autumn coloration and winter leaf retention. The experiment included eight provenances from Bosnia and Herzegovina, four from Germany, three from Serbia, two from Croatia, Romania and Switzerland each and one from Hungary. The experiment was established in a randomized block system with three repetitions.

The goal of this research was to establish whether there were differences in variability between the investigated provenances and what features would be important for future management and regeneration of beech forests, as well as for increased production and adaptability.

Based on the investigated phenological properties, statistically significant variability was found between all the studied provenances. The established differences included the beginning, duration and end of particular phenophases. There are considerable statistically significant differences in flushing among the studied provenances. Provenances from Croatia and Bosnia as a rule flush earlier than foreign provenances. April 10 can be considered as the beginning of common beech vegetation in this international trial, because this is the date on which bud opening occurred in twelve out of 22 provenances.

The analysis of leaf colour of different common beech provenances revealed variations in the presence of different groups of colours with varying shades in a particular period. Current differences in winter leaf retention indicate that 65.92 % of the plants in the entire experiment do not retain the leaves. The highest proportion of leaf non-retention, amounting to 91.05 %, was manifested by the provenance from Germany (Hoellerbach). It can be concluded that common beech is genetically differentiated according to ecological and vegetation areas and that it shows different morphological and physiological characteristics from different provenance areas. This proves that morphological and physiological properties of common beech are influenced by a number of factors and that each of the investigated properties is determined by a large number of genes.

## THE IMPACTS OF FOREST FIRES IN THE CURRENT STATE OF FOREST RESOURCES IN KOSOVO

*Thaci Bashkim<sup>1</sup>, Bojaxhi Faruk<sup>2</sup>*

<sup>1</sup> Ministry of Agriculture, Forestry and Rural Development, Republic of Kosovo

<sup>2</sup> Kosovo Forest Agency, Republic of Kosovo

E-mail: bashkimt2002@yahoo.com; faruk.bojaxhi@rks-gov.net

**Keywords:** forest fires, economic damages, vulnerability, preventing measures

Kosovo is situated in the central Balkan peninsula. It has an area of 10,887 square kilometer and an overall forest area of 481,000 hectare. Forests and pastures in Kosovo occupy around 50 % of the whole Kosovo territory. Based on the National Forest Inventory carried out on 2012, the total standing volume is 46.3 Million m<sup>3</sup>. Natural broadleaves forests cover 90 % of the whole forest area, while conifer forests cover about 7 % represented by silver fir (*Abies alba* Mill.), norway spruce (*Picea abies* H. Karst.) and pine species. More than 60 % of the forests are located in various altitudes ranging from 600 m to 1000 m. The aim of the study is to assess the vulnerability of forest fires, their causes as well as measures to prevent them. Kosovo forests during last years have been affected by severe forest fires causing many damages with a high economic bill. The most severe fire's years were 2000, 2004, 2007 and 2012, burning about 10,955 ha, with an economic damage about 4.65 Million €.

## RETREATING SESSILE OAK FOREST WITH IMPROVING VITALITY – INCLUDING TREE MORTALITY IN VITALITY ASSESSMENT

Imre Berki<sup>1</sup>, Ervin Rasztoivits<sup>1</sup>, Norbert Móricz<sup>2</sup>, Gábor Veperdi<sup>3</sup>, László Kolozs<sup>4</sup>

<sup>1</sup> Institute of Environmental and Earth Sciences, University of West Hungary, Sopron, Hungary

<sup>2</sup> Forest Research Institute, National Agricultural Research and Innovation Centre, Sárvár, Hungary

<sup>3</sup> Institute of Forest Resource Management and Rural Development, University of West Hungary, Sopron, Hungary

<sup>4</sup> National Food Chain Safety Office, Budapest, Hungary

E-mail: rasztovits.ervin@emk.nyme.hu

**Keywords:** *Quercus petraea*, drought, mortality, stand density, forest monitoring

The drought induced vitality loss of sessile oak (*Quercus petraea*) has been continuously observed in Hungary for more than three decades. Although stand density is an important indicator of the stand stability, the decreasing stand density as a consequence of drought induced mortality is not considered in most of the monitoring methods (e.g. ICP forest monitoring network). We introduced a novel health assessment method based on the vitality of the living trees and on the relative stand density, expressed as the ratio of the current density to the fully stocked density. Stands for the assessment were selected along a climatic transect from the humid region in SW-Hungary to the continental-semiarid region in NE-Hungary where no forest intervention was applied at least during the last three decades. The results showed that the calculated health status of the sessile oak stands are considerably below the ones measured by the ICP, lying between 70-90 % in SW Hungary and below 50 % close to its xeric limit.

## GENETIC ANALYSIS OF NATIVE EUROPEAN BLACK POPLAR (*populus nigra* L.) POPULATIONS IN SLOVENIA AND CROATIA BY USING SSR MARKERS

Gregor Božič<sup>1</sup>, Marjana Westergren<sup>1</sup>, Davorin Kajba<sup>2</sup>, Andreja Ferreira<sup>1</sup>, Hojka Kraigher<sup>1</sup>, An Vanden Broeck<sup>3</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI-1000 Ljubljana, Slovenia

<sup>2</sup> Faculty of Forestry University of Zagreb, HR-10000 Zagreb, Croatia

<sup>3</sup> Research Institute for Nature and Forest, BE-9500 Geraardsbergen, Belgium

E-mail: gregor.bozic@gozdis.si

**Keywords:** *Populus nigra*, genetic resources, genetic diversity, introgression, microsatellite loci

The European black poplar (*Populus nigra* L.) is one of the most important tree species of alluvial forests in Europe. In Slovenia and northern Croatia, ecosystems, where European black poplar is the keystone species, are fragmented and endangered mostly because of river regulations, pollution, urbanisation and felling of vital trees for wood. European black poplar is also an indicator species for the vitality and intactness of alluvial ecosystems. To inventory genetic diversity and structure of the remaining natural populations, a study using nuclear microsatellites was conducted.

We analysed genetic diversity and structure of 10 native black poplar populations (200 trees) along the main river systems across Slovenia and Croatia (rivers Soča, Sava and Mura) and in the submediterranean using 6 microsatellite loci (WPMS16, WPMS20, WPMS14, PMGC14, WPMS09, WPMS18). Eleven common cultivated clones of *Populus x canadensis* were also included in the analysis in order to assess the level of introgression of genes of the American Eastern cottonwood, *P. deltoides*, into the gene pool of native black poplar. Standard methods using programmes GenAlEx, SpaGeDi and Structure were used to obtain estimates of genetic diversity and genetic structure including presence of hybrids.

Three genepools were observed for our dataset, mainly following the river systems (Soca, Mura, Sava with two genepools). No clear accumulation of allelic richness nor gene diversity downstream neither for Sava nor Mura was observed. In natural populations, the introgression of *P. deltoides* genes was very low. The results indicate that the gene pool of the remaining *P. nigra* populations maintains high genetic connectivity along rivers even if fragmented today. The obtained results can be used to further develop dynamic *P. nigra* conservation in endangered forest ecosystems at regional and national levels.

### Acknowledgments

This study took place within the framework of research tasks of project **V4-1438** and Research Programme P4-0107.

## IMPLEMENTING FOREST GENETIC MONITORING PLOTS

*Domen Finžgar<sup>1</sup>, Melita Hrenko<sup>1</sup>, Gregor Božič<sup>1</sup>, Hojka Kraigher<sup>1</sup>*

<sup>1</sup>Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: domen.finzgar@gozdis.si

**Keywords:** forest genetic resources, genetic diversity, monitoring, Smolarjevo, Pri studencu

Within the LIFE GENMON project, two forest genetic monitoring (FGM) plots were installed in Slovenia in 2015. Such monitoring might be a possible tool for gathering information, whether the forest ecosystem can adapt to the future changes of its environment (Namkoong et al. 1996, Konnert et al. 2011). FGM is going to be conducted for two species - *Fagus sylvatica* L. and *Abies alba* Mill. The sites were considered based on the pre-prepared LIFE GENMON protocol drafts, meteorological and environmental data availability, site conditions, local forestry initiative, legal status of selected forests and ownership. Upon several candidate plots, two seed stands: "Smolarjevo" and "Pri studencu" were chosen. The poster presents main reasons for this decision, as well as rich history of forest management on both FGM plots. The pre-prepared LIFE GENMON protocol drafts for FGM seem to be well considered so far and only minor changes need to be discussed within LIFE GENMON partners when preparing the final protocol.

### Acknowledgements

Research Program P4-0107, Public Forestry Service, CRP V4-1438, EUFORGEN WG on Genetic monitoring and LIFE GENMON project, financed by LIFE+, MOP, MKGP and SFI.

Konnert M, Maurer W, Degen B, Kätzel R (2011) Genetic monitoring in forests - early warning and controlling system for ecosystemic changes. *JForest* (4): 77-81.

Namkoong G, Boyle T, Gregorius HR, Joly H, Savolainen O, Ratnam W, Young A (1996) Testing criteria and indicators for assessing the sustainability of forest management: genetic criteria and indicators. Working paper No. 10, Bogor, Centre for International Forestry Research (CIFOR): 12pg.

## TESTING OF SALT TOLERANCE ASSOCIATED GENES IN SERBIAN POPLAR CLONES

*Vladislava Galović<sup>1</sup>, Saša Orlović<sup>1</sup>, László Szabados<sup>2</sup>, Imma Perez<sup>2</sup>*

<sup>1</sup> Institute of Lowland Forestry and Environment, Antona Cehova 13d, 21000 Novi Sad, Serbia

<sup>2</sup> Institute of Plant Biology, Biological Research Center, Temesvári krt. 62., 6726-Szeged, Hungary

E-mail: vladislava.galovic@gmail.com

**Keywords:** poplar, gene expression, stress inducible genes, quantitative RT-PCR

Trees from Pannonian region are exposed to a combination of environmental stress conditions, especially high salinity. Poplar is extensively used for forestation in this region due to its importance in carbon sequestration, bioremediation, nutrient cycling and wood production. Rapid growth, facile transformation and small genome size makes poplar an ideal model for basic and applied research targeting stress responses in woody plants.

This research work employs functional genomics strategies to analyze oxidative stress responses of poplar woody plant species in order to improve its tolerance to extreme environmental conditions. The experiment was started with two poplar clones (*P. deltoides* and *P. x euramericana*), with different tolerance level, exposed to salt stress where different concentration of NaCl as a stress agent with a different time point of stress were monitored. Transcript levels of stress-induced marker genes, transcription factors and other regulatory genes were analyzed. In order to test several stress inducible genes of different biosynthetic pathways on particular poplar clones quantitative RT-PCR was used. The isolation of total RNA from poplar tissue, derived from control and stress samples was done using TRIzol method, followed with first strand DNA synthesis. Successful cDNA synthesis was confirmed by amplification of actin gene (PtACTIN9 *Populus trichocarpa* homolog) for both species as endogenous gene of choice (internal standard). The Ct value of the genes of interest were tested on sampled cDNA by quantitative RT-PCR using SYBR green. Different level of expression has been occurred for different stress induced genes. The best induction was detected in the gene belonging to the dehydrin protein family with lysine rich segments in its sequence. It is a regulatory gene which amino acid sequence shows homology with group II LEA (late embryogenesis abundant) proteins. Also responds to osmotic stress, ABA, dehydration and inhibits E.coli growth while overexpressed. The most important segment of this work represents getting successful strategy for approaching the climate change problem with functional genomic tools using qPCR for testing gene expression in order to obtain the first applicable insights into the future protection of poplar species in the area of Srem, Vojvodina region, Serbia.

### Acknowledgements

This work were supported by the project of Ministry of Education, Science and Technological Development Republic of Serbia No III43002 and by the project funded by Autonomous Province of Vojvodina, Provincial secretariat for science and technological development 2015.

## MOLECULAR ANALYSES OF ITS rDNA OF THREE AUTOCHTHONOUS FUNGAL SPECIES FROM MOUNTAIN FORESTS IN SERBIA

*Vladislava Galović<sup>1</sup>, Maja Karaman<sup>2</sup>, Saša Orlović<sup>2</sup>, Bojan Šeguljev<sup>2</sup>, Hojka Kraigher<sup>3</sup>, Marko Bajc<sup>3</sup>*

<sup>1</sup> Institute of Lowland Forestry and Environment, University of Novi Sad, Serbia

<sup>2</sup> Department of Biology, Faculty of Sciences, University of Novi Sad, Serbia

<sup>3</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: vladislava.galovic@gmail.com

**Keywords:** mountain forest, macrofungi, diversity, molecular taxonomy, molecular phylogeny

Fungi and mushrooms represent an essential structural element of forests ecosystems. In the frame of national project of biodiversity examination in mountain forests of Serbia during three investigation years (2011-2013), autochthonous macrofungal diversity have been examined on the following areas: Tara Mt (Mitrovac), Kopaonik Mt (Metode) and Stara planina Mt (Vidlič). The most frequently 3 species found in all three areas were chosen for analyses, including selected isolates of each, namely 6 isolates of *Fomitopsis pinicola* (Sw.) P. Karst., 8 isolates of *Polyporus varius* (Pers.) Fr. and 3 isolates of *Hymenopellis radicata* (Relhan) R.H. Petersen.

Since molecular methods have been contributed to a great extent to the taxonomic and phylogenetic studies of fungi in the last decades and DNA barcoding has become a standard method in species identification, the aims of this study were: 1) confirmation of selected taxa, previously determined by classical morphological procedures, using molecular methods; 2) evaluation of their phylogenetic origin using sequence analysis of the ITS1-5.8S-ITS2 region rDNA, by comparing it with sequences from the GenBank® database using BLAST. UNITE sequences from databases, that upon BLAST search showed concordance of over 97 %, were used for the formation of the cluster tree dendrogram constructed using RAxML (Randomized Axelerated Maximum Likelihood) software.

Based on the results obtained, the ITS rDNA region proved to be adequate bar-coding marker for molecular determination of tested fungal species and construction of phylogenetic trees.

According to sequence-based analysis from dendrogram constructed for *F. pinicola* clade, autochthonous mushrooms species have a common origin with European species. Based on the sequences of *P. varius* two distinct clades can be singled out, of which one consists of species *Polyporus varius*, *Polyporus melanopus* and *Royoporus badius* what is in correspondence with morphologically defined group *Melanopus*. Since all three sequences of autochthonous *H. radicata* grouped in one of two distinct subclusters, phylogenetic relatedness of the isolates can be concluded as well as the separation from European species sequences of the same fungal species.

### Acknowledgements

This work was supported by the project of Ministry of Education, Science and Technological Development Republic of Serbia No III43002.

## HYPOGEOUS FUNGI DIVERSITY AND ECOLOGY IN SE EUROPE

*Tine Grebenc*<sup>1</sup>, *Maria P. Martin*<sup>2</sup>, *Marcelo Sulzbacher*<sup>3</sup>, *Mitko Karadelev*<sup>4</sup>, *Gian Maria Niccolo Benucci*<sup>5</sup>, *Dalibor Ballian*<sup>6</sup>, *Tomislav Lukic*<sup>7</sup>, *Jelena Lazarević*<sup>8</sup>, *Hojka Kraigher*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI-1000 Ljubljana, Slovenia

<sup>2</sup> Jardín Botánico, CSIC, Plaza de Murillo 2, 28014 Madrid, Spain

<sup>3</sup> Universidade Federal de Pernambuco, Departamento de Micologia/CCB, Av. Prof. Nelson Chaves, s/n, CEP: 50670-901, Recife, PE, Brazil

<sup>4</sup> Mycological Laboratory, Institute of Biology, Faculty of Natural Science and Mathematics, Cyril and Methodius University, Arhimedova 5, 1000 Skopje, The Republic of Macedonia

<sup>5</sup> Department of Plant, Soil and Microbial Sciences, Michigan State University, 1066 Bogue Street 48825 East Lansing, MI, USA

<sup>6</sup> University of Sarajevo, Faculty of Forestry, Zagrebačka 20, BA-71000 Sarajevo, Bosnia and Herzegovina

<sup>7</sup> no address

<sup>8</sup> University of Montenegro, Biotechnical faculty, Mihaila Lalića 1, 81000 Podgorica, Montenegro

**Keywords:** truffles, distribution, species concept, cultivation perspectives

In Europe there is over 50 genera of hypogeous fungi with best known genus *Tuber* (true truffles), represented with over 30 species published for central and western Europe. Truffles were only recently popularised in the SE Europe, namely in the Balkan Peninsula and their presence, diversity and ecology in the region remains understudied. We aimed to analyse the diversity of true truffles and to certain level also false truffles (all non-*Tuber* hypogeous fungi) in selected countries in the SE Europe and compare it with the rest of Europe and globally.

Using specialised purposive oriented tracing approaches we collected over 2000 collections of hypogeous fungi, all ecologically and morphologically analysed. The less known genera were also characterised using molecular markers.

The diversity of hypogeous fungi in the SE Europe is comparable to other parts of Europe (Marjanović et al. 2010) with a high number of endemic genotypes, among which at least some can be identified as separate taxa following the Species hypothesis approach (Köljalg et al., 2013). The distribution and ecology of selected species is pointed out, so as the importance of commercial truffle species in the region.

### Acknowledgements

The research project was financed in part by EUREKA E!3835 EUROAGRI- CULTUBER “Improvement of truffle cultivation via novel quality control, soil analysis and inoculation methods” and by the SYNTHESYS grant ES-TAF-1729 - financial support of the EC’s Programme “Structuring the European Research Area”. The partners from the Slovenian Forestry Institute were co-financed by the Ministry of Higher Education, Science and Technology through Research Programme P4-0107.

Marjanović Ž, Grebenc T, Marković M, Glišić A, Milenković M. 2010. Ecological specificities and molecular diversity of truffles (genus *Tuber*) originating from mid-west of the Balkan Peninsula. *Sydowia* 62(1): 273-291.

Köljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Grebenc T, et al. (2013). Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology* 22: 5271–5277.

## TYPES OF ECTOMYCORRHIZA IN THE 34 YEARS OLD *Pinus sylvestris* L. SEED PLANTATION IN THE LOWLAND FOREST SITE "MURSKA ŠUMA"

*Melita Hrenko*<sup>1</sup>, *Gregor Božič*<sup>1</sup>, *Tine Grebenc*<sup>1</sup>, *Anita Mašek*<sup>1</sup>, *Hojka Kraigher*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI-1000 Ljubljana, Slovenia

E-mail: melita.hrenko@gozdis.si

**Keywords:** *Pinus sylvestris*, *Quercus robur*, forest plantation, ectomycorrhiza, monitoring

The objective of this study was to evaluate the types of ectomycorrhiza of *Pinus sylvestris* L. seed plantation in SE part of Slovenia at 158 m a.s.l. and establish a longterm monitoring of ectomycorrhizal types in new developing forest plantation of *Quercus robur* L.. Samples were taken in April 2015. *Quercus robur* was sampled in forest nursery Polana in Banuta site and *Pinus sylvestris* in its former plantation in the Murska šuma site. At each of two sites soil cores were taken and the soil structure was analysed. Roots from the both species were sampled in different sites. After washing, all roots were differentiated into non-woody-plant roots, non-ectomycorrhizal, non-turgescent and different vital ectomycorrhizal morphotypes. Morphological and anatomical characteristics of mycorrhiza is under study. Ectomycorrhizal roots will be identified by using classical ITS-PCR and sequencing.

### **Acknowledgments**

This study took place within the framework of the Research Programme P4-0107.

## SOIL PROPERTIES IN RELATION TO SELECTIVE LOGGING IN BEECH AND FIR FOREST STANDS - CASE STUDY AT BJELAŠNICA MOUNTAIN

*Emira Hukić<sup>1</sup>, Primož Simončič<sup>2</sup>, Hamid Čustović<sup>3</sup>*

<sup>1</sup> Faculty of Forestry, University of Sarajevo, Zagrebačka 20, BiH – 71000 Sarajevo, Bosnia and Herzegovina

<sup>2</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

<sup>3</sup> Faculty of Agriculture and Food Science, University of Sarajevo, Zmaja od Bosne 8, BiH – 71000 Sarajevo, Bosnia and Herzegovina

E-mail: e.hukic@sfsa.unsa.ba

**Keywords:** SOM, topsoil, site factors measurements, Mt Bjelašnica

Quality of forest soils is associated with the range of stand disturbance due to applied forest management practice. Selective logging, which is characteristic for mixed beech and fir forests (*Abieti-fagetum dinaricum s. lat.*), rises globally in its intensity in regard to the growing needs for wood products and energy resources. Understanding how the soil quality alters relative to selective logging operations in a successional cycle will help determine sustainability of forest management system. Mixed beech and fir forests are covering large areas along the Dinaric mountain range where limestone and dolomites occur as predominant geological substrates. In Bosnia and Herzegovina, they comprise more than 50 % of all high forests and they are therefore considered as the most important forest types. In addition to the principles of the forest management strategy which imply maintenance of their structural elements, we are investigating soil chemical and microbiological properties associated with alternations in forest structure, micrometeorological parameters, and light conditions in a different chronosequences after logging. Regarding the intensity and temporal premises, the results should answer the question of indirect influence of logging on forest soil organic matter quality, storage ability for the organic carbon and nitrogen, nutrient ability, microbial properties etc. Soil organic matter is vitally important for ecosystem functions on shallow calcareous soils and further critical for maintaining natural regrowth and forest ecosystem sustainability. In that view the study attempts to valorize the forest management system using soil quality parameters.

## SEASONAL VARIATION OF ECTOMYCORRHIZAL COMMUNITY FROM MATURE POPLAR PLANTATION

*Marina Katanić<sup>1</sup>, Saša Orlović<sup>1</sup>, Tine Grebenc<sup>2</sup>, Marko Bajc<sup>2</sup>, Branislav Kovačević<sup>1</sup>, Milan Matavuly<sup>3</sup>, Hojka Kraigher<sup>2</sup>*

<sup>1</sup> University of Novi Sad, Institute of Lowland Forestry and Environment, Antona Čehova 13, 21000 Novi Sad, Serbia

<sup>2</sup> Slovenian Forestry Institute, Večna pot 2, 1000 Ljubljana, Slovenia

<sup>3</sup> European University, Faculty of Pharmacy, Trg mladenaca 5, 21000 Novi Sad, Serbia

E-mail: marinakatanic44@gmail.com

**Keywords:** *Populus alba* L., ectomycorrhizal diversity, anatomophic characterization, molecular identification, seasonal dynamics

Ectomycorrhizal communities are rarely studied at the seasonal basis. For this reason we analysed the seasonal variation of the ectomycorrhizal community from mature twenty years old white poplar plantation growing on native poplar site. Using anatomical and morphological identification in combination with molecular approach, 29 distinct ectomycorrhizal types mainly belonging to Basidiomycota were identified. The seasonal differences were most profound between winter and spring on one side and summer and autumn on the other side. Number of ectomycorrhizal types and percentage of vital ectomycorrhizal roots were the highest in winter and spring. The diversity indices of ectomycorrhizae and number of fine roots from the studied poplar plantation were similar to other studies on poplars. Persistent dominance of types from family Inocybaceae was observed at studied site. While short distance exploration type dominated overall ectomycorrhizal community in all seasons, the abundance of ectomycorrhizal root tips belonging to the medium distance exploration type was significantly higher in spring in comparison with autumn and winter.

### Acknowledgements

The study was financed by the Slovenian Research Agency through Programme P4-0107, Scholarship Ad futura and project III43007 of Ministry of ESTD of the Republic of Serbia.

## POSSIBILITIES FOR THE USE OF LASER VIBROMETRY IN THE MANAGEMENT OF INSECT PESTS

*Andreja Kavčič*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: andreja.kavcic@gozdis.si

**Keywords:** project EUFORINNO, final conference, 7FP, invasive insect pests, Laser vibrometry

Nowadays, invasive alien species present one of the main threats to natural and semi-natural ecosystems worldwide. In addition to global trade and tourism, which have escalated the spread of alien species to unprecedented levels, climate change is becoming an increasingly important factor as it further facilitates expansion of species' natural ranges and their spread to new areas. Until recently, control of pests was based mainly on the use of chemical pesticides. Since these often present risk to non-target organisms, researchers seek alternative methods for pest management. Among these, biological control through the use of the species's natural enemies has attracted considerable attention, and currently shows potential to play a pivotal role in the sustainable management of invasive species in the future.

Among insects, some of the most important pests come from the family of plant-dwelling stink bugs (Heteroptera, Pentatomidae), which feed on a wide variety of host-plants. Given the high amount of damage that these species cause mainly in agriculture, it is not surprising that they are subject to extensive research. Additionally, the family Pentatomidae is important also for certain predatory species that it includes. However, these are less well known and have drawn attention of the scientific community only recently, namely because they show high potential as biocontrol agents. It is now well established that in Pentatomidae species vibrational communication represents a communication channel that is vital for successful reproduction for it enables sexual partners to recognise each other, to meet and to mate. Studies have shown that males and females emit species- and sex-specific substrate-borne vibrational signals, which are produced by dorso-ventral vibrations of the abdomen and have distinct temporal, frequency and amplitude characteristics. Vibrations of the abdomen are transmitted through legs and induce vibrations of the substrate. In addition to this, research have shown that vibrational signals play an important role also in interspecific interactions, such as interactions between prey and their natural enemies. It has been shown that certain parasitoids exploit vibrations produced by their prey as cues for localization of the prey. Similarly, in some prey species vibrations produced by predators evoke escape reactions. Given the importance of vibrational communication in a variety of economically important insects and implication of vibrational signals in predator – prey interactions, a wide variety of research has focused on vibrational communication within and between species, trying to better understand their biology and gain new insights, which would help in obtaining new tools for effective control of insect pests.

Detection and measurement of vibrational signals is possible by different bioacoustic detection methods. However, Laser vibrometry has proven especially suitable for registration of vibrational signals of small insects. This is a novel, non-contact method based on the frequency shift of the laser beam due to the motion of the surface (Doppler effect). The amplitude and frequency of vibrational signals are calculated from the frequency change of the test beam, which is obtained from the interference of the test and reference beams. The main advantages of this method are that it does not load the substrate, has a wide range of frequency sensitivity, and enables measurement of vibrational signals on small and delicate natural as well as artificial substrates. Laser vibrometry has significantly extended the area of possible

research of insect pests. Current research focus on vibrational repertoires of pest species and their potential predators, signal transmission and the nature of vibrational signals on different substrates, and the effect of heterospecific vibrational signals on insect development and behaviour. Together with the latest study showing that Laser vibrometry could be used also as a diagnostic tool for detection of wood-boring insect pests, this method shows high potential in the development of novel approaches to the management of insect pests.

- Castellanos I, Barbosa P (2006) Evaluation of predation risk by a caterpillar using substrate-borne vibrations. *Anim Behav*, 72: 461-469.
- Čokl A (2008) Stink bug interaction with host plants during communication. *J Insect Physiol* 54: 1113-1124.
- Čokl A, Millar JG (2009) Manipulation of insect signaling for monitoring and control of pest insects. In: Ishaaya I., Horowitz A.R. (Eds.). *Biorational Control of Arthropod Pests: Application and Resistance Management*. Dordrecht, Heidelberg, London, New York, Springer: pp. 279-316.
- Čokl A, Virant-Doberlet M (2003) Communication with substrate-borne signals in small plant-dwelling insects. *Annu Rev Entomol* 48: 29-50.
- Čokl A, Žunič A, Virant-Doberlet M (2011) Predatory bug *Picromerus bidens* communicates at different frequency levels. *CEJB* 6: 431-439.
- De Clercq P (2000) Predaceous stinkbugs (Pentatomidae: Asopinae). In: Schaefer C.W., Panizzi A.R. (Eds.). *Heteroptera of Economic Importance*. Boca Raton, Florida, CRC Press: pp. 737-789.
- Horowitz AR, Ellsworth PC, Ishaaya I (2009) Biorational pest control – an overview. In: Ishaaya I., Horowitz A.R. (Eds.). *Biorational Control of Arthropod Pests: Application and Resistance Management*. Dordrecht, Heidelberg, London, New York, Springer: pp. 1-20.
- Kavčič A (2014) Recognition of predators by vibrational signals in the stink bug *Nezara viridula* (L.) (Doctoral Dissertation). University of Ljubljana, Slovenia.
- Kavčič A, Čokl A, Laumann RA, Blassioli-Moraes MC, Borges M (2013) Tremulatory and abdomen vibration signals enable communication through air in the stink bug *Euschistus heros*. *PLoS ONE* 8, 2: e56503. doi:10.1371/journal.pone.0056503
- Knight KMM, Gurr GM (2007) Review of *Nezara viridula* (L.) management strategies and potential for IPM in field crops with emphasis on Australia. *Crop Prot* 26: 1-10.
- Kuštor V (1989) Activity of muscles of the vibration producing organ of the bug *Nezara viridula* (Master's Thesis). University of Ljubljana, Slovenia.
- Laumann RA, Blassioli Moraes MC, Čokl A, Borges M (2007) Eavesdropping on sexual vibratory signals of stink bugs (Hemiptera: Pentatomidae) by the egg parasitoid *Telenomus podisi*. *Anim Behav* 73: 637-649.
- Laumann RA, Čokl A, Lopes APS, Ferreira JBC, Moraes MCB, Borges M (2011) Silent singers are not safe: selective response of a parasitoid to substrate-borne vibratory signals of stink bugs. *Anim Behav* 82: 1175-1183.
- Laumann RA, Kavčič A, Moraes MCB, Borges M, Čokl A (2013) Reproductive behaviour and vibratory communication of the neotropical predatory stink bug *Podisus nigrispinus*. *Physiol Entomol*, 38: 71-80.
- Murphy ST, Evans HC (2009) Biological control of invasive species. In: Clout M.N., Williams P.A. (Eds.). *Invasive Species Management: A Handbook of Principles and Techniques*, (Techniques in Ecology and Conservation Series). Oxford: Oxford University Press, pp. 77-92.
- Panizzi AR, McPherson JE, James DG, Javahery M, McPherson RM (2000) Stink bugs (Pentatomidae). In: Schaefer C.W., Panizzi A.R. (Eds.). *Heteroptera of Economic Importance*. Boca Raton, Florida: CRC Press, pp. 421-474.
- Polajnar J, Kavčič A, Žunič Kosi A, Čokl A (2013) *Palomena prasina* (Hemiptera: Pentatomidae) vibratory signals and their tuning with plant substrates. *CEJB* 8: 670-680.
- Zorović M, Čokl A (1995) Laser vibrometry as a diagnostic tool for detecting wood-boring beetle larvae. *J Pest Sci* 88: 107-112.
- Žunič A, Čokl A, Virant-Doberlet M, Millar JG (2008) Communication with signals produced by abdominal vibration, tremulation and percussion in *Podisus maculiventris* (Heteroptera: Pentatomidae). *Ann Ent Soc Am* 101: 1169-1178.

## FOREST PHENOLOGY IN CENTRAL-EUROPE BASED ON MODIS DATA

*Anikó Kern<sup>1</sup>, Hrvoje Marjanović<sup>2</sup>, Zoltán Barcza<sup>3</sup>, Laura Dobor<sup>3</sup>*

<sup>1</sup> Department of Geophysics and Space Science, Eötvös Loránd University, 1117 Budapest, Pázmány P. st. 1/A, Hungary

<sup>2</sup> Croatian Forest Research Institute, Cvjetno naselje 41, HR-10450 Jastrebarsko, Croatia

<sup>3</sup> Department of Meteorology, Eötvös Loránd University, 1117 Budapest, Pázmány P. st. 1/A, Hungary

E-mail: hrvojem@sumins.hr

**Keywords:** MODIS, EVI, NDVI, forest activity, forest phenology

Forest phenology and productivity is intimately linked with the actual weather conditions, and in the long term with the local climate. Our current understanding on the environmental control on spring leaf-out and autumn senescence is incomplete. Causes of the interannual variability of tree growth and forest carbon balance are not well understood as well. Satellite remote provides a feasible way to monitor and study the changes of forest activity in general and to understand its relationship with the climate fluctuations.

In this study the Enhanced Vegetation Index (EVI) and the Normalized Difference Vegetation Index (NDVI) calculated from measurements of the MODIS sensor onboard the NASA Terra satellites are used to characterize forest activity in Central Europe (Hungary and Croatia). The applied EVI and NDVI dataset is part of the so-called MOD13 product of NASA and covers the 2000-2014 time period. Based on long term climatologies and data availability we selected a few forest sites in Hungary and Croatia that represent contrasting climatic conditions in the temperature-precipitation space. The selected two Croatian sites are data-rich in terms of information on management, productivity and groundwater level. Using the FORESEE climatological database as archive meteorological data the effects of drought is studied on the NDVI variations. Possible lagged effect of severe drought on NDVI variability during the consecutive years is studied at the selected sites. The effect of water availability in the form of groundwater during dry periods is studied as well as a possible mechanism to mitigate the negative effect of the dry periods. The presented methodology might be used to identify forested regions which are susceptible to climate variations in terms of forest health and production.

### Acknowledgments

The research has been supported in part by the Croatian Science Foundation (HRZZ UIP-11-2013-2492) and Hungarian Scientific Research Fund (OTKA PD-111920 and K-104816).

## THE IMPORTANCE OF SEEDLINGS QUALITY IN TIMBER AND BIO-ENERGY PRODUCTION

*F. Kiourtsis<sup>1</sup>, Ch. Sarvani<sup>1</sup> and P. Bekiaroglou<sup>1</sup>*

<sup>1</sup> Decentralised Administration of Macedonia & Thrace, Prof.Oikonomidi 11str.Thessaloniki, Greece

E-mail: fkiourts@gmail.com

**Keywords:** Seedlings, genetic resources, timber, bio-energy, wood production

One of the main issues that the forest sector is facing is to achieve a balance between the demand for wood production and the need to preserve the sustainability and biodiversity of forest ecosystems. The purposes of the new approaches are to ensure more efficient management of ecosystems and implement intensive forestry that will increase timber yields. To achieve this, we need to determine the macroeconomic potential of the various options available, including the use of biotechnology and genetics. The degradation of the forests natural regeneration capacity may be solved through forest certification, as described below: a) Conservation of biodiversity and stabilization of the forests structure. b) Hierarchy of timber production in the forest's management process. c) Organization and implementation of effective reforestation. d) Maintenance or increase of forest productivity by introducing new items as and when they are required. e) Prevention of the properties that reduce the basic parameters of the trees in seed production stands. It is important to evaluate of the influence of factors such as the quality of reforestation areas, the utilization of the genetic resources and the management of forest operations with the environmental economic criteria such as net present value of benefits (N.P.V) and the corresponding flow annuities (E.A.C.F).The existing evaluations studies showed that the quality of the reforestation areas has the most influence and through validated quality seed production can generate an increase in the NPV up to 73%. The importance of seedlings quality in timber and bio-energy production based on the literature it is estimated according to the heredity of the characteristics of the wood structure (except shrinkage) and it was always higher than that on the characteristics of growth and shape. This clearly indicate that seedlings with the appropriate morphological characteristics can significantly improve the growth performance and help to support the development of reforestation oriented in tailor-made timber and bio-energy production.

Mayfield C, C Smith, B Lattimore (2007) Conserving Biodiversity in Forest Bioenergy Production Systems. Pages 261–266.

Shakti S Chauhan, John CF Walker (2011).Wood quality in artificially inclined 1-year-old trees of *Eucalyptus regnans* — differences in tension wood and opposite wood properties. Canadian Journal of Forest Research, 41(5): 930-937.

Gélinas N,Petrinovic PJF,Beaulieu J (2009) Benefits of using genetically improved white spruce in Quebec - The forest landowner's view point. Forestry Chronicle 85(4): 571-582:

Williams RD and SH Hanks 1976 (slightly revised 1994). Hardwood nursery guide. USDA Forest Service, Agriculture Handbook 473, 78 p.

Douglass F. Jacobs. Nursery Production of Hardwood Seedlings. Hardwood Tree Improvement and Regeneration Center, North Central Research Station, USDA Forest Service ,Department of Forestry and Natural Resources, Purdue University 9/03.

Grigoriou A (2006) Wood based panels products. Production Technology, Properties and Uses. 2<sup>nd</sup> edition. Thessaloniki.

## ROCKFALL MONITORING BY THE COMBINATION OF LIDAR AND UNMANNED AERIAL VEHICLE TECHNOLOGY

*Milan Kobal*<sup>1</sup>

<sup>1</sup> University of Ljubljana, Biotechnical Faculty, Department of Forestry and Renewable Forest Resources, Večna pot 83, 1000 Ljubljana, Slovenia

E-mail: milan.kobal@bf.uni-lj.si

In recent years, the application of unmanned aerial vehicles (UAVs) in disaster monitoring and management operations has become more common due to its advantages, such as low-cost, great mobility and high-resolution acquisitions of real-time aerial photos. Additionally, the areas, relevant for rockfall activity are often difficult to access due to their typical morphology, since the concept of manual controlled quad-rotor systems have been proved to be well suited for terrain changes monitoring in difficult alpine terrain.

In this study, a combination of lidar technology and remote sensing approach, based on UAVs and digital compact cameras, will be presented in the context of rockfall, which occurred in 17. 2. 2014 in the main road Kobarid – Drežnica (Slovenian Alps). In addition to previously existing airborne laser scanning (ALS) data from 2007, significant numbers of airborne photographs of rockfall have been acquired. These images were used for dense point cloud generation, using the structure from motion approach (SfM). By using the original point clouds of both technology, the digital terrain model (DTM) of 0.5 × 0.5 m before and after the rockfall event were derived and different raster maps processing were done to analyses rockfall activation areas.

The results of our analyses provide important perspectives for rockfall activity research: a) detailed geomorphological analyses of rockfall activation areas, b) estimation of the total volume of the rockfall event by a reconstruction of the pre-event surface at the detachment zone and c) for monitoring rockfall activity in time.

***Leotia* cf. *lubrica* FORMS ARBUTOID MYCORRHIZA WITH *Comarostaphylis arbutoides* (ERICACEAE)**

*Katja Kühdorf*<sup>1</sup>, *Babette Münzenberger*<sup>1</sup>, *Dominik Begerow*<sup>2</sup>, *Jorge Gómez-Laurito*<sup>3</sup>, *Reinhard F. Hüttl*<sup>4,5</sup>

<sup>1</sup> Institute of Landscape Biogeochemistry, Leibniz Centre for Agricultural Landscape Research (ZALF), 15374 Müncheberg, Germany

<sup>2</sup> AG Geobotany, Ruhr-University of Bochum, 44780 Bochum, Germany

<sup>3</sup> Escuela de Biología, University of Costa Rica, CP11501-2060 San José, Costa Rica

<sup>4</sup> Chair of Soil Protection and Recultivation, Brandenburg University of Technology Cottbus-Senftenberg, 03013 Cottbus, Germany

<sup>5</sup> German Research Centre of Geosciences Potsdam (GFZ), 14473 Potsdam, Germany

E-mail: Katja.Kuehdorf@zalf.de

**Keywords:** cloud forest, Costa Rica, description of mycorrhiza

Arbutoid mycorrhizal plants are commonly found as understory vegetation in forests worldwide where ectomycorrhiza-forming trees occur. *Comarostaphylis arbutoides* (Ericaceae) is a tropical woody plant and common in tropical Central America. This plant forms arbutoid mycorrhiza, whereas only associations with *Leccinum monticola* (Osmundson et al. 2007) as well as *Sebacina* sp. (Kühdorf et al. 2014) are described so far. We collected arbutoid mycorrhizas of *C. arbutoides* from the Cerro de la Muerte (Cordillera de Talamanca), Costa Rica, where this plant species grows together with *Quercus costaricensis*. We provide here the first evidence of mycorrhizal status for the Ascomycete *Leotia* cf. *lubrica* (Helotiales) that was so far under discussion as saprophyte or mycorrhizal. This fungus formed arbutoid mycorrhiza with *C. arbutoides*. The morphotype was described morphologically and anatomically. *Leotia* cf. *lubrica* was identified using molecular methods, such as sequencing the internal-transcribed spacer (ITS) and the large subunit (LSU) ribosomal DNA regions, as well as phylogenetic analyses. Specific plant primers were used to confirm *C. arbutoides* as the host plant of the leotiid mycorrhiza.

Kühdorf K, Münzenberger B, Begerow D, Karasch-Wittmann C, Gómez-Laurito J, Hüttl RF (2014) *Sebacina* sp. is a mycorrhizal partner of *Comarostaphylis arbutoides* (Ericaceae). *Mycol Prog.* 25: 109–120.

Osmundson TW, Halling RE, den Bakker H (2007) Morphological and evidence supporting an arbutoid mycorrhizal relationship in the Costa Rican páramo. *Mycorrhiza* 17: 217–222.

## EFFECTS OF DIFFERENT FOREST MANAGEMENT INTENSITIES ON BIODIVERSITY IN NATURA 2000 FOREST HABITATS

Lado Kutnar<sup>1</sup>, Klemen Eler<sup>1, 2</sup>, Aleksander Marinšek<sup>1, 3</sup>

<sup>1</sup> Slovenian Forestry Institute, Department of Forest Ecology, Večna pot 2, Ljubljana, Slovenia

<sup>2</sup> University of Ljubljana, Biotechnical Faculty, Jamnikarjeva 101, Ljubljana, Slovenia

<sup>3</sup> Higher Vocational College for Forestry and Hunting, Ljubljanska 3, Postojna, Slovenia

E-mail: lado.kutnar@gozdis.si

**Keywords:** forest management, nature conservation, plant diversity, species turnover, gap colonization, Illyrian *Fagus sylvatica* habitat type

In Slovenia, the Natura 2000 network covers more than 37 % of the country. Forests dominate more than 70 % of this area, and forest management is a significant driver of diversity. Depending on the options applied, forest management may enhance or decrease forest biodiversity. Dinaric fir-beech forests (part of Natura 2000 Illyrian *Fagus sylvatica* habitat type) with significant nature-conservation interest and timber production functions were selected for this study. With the aim of testing the effects of different forest management measures on plant diversity in these forests, and consequently on biodiversity in a broader sense, three sites in the Slovenian part of a Dinaric fir-beech forest range were studied. The plant species diversity was assessed before and after the implementation of forest management measures of three intensities: 1. control plots – no logging; 2. logging of 50 % of the growing stock; and 3. logging of 100 % of the growing stock. Before the implementation of the management measures, the mean number of plant species per 400 m<sup>2</sup> vegetation plots was 48.8, and the mean value of the Shannon diversity index was 2.41. Two years after the measures were implemented, different magnitudes of plant species turnover were observed. There were no significant changes in plant diversity status and vegetation composition in the control plots. Two years after 50 % of the growing stock was logged, the mean number of species was 73.3, and the mean value of the Shannon index was 3.21. In the plots where all the trees were removed, the mean number of species was 87.4, and the mean value of the Shannon index was 3.42. In parallel with the increases in the diversity parameters, the cover of the herbaceous layer increased significantly with an increase in the management intensity, indicating that short-term species turnover can mostly be attributed to herbaceous plant species. As a result of changed stand and ecological conditions, in addition to the increased plant diversity, greater biodiversity in a broader sense and improved habitat suitability for different animal species could be expected.

### Acknowledgements

This study was supported by the European LIFE ManFor C.BD (Managing forests for multiple purposes: carbon, biodiversity and socio-economic wellbeing) Project (LIFE09 ENV/IT/000078). Part of this work was performed within the Research Programmes P4-0107 and P4-0085 (financed by Slovenian Research Agency).

Bouwma IM, van Apeldoorn R, Kamphorst DA (2010) Current practices in solving multiple use issues of Natura 2000 sites: Conflict management strategies and participatory approaches. Alterra, Wageningen, the Netherlands, pp. 77.

Decocq G et al. (2004) Plant diversity in a managed temperate deciduous forest: understory response to two silvicultural systems. *J. Appl. Ecol.* 41: 1065–1079.

Durak T (2012) Changes in diversity of the mountain beech forest herb layer as a function of the forest management method. *For. Ecol. Manage.* 276: 154–164.

Kelemen K et al. (2012) Dynamic response of herbaceous vegetation to gap opening in a Central European beech stand. *Silva Fenn.* 46: 53–65.

## XYLEM SAP FLOW AND RADIAL GROWTH IN *QUERCUS PUBESCENS* WILLD. FROM ABANDONED GRASSLANDS IN SLOVENIAN KARST REGION

*Martina Lavrič<sup>1</sup>, Klemen Eler<sup>2</sup>, Mitja Ferlan<sup>1</sup>, Dominik Vodnik<sup>2</sup>, Jožica Gričar<sup>1</sup>*

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI-1000 Ljubljana, Slovenia

<sup>2</sup> Biotechnical Faculty, Department of Agronomy, University of Ljubljana, Jamnikarjeva 101, SI-1000 Ljubljana, Slovenia

E-mail: martina.lavric@gozdis.si

**Keywords:** pubescent oak, karst, xylem sap flow, radial growth, phenology

Karst region, which is located in SW part of Slovenia, has been significantly changed during the last decades due to the abandonment of marginal agricultural lands, mostly dry calcareous grasslands. In sub-Mediterranean climate, characterized by rather harsh winter and dry and hot summer drought periods frequently occur leading to a great impact on plant and ecosystem functioning.

Pubescent oak (*Quercus pubescens* Willd.), hophornbeam (*Ostrya carpinifolia* Scop.) and manna ash (*Fraxinus ornus*) are the dominant native tree species of the Slovenian Karst region. To survive in such environment, pubescent oak has developed various mechanisms and adaptations, such as conservative water use, tolerance of plant water potential drop, deep and extensive root system, narrow xylem increment etc.

Research was conducted in abandoned karst grassland that is being overgrown with different tree species among which pubescent oak dominates. In the current study we evaluated intra-annual regulation of water of pubescent oak with eco-physiological and wood-anatomical analysis. In the growing season of 2014 (i.e., in the period April–October), we performed the following measurements and analyses: 1) xylem sap flow measurements according to the Heat Ratio Method, 2) micro-core sampling at weekly intervals using Trephor to study the timing of xylem and phloem formation, and the widths and structure of completed xylem and phloem increments, 3) leaf phenological observations, 4) collection of weather data.

Preliminary results show that xylem sap flow is closely related to the climate factors and to internal tree factors. With the development of new shoots and with leaf unfolding at the beginning of May xylem sap flow increased. Cambial cell production and, consequently, xylem and phloem formation started in the third week of March 2014. By the second half of April, the first ring of the earlywood vessels was fully developed and thus ready for water transport. In mid-May the second ring of earlywood vessels was forming in wider xylem growth rings, while latewood formation began in the third week of May. The transition between early and late phloem occurred at the end of May.

In order to get a better insight into tree water relations and radial growth of *Quercus pubescens*, we have to extend measurement period over several years, which will include dry and wet periods.

### Acknowledgements

The study was supported by the Slovenian Research Agency, programs P4-0107, P4-0085 and young researcher program. The authors would like to acknowledge to Boštjan Zupanc from Slovenian Forestry Institute for his help during fieldwork and to Zlatko Rojc for his permission to perform the study on his land.

Burgess S S O, Adams M A, Turner N C, Beverly C R, Ong C K, Khan A A H, Bleby T M (2001) An improved heat pulse method to measure low and reverse rates of sap flow in woody plants. *Tree Physiology* 21: 589-598.

Damesin C, Rambal S (1995) Field study of leaf photosynthetic performance by a Mediterranean deciduous oak tree (*Quercus pubescens*) during a severe summer drought. *New Phytologist* 131: 159-167.

Eilmann B, Zweifel R, Buchmann N, Fonti P, Rigling A (2009) Drought-induced adaptation of the xylem in Scots pine and pubescent oak. *Tree Physiology* 29: 1011-1020.

- Ferlan M, Alberti G, Eler K, Batič F, Peressotti A, Miglietta F, Zaldei A, Simončič P, Vodnik D (2011) Comparing carbon fluxes between different stages of secondary succession of a karts grassland. *Agriculture, Ecosystems and Environment* 140: 199-207.
- Poyatos R, Llorens P, Piñol J, Rubio C (2008) Response of Scots pine (*Pinus sylvestris* L.) and pubescent oak (*Quercus pubescent* Willd.) to soil and atmospheric water deficits under Mediterranean mountain climate. *Annals of Forest Science* 65: 306.
- Sass-Klaassen U, Sabajo C R, den Ouden J (2011) Vessel formation in relation to leaf phenology in pedunculate oak and European ash. *Dendrochronologia* 29: 171-175.
- Steppe K, Sterck F, Deslauriers A (2015) Diel growth dynamics in tree stems: linking anatomy and ecophysiology. *Trends in Plant Science* 20 (6): 335-343.
- Takahashi S, Okada N, Nobuchi T (2013) Relationship between the timing of vessel formation and leaf phenology in ten ring-porous and diffuse-porous deciduous tree species. *Ecological Research* 28: 615-624.
- Tessier L, Nola P, Serre-Bachet F (1994) Deciduous *Quercus* in the Mediterranean region: tree-ring/climate relationships. *New Phytologist* 126: 355-367.
- Tognetti R, Cherubini P, Marchi S, Raschi A (2007) Leaf traits and tree rings suggest different water-use and carbon assimilation strategies by two co-occurring *Quercus* species in a Mediterranean mixed-forest stand in Tuscany, Italy. *Tree Physiology* 27: 1741-1751.
- Wullschlegel S D, Hanson P J, Todd D E (2001) Transpiration from a multi-species deciduous forest as estimated by xylem sap flow techniques. *Forest Ecology and Management* 143: 205-213.

## WILD EDIBLE MUSHROOMS AND MEDICAL HERBS ON MONTENEGRIN MOUNTAINS AND KATUNS

*Jelena Lazarević<sup>1</sup>, Dragica Bojović<sup>2</sup>, Olgica Perić<sup>3</sup>*

<sup>1</sup> University of Montenegro, Biotechnical faculty, Mihaila Lalića 1, 81000 Podgorica, Montenegro

<sup>2</sup> Galenika Crna Gora d.o.o., Ul. 8. mart 55a, Podgorica, Montenegro

<sup>3</sup> *Đulje Jovanova 16*, 81000 Podgorica, Montenegro

E-mail: ena.lazarevic@gmail.com

**Keywords:** mountain regions, inventory, protection, commercial use, rural communities

Mountain regions of Montenegro are rich in forests and biodiversity, including rare and European red listed plant and fungal species. The inventory and protection of them are among the highest priorities. On the other hand, collection of wild edible mushrooms, berries, medical and aromatic herbs could be important sources of income for rural communities. In order to provide a balance between commercial usage and needs for protection, best management practice is necessary to be established.

Ongoing KATUN project is focused on Kuči mountains (SE Montenegro) and Mt. Durmitor (NW Montenegro), where identification of the most valuable medical herbs and wild edible mushrooms are under study. Electronic database with GIS positioning are created for inventory of forests, plants and diversity of fungi.

Raising the public awareness and social responsibility on sustainable use of the resources currently are the highest priority. Hence the workshops and trainings on their sustainable use will be organized for farmers and wider public as well as the manual for collectors will be prepared.

### Acknowledgements

Study was supported by the grant from the Ministry of Science of Montenegro, INVO HERIC No: 01-646.

Lazarević J, Perić O, Perić B (2011) Ectomycorrhizal fungi in Montenegro - diversity and distribution. *Mycologia Montenegrina* XIV (2011): 85-115.

Perić B (2011) Fungi and Flowering plants of Montenegro. Montenegrin Academy of science and Arts, Monographies and Studies, Volume 77, Section of Natural Sciences Volume 34, p. 391.

PERIĆ B, PERIĆ O (2004) Preliminary red list of macromycetes of Montenegro 2°. *Mycol. Monten VII*: 7-33.

Stešević D, Jovović Z. (2008) Plant genetic resources of Montenegro – Medicinal and aromatic plant. University of Montenegro, Biotechnical faculty Podgorica, Podgorica.

Perić O, Perić B (2000) Some most frequent wild edible and medicinal fungi of Montenegro. *Agriculture and Forestry* 46(3-4): 97-119.

## APPLICATION OF AUTOCHTHONOUS FUNGI FOR MYCORRHIZATION OF CONIFERS –TRIALS IN MONTENEGRO

Jelena Lazarević<sup>1</sup>

<sup>1</sup> University of Montenegro, Biotechnical faculty, Mihaila Lalića 1, 81000 Podgorica, Montenegro

E-mail: ena.lazarevic@gmail.com

**Keywords:** *Pisolithus arhizus*, *Suillus granulatus*, forest nurseries, *Pinus nigra*

Mycorrhiza improves water uptake, mineral nutrition, growth and adaptation of forest trees, and provides a protection of seedlings against pathogens. Seedling mycorrhization could be of importance for South-eastern Europe, due to unfavorable environmental conditions: prevailing high summer temperatures of air and soil, low humidity and low precipitation during growing season.

Several thousands of fungi are thought to form ectomycorrhizas (ECM) globally. More than 250 putative ECM species have been recorded in Montenegro until now. Value of fungal species in seedling mycorrhization is different. Compatibility of fungus-host combinations for nursery inoculation and source of inoculums adapted to the environmental conditions of the transplantation site are among the most important criteria for selection.

Possibilities for involvement of different techniques for ECM inoculation of containerized *Pinus nigra* seedlings in nursery production in open field were studied. Ten native ECM fungi were examined as vegetative and spore inoculums. Effect of inoculations on ECM formation and seedling growth characteristics were tested.

The utilization of spore ( $10^6$ ,  $10^7$ ,  $10^8$ ) and vegetative (1:4, 1:8, 1:16) inoculums of *Pisolithus arhizus* and *Suillus granulatus* proved to be an effective method for obtaining containerized ECM *P. nigra* seedlings in open field conditions.

Additional trials also included: use of the soils from four unmanaged forests for coniferous mycorrhization, as well as the use of vegetative inoculums of *Sclerodermatace* in mycorrhization of oaks.

### Acknowledgements

Study was supported by the grant from the Ministry of Science and Ministry of Agriculture and Rural Development of Montenegro, project cont. nmb 01-406/01-558.

Lazarević J, Perić O, Perić B (2011) Ectomycorrhizal fungi in Montenegro- diversity and distribution. *Mycologia Montenegrina* XIV (2011): 85-115.

Lazarević J, Keča N, Martinović A (2012) Mycorrhization of containerized *Pinus nigra* seedlings with *Suillus granulatus* under open field conditions. *Forest Systems* 21(3): 498-507.

Lazarević J, Vilotić D, Keča N (2015) Mycorrhization and use of super absorbent polymers in targeted production of hardwoods planting material. *Agriculture and Forestry* 61/1: 295-307.

## STUDENTS' ATTITUDE TOWARDS COMMERCIALIZATION OF TRANSGENIC FOREST TREES IN SLOVENIA

*Zlata Luthar<sup>1</sup>, Gregor Božič<sup>2</sup>, Robert Brus<sup>1</sup>, Vassiliki Kazana<sup>3</sup>*

<sup>1</sup> University of Ljubljana, Biotechnical Faculty, Jamnikarjeva 101, SI - 1000 Ljubljana, Slovenia

<sup>2</sup> Slovenian Forestry Institute, Večna pot 2, SI - 1000 Ljubljana, Slovenia

<sup>3</sup> Eastern Macedonia and Thrace Institute of Technology, Department of Forestry & Natural Environment Management, GR - 66100 Drama, Greece

E-mail: zlata.luthar@bf.uni-lj.si

**Keywords:** *questionnaire, students' towards, transgenic, forest tree*

Genetically modified organisms (GMOs) are organisms that have been transformed by the insertion of one or more isolated genes. Genetic modification in forestry is much more than a technical issue; sociocultural values and the multiple uses of forests need to be taken into account and public acceptance is necessary if genetically modified forest trees are to be deployed (El-Lakany, 2004). In Slovenia, information on public attitudes towards the use of transgenic forest trees is lacking. To address this issue, a pilot survey was organized using as sampling subjects university students. Specifically students were selected from three different disciplines: forestry, agriculture and biotechnology. The Slovenian survey was conducted in the frame of a cross country pilot survey through the working program of the EU COST Action FP0905. The survey attempted to provide information about the young peoples' attitude regarding: 1) the their knowledge about the use of transgenic forest trees in forest plantations, 2) whether they would agree with the use of transgenic forest trees in forest plantations, and 3) issues related to concerns about using transgenic forest trees in forest plantations. The questionnaires were delivered to selected classes of students. The students completed the questionnaires on site; 180 questionnaires were collected and subject to statistical analysis. Most respondents (86 %) knew what forest transgenic trees are, 70 % did not know whether transgenic forest plantations are grown for commercial purposes in Slovenia and 68 % did not know whether final products of transgenic plantations are sold on the market. About a third of the respondents were highly positive towards the transgenic changes that concern the greater resistance of forest trees and the altered lignin content. The majority of respondents (81 %) would approve funding of transgenic forest plantations for commercial purposes and 86 % appeared willing to purchase products originating from transgenic forest plantations. Most respondents (82 %) agreed that the final products derived from transgenic forest plantations should be labelled and 80 % of respondents agree that labelling such products should be legally mandatory. In terms of the potential benefits of transgenic forest plantations, 66 % of the respondents stated as very important for Slovenia the potential use of fewer pesticides, 63 % the potential better quality wood, 54 % the use of less chemicals and energy for cellulose pulping and 52 % the need for fewer treatments with herbicides of the forest plantations, as well as the increased stress resistance. In relation to the safety of adopting transgenic forest crops: the potential loss of biodiversity due to genetic flows between transgenic and wild trees was the main concern of the great majority of the respondents (72 %), followed by the potential adverse effects of bio-trophic processes of host ecosystems. New genetic traits entering these ecosystems was considered by 46 % of the respondents to be the most potentially serious hazard in Slovenia in relation to adoption of transgenic forest trees.

### Acknowledgments

This study took place within the framework of research tasks of Research Programmes P4-0059, P4-0077, P4-0107 and EU COST Action FP0905.

## FUNGAL AND BACTERIAL COMMUNITIES IN SPRUCE WOOD SAMPLES EXPOSED TO OUTDOORS ENVIRONMENTAL CONDITIONS

*T. Martinović<sup>1,3</sup>, M. Bajc<sup>1</sup>, N. Thaler<sup>2</sup>, M. Humar<sup>2</sup>, H. Kraigher<sup>1</sup> and B. Kraigher<sup>3</sup>*

<sup>1</sup> Department of Forest Physiology and Genetics, Slovenian Forestry Institute, Večna pot 2, 1000 Ljubljana, Slovenia

<sup>2</sup> Department of Wood Science and Technology, Biotechnical Faculty, University of Ljubljana, Jamnikarjeva 101, 1000 Ljubljana, Slovenia

<sup>3</sup> Chair of Microbiology, Department of Food Science and Technology, Biotechnical Faculty, University of Ljubljana, Jamnikarjeva 101, 1000 Ljubljana, Slovenia

E-mail: [tijana.martinovic@gozdis.si](mailto:tijana.martinovic@gozdis.si)

**Keywords:** wood degradation, construction wood, fungi, bacteria, community structure

The problem of industrial wood biodegradation is still an up-to-date topic. Scarcity of more resistant tree species is a limiting factor in most parts of Europe, thus one of the commonly used tree species in building industry is Norway spruce (*Picea abies*). Our aim was to evaluate the differences in fungal and bacterial communities of spruce samples that were treated differently prior to environmental exposure. The first treatment was thermal wood modification and the second one was based on the biocides - alkaline copper quaternary (ACQ, Silvanolin©) for limiting microbial growth. Collected samples were processed mechanically and wood meal was used for DNA extraction. The target regions of the ribosomal operon were amplified in polymerase chain reaction (PCR): ribosomal ITS1 region for fungal communities and 16S rRNA for bacterial communities. The communities were profiled using denaturing gradient gel electrophoresis (DGGE) method and the profiles were analyzed in BioNumerics© program. Representative bands from fungal profiles were processed for sequencing by Sanger method. DGGE profiles showed considerable differences between treated samples and the untreated control for both fungi and bacteria. However, preliminary sequencing results for fungi revealed that most of the sequences do not represent typical biodegrading fungi and the profiles could not be distinguished based on these sequences. Sequences were compared against Unite and INSD databases. Out of 30 sequences in total, roughly 35% corresponded to uncultured fungi from family Herpotrichielaceae (genus *Phaeococcomyces*), 14% family Tremellaceae (genera *Tremella* and *Dioszegia*), 10% family Coniochaetaceae (genus *Coniochaeta*) and 10% family Dothioraceae (genus *Aureobasidium*). There were some representatives of the order Helotiales, as well as genus *Peniophora* and species *Rhodotorulla lamellibrachiae* and *Cryptococcus dimennae*. Most of the identified fungal genera do not belong to the extremely degrading fungi, but predominately to wood-staining fungal species. It should be noted that a number of the sequences recovered in this study exhibited a low level of identity with sequences of known fungi, thus making determination of their identity and biological function within the community impossible. Based on the DGGE profiles of bacterial communities we can speculate about the community shift, especially in case of ACQ treatment, but this needs further validation. In order to improve the results, we plan on prolonging the environmental exposure period and analyzing the samples using next generation sequencing approach (Illumina MiSeq platform).

### Acknowledgements

The contribution was financed by the EUFORINNO project (RegPot No. 315982), L4-5517 project and Research Program P4-0407. This work is also part of master thesis (Martinović T.), mentored by Kraigher B. and Kraigher H.

Lesar B, Humar M, Kamke F A, Kutnar A (2013) Influence of the thermo-hydro-mechanical treatments of wood on the performance against wood-degrading fungi. *Wood Sci. Technol.* **47**, 977–992.

Prewitt L, Kang, Y, Kakumanu M L, Williams M (2014) Fungal and Bacterial Community Succession Differs for Three Wood Types during Decay in a Forest Soil. *Microb. Ecol.* **68**, 212–221.

## COMPARISON OF STAND STRUCTURE IN MANAGED AND VIRGIN EUROPEAN BEECH FORESTS IN SERBIA

*Bratislav Matović<sup>1</sup>, Miloš Koprivica<sup>2</sup>, Bratislav Kisin<sup>3</sup>, Dejan Stojanović<sup>1</sup>, Milan Drekić<sup>1</sup>*

<sup>1</sup> University of Novi Sad, *Institute of Lowland Forestry and Environment, Novi Sad, Serbia*

<sup>2</sup> Institute of Forestry, Belgrade, Serbia

<sup>3</sup> State Enterprise for Forest Management "Srbijašume", Belgrade, Serbia

E-mail: bratislav.matovic@gmail.com

**Keywords:** beech, stand structure, managed forest, virgin forest, Serbia

This study is presenting results of the research of structural and production characteristics of European beech forests in Serbia. Analysed were five managed and three virgin stands. The aim of this study was to perform a comparison of structural and production characteristics of managed and virgin beech forests on the most productive sites in Serbia.

The observed stands are located in the mountainous range of 400 to 1200 meters above sea level. For the gathering and processing of data in the managed stands systematic sample was applied. Sample plots 0.05 ha in size, circled shaped, which are spaced at a distance of 100 x 100 m were set. In all test areas diameter at breast height and height of all trees were measured. In two virgin forests stands total survey of diameters and tree heights were performed, while in the third virgin forest set of seven experimental fields of different sizes approximately square and rectangular shapes where measured. In all stands trees with dbh>10 cm were measured. For comparison of structural characteristics following parameters were used: height curves, diameter distribution, Lorenz index, Gini index, Coefficient of variation, ratio of the height and diameter at breast height and maximum dimensions of trees. Density and production of stands are analysed with taxation elements: number of trees, basal area, volume, biomass, carbon stock, stand quadratic mean diameter and Lorey's mean height.

Height curves of managed and virgin type stands differ to a diameter of 30 cm because the slope of the curve is higher in virgin stands, while the average height of medium thick and thick trees are without significant difference. Diameter distribution of managed type stands can be roughly identified with the selective managed stands. Diameter distribution of virgin type stands Danilova kosa and Vinatovača approximately can be described as the virgin forest stand structure, while diameter distribution of Kukavica stand has a shape that is characteristic of two-story stands. Diameter and height diversity in stands of managed and virgin type in Serbia are similar, which is diametrically different to the most of the managed beech forest in Europe. For all taxation elements there is a statistically significant difference between the average values for managed and virgin forest stands. General conclusion of this research is that the structural characteristics of managed beech forests in Serbia are much closer to their primeval state in comparison to the most of managed beech forests in Europe.

### Acknowledgments

This study was supported by the project "Studying climate change and its influence on the environment: impacts, adaptation and mitigation" (III 43007) financed by the Ministry of Education and Science of the Republic of Serbia within the framework of integrated and interdisciplinary research for the 2011–2014 period.

## ECTOMYCORRHIZA BETWEEN *Scleroderma Areolatum* EHRENB. AND *Fagus sylvatica* L.

*Tanja Mrak*<sup>1</sup>, *Katja Kühdorf*<sup>2</sup>, *Tine Grebenc*<sup>1</sup>, *Ines Štraus*<sup>1</sup>, *Marko Bajc*<sup>1</sup>, *Nada Žnidaršič*<sup>3</sup>, *Babette Münzenberger*<sup>2</sup>, *Hojka Kraigher*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

<sup>2</sup> Institute of Landscape Biogeochemistry, Leibniz Centre for Agricultural Landscape Research (ZALF), Eberswalder Strasse 84, 15374 Müncheberg, Germany

<sup>3</sup> Department of Biology, Biotechnical Faculty, University of Ljubljana, Večna pot 111, 1000 Ljubljana, Slovenia

E-mail: tanja.mrak@gozdis.si

**Keywords:** description of mycorrhiza, morphotyping of mycorrhiza

*Scleroderma* is a genus of widely distributed gasteromyceous species (Sims et al. 1995), many of them are reported to persist in extreme habitats, withstand drought and increased temperatures and being a competitive primary colonizer (Jeffries 1999). They have a potential application as inoculants for forest plantations in areas with poor mycorrhizal status (Chen et al. 2014). *Scleroderma* ectomycorrhizas were obtained on fine roots of beech (*Fagus sylvatica* L.) seedlings from tree nursery and peristed for several years in conditions of increased soil temperatures on seedlings grown in rhizotrons under controlled conditions. The morphotype collected from rhizotrons was described morphologically and anatomically and identified using molecular methods, namely sequencing of the internal-transcribed spacer followed by a genus-based phylogenetic analyses. Based on phylogenetic outcome, the morphotype was assigned to *Scleroderma areolatum* Ehrenb. morphological species which forms a subcluster in SH005470.07FU species hypothesis according to criteria set up by Kõljalg et al. (2013).

Chen YL, Liu RJ, Bi YL, Feng G (2014) Use of Mycorrhizal fungi for Forest Plantations and Minesite Rehabilitation. In: Solaiman Z, Abbott LK, Varma A (Eds.) Mycorrhizal Fungi: Use in Sustainable Agriculture and Land Restoration. Soil Biology, vol 41. Springer-Verlag Berlin Heidelberg, pp. 325-355.

Jeffries P (1999) *Scleroderma*. In: Ectomycorrhizal Fungi Key Genera in Profile. Springer Berlin Heidelberg, pp. 187-200.

Kõljalg U et al. (2013) Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology* 22: 5271-5277.

Sims KP, Watling R, Jeffries P (1995) A revised key to the genus *Scleroderma*. *Mycotaxon* 56: 403-420.

## TOTAL AMOUNT OF CARBON HOLD UNDER DIFFERENT AGES EUCALYPTUS TREES, PLANTS WITH DIFFERENT SOIL AND CROP MANAGEMENT SYSTEMS IN COST OF THE MEDITERRANEAN CONDITIONS

*Ibrahim Ortas*<sup>1</sup>, *Cengiz Darici*<sup>3</sup>, *Sedat Tufekci*<sup>2</sup>, *Sahin Cenkseven*<sup>3</sup>, *Ahu Kutlay*<sup>3</sup>, *Cagdas Akpinar*<sup>1</sup>, *Nacide Kizildag*<sup>3</sup>, *Murat Simsek*<sup>1</sup>, *Ahmet Demirbas*<sup>1</sup>

<sup>1</sup> University of Cukurova, Department of Soil Science, Adana, Turkey

<sup>2</sup> Eastern Mediterranean Forestry Research Institute, Tarsus, Turkey

<sup>3</sup> University of Cukurova, Department of Biology, Adana, Turkey

E-mail: iortas@cu.edu.tr

The biggest soil carbon (C) sink and C sequestration sources in terrestrial ecosystems is mainly coming from forestry area. Forest ecosystems store C as lignin and other relatively resistant C compounds. The forest C is sequestered not only in the harvestable timber and other branched. It is important to keep the atmospheric CO<sub>2</sub> in the plant tissue for long term sequestration. Recently fiber forest farming and agro forestry also getting more used. One of the fiber forestry is eucalyptus. Eucalyptus is a common plantation in Mediterranean cost conditions. Since eucalyptus timbers are used for ecological and economical it is sound to calculate effect of several years old tree on total biomass and total C sequestration.

The experiment was carried out on three age's groups eucalyptus plantation in the Regional Forest and Forestry Management area which is located in Mersin-Tarsus, Southern of Turkey.

The eucalyptus (*E. camaldulensis*) tree in the experimental plan for each age group was harvested from the soil surface with motorized saws. There were three different treatments such as (SI) shaved the three trees from soil surface and ploughed, (SII) Shaved trees without plowing (SIII) Control without shaved the trees.

Research areas that are connected to three different age groups 5, 7 and 10 years old which have different diameter at breast height of trees high and mass. Branch and leaf, fresh and dry weight and total moisture content were also determined by calculating the total biomass. For each age group all the tree diameters and height taken into consideration with the *E. camaldulensis* of double-entry volume tables by utilizing a single tree volume calculating on hectare.

According to the research findings in five years old eucalyptus total biomass weighing for SI, shaved was 82 tons ha<sup>-1</sup> and shaved and plowed was 105 ton ha<sup>-1</sup> respectively. In the 7 age group SI was 107 ton ha<sup>-1</sup>, and the SII was 125 ton ha<sup>-1</sup> biomass produced. In the 10 age group in SI was 151 ton ha<sup>-1</sup> and SII was 150 ton ha<sup>-1</sup> bio-mass production was obtained.

Also the soil organic carbon (SOC) stock was calculated for each ages and management systems. Apart from SOC, total soil nitrogen was calculated as well. Soils have collected at 5 years old eucalyptus plantation total 40.46 ton ha<sup>-1</sup> SOC, and respectively to 7 and 10 years old plantation they have accumulated 56.04 and 82.66 ton ha<sup>-1</sup> SOC. The SOC accumulation is strongly depends on plantation ages and their duration on the soil.

## INFLUENCE OF HEAVY FOREST MACHINERY ON FINE ROOTS: OBSERVATIONS ON MINIRHIZOTRON DATA

*Yasmine Pinuela Samaniego*<sup>1</sup>, *Peter Železnik*<sup>2</sup>, *Hojka Kraigher*<sup>2</sup>

<sup>1,2</sup> Technical University of Madrid (UPM).

<sup>2</sup> Slovenian Forestry Institute

yasmin.pinuela@hotmail.com

**Keywords:** Fine root turnover, *Picea abies*, heavy forest machinery

Between 10% and 20% of the total roots in forest are fine roots (Jackson et al. 1997), therefore the amount of carbon (C) stored in fine roots is significant and relevant in forest C cycle. Fine roots play important role in long-term changes in soil organic matter and ecosystem carbon balance.

In a managed forest, the influence of soil compaction by heavy forest machinery on fine roots dynamics is not well measured, but still observed in some research (Železnik et al. 2015). We observed impact of logging activities on fine root dynamics with minirhizotrons. Fine root turnover and longevity of Norway spruce (*Picea abies* (L.) H.Karst), situated on mountain region Pohorje (NE Slovenia), was analyzed.

Minirhizotron tubes were installed on two different locations. On each site, tubes were placed in and beside wheel rut and in undisturbed soil profiles. Under the most adverse conditions (in the wheel ruts made by harvester and forwarder) fine roots were almost absent. In average, the highest fine root longevity was observed by wheel rut and the lowest in undisturbed soils.

### Acknowledgments

The research was funded by the Young Researcher funding scheme of the Slovenian Research Agency (ARRS), the Research Programme P4-0107, and projects L4-2265 and L4-4318, co-funded by the Ministry responsible for forestry. The final analyses were done within the 7FP EU project EUFORINNO (Regpot No. 315982).

Železnik P, Vilhar U, Starr M, de Groot M, Kraigher H (2015) Fine root dynamics in Slovenian beech forest in relation to soil temperature and water availability. *Trees-Struct Funct* (doi:10.1007/s00468-015-1218-z)

Jackson RB, Mooney HA, Schulze ED (1997). A global budget for fine root biomass, surface area, and nutrient contents. *Ecology* 94: 7362 – 7366.

## VISIBILITY AND DISSEMINATION IN FOREST GENETIC MONITORING

*Boris Rantaša*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: boris.rantasa@gozdis.si

**Keywords:** Communications, dissemination, forest, genetic, monitoring

Communication skills are very important for scientists and practitioners in the field of forest genetic monitoring (FGM). In the past, communication and dissemination activities were often avoided by scientists, experts and professionals because they were perceived as difficult, time consuming, stressful and tiresome. Today, the best path to achieve the implementation and ensure long-lasting results of FGM is through good communication and appropriate dissemination. It is the duty of any FGM scientist, expert or professional to disseminate their knowledge in an ethical way, because: (i) FGM research is mostly publicly funded; (ii) the public deserves an understandable presentation of FGM; (iii) feedback from different stakeholder groups allows FGM knowledge and future research to be suited to its practical users and therefore more relevant; (iv) new opportunities and synergies can be fostered through communication activities and channels. The willingness to communicate and personal communication skills of FGM scientists, experts and professionals will be a crucial factor in the evolution and implementation of FGM systems in the future.

### **Acknowledgements**

Research Program P4-0107, Public Forest Service, CRP V4-1438, EUFORGEN WG on Genetic monitoring, FP7 EUFORINNO and LIFE GENMON project, financed by LIFE+, MOP, MKGP and SFI.

## OZONE IN URBAN FORESTS

*Matej Rupel<sup>1</sup>, Daniel Žlindra<sup>1</sup>*

<sup>1</sup> Department of Forest Ecology, Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

E-mail: matej.rupel@gozdis.si

Ozone is still the important air pollutant, dangerous for plants, animals and humans.

The Slovenian Forestry Institute (SFI) have more than ten years experiences (2003 – 2015), assessing ozone visible injury on forest vegetation in Slovenia, observed as foliage damage on the forest trees and shrubs species. Our first observations of negative impacts of ozone on forest trees were in year 2003. During these years the visible ozone damages were observing at 5 to 11 intensive monitoring plots (ICP Forests Level II) and in the years 2013 and 2014 also in Ljubljana urban and peri-urban forests.

In the same time span the SFI has been monitoring ozone concentrations in the air with passive samplers. In 2013, we expanded air monitoring to three other pollutant gases: sulphur dioxide, nitrogen dioxide and ammonia. We monitor air pollution with passive samplers also in the capital of Slovenia – Ljubljana. We perform measurements in urban areas and urban and peri-urban forests (city parks, nature park Tivoli, Rožnik, Šišenski hrib, forest covered hill Golovec, riparian forests).

Identification of visible ozone or visible ozone like symptoms in small trees and shrubs in Ljubljana urban forests was performed. Visible foliar injury by ozone we surveyed on 3 *off-plots*, on light-exposed sampling sites at urban forest edge according to the ICP Forests manual. We looked at 34 different species.

Different species show different sensibility to the same ozone dose.

Due to the environment pollution and impacts on the forest, urban and peri-urban forests and vegetation, we present the acquired conclusions to the wider public.

## SOIL RESPIRATION VARIABILITY IN BOREAL PINE FOREST IN VÄRRIÖ, FINLAND

*Iztok Sinjur*<sup>1</sup>, *Liisa Kulmala*<sup>2</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana

<sup>2</sup> University of Helsinki, P.O. Box 33 (Yliopistonkatu 4), FI – 00014 University of Helsinki, Finland

E-mail: iztok.sinjur@gozdis.si

**Keywords:** soil respiration, carbon dioxide, Boreal forest, weather

Soil respiration stands for carbon dioxide (CO<sub>2</sub>) emissions from soil to atmosphere. It is a sum of CO<sub>2</sub> that is emitted by soil microbes, soil animals and plant roots. During summer days, forest floor vegetation photosynthesizes and takes up CO<sub>2</sub> from the atmosphere. Then the observed net exchange at the forest floor can be much less than the soil respiration only since the canopy especially in northern forest is open providing considerably light also on the forest floor. The influx and efflux ratio is driven by various factors i.e. atmospheric and soil conditions, daylight as well as flora, fauna and disturbances caused by human or animals. In high latitude Boreal forests, where ecosystem productivity, decomposition and accumulation of organic matters are low, high spatial diversity in soil conditions and weather variability significantly affect the amount of released carbon dioxide into the atmosphere. During the summer and autumn 2014, we conducted a periodical chamber measurements of carbon dioxide emissions on twelve collars, distributed in two transects inside the fence of SMEAR I research site using close-path system with portable chambers. The measurements are repeated in ambient light conditions and in darkness with transparent and opaque chamber, respectively. The site is located 220 km northeast from Rovaniemi, Finland, at the top of hill Kotovaara (390 m a.s.l., 67.75516° N, 29.60997° E), inside of uneven-aged Scots pine (*Pinus sylvestris* L.) stand and with some mountain birch (*Betula pubescens* subsp. *Czerepanovii*). The measurement site is fenced in order to protect from reindeer grazing. We analyzed readings of eight measuring campaigns from June to October 2014 with sample rate of fifteen days and dataset of continuous close-path measuring system using two chambers. The analyses showed significant differences between collars and pointing out effects of (under) ground variability as well as weather conditions.

### Acknowledgement

We would like to acknowledge a staff members of the Research station Värriö for their help during field campaigns.

## **RADIAL GROWTH RESPONSES OF PINUS HALEPENSIS MILL. AND *Pinus Pinea* L. FORESTS TO CLIMATE VARIABILITY IN WESTERN ALBANIA**

*Elvin Toromani*<sup>1</sup>, *Edmond Pasho*<sup>1</sup>, *Arben Alla*<sup>1</sup>, *Vasillaq Mine*<sup>1</sup>, *Nehat Çollaku*<sup>1</sup>

<sup>1</sup> Faculty of Forestry Sciences, Agricultural University of Tirana, 1029 Kodër-Kamëz, Tirana, Albania

E-mail: etoromani@ubt.edu.al

In this study are presented chronologies of earlywood (EW), latewood (LW) and tree-ring widths (RW) of a *Pinus halepensis* (*P. halepensis*) and *Pinus pinea* (*P. pinea*) natural forest stand growing in western Albania. Bootstrapped correlations and pointer year analysis were combined in a dendroclimatological study to evaluate climate-growth relationships in both pine species as well as to assess the spatial outreach of our chronologies evaluating them with those of the same species from other Mediterranean countries. We found that both species responded positively to precipitation and Indexed Percentage Average Precipitation (% AvP) in late summer-early autumn, particularly the LW, whereas summer temperatures constrained the growth of *P. halepensis* tree-ring features. Current January temperature and Potential Evapotranspiration (PET) showed positive relationship with *P. pinea* LW and RW. The same association was observed when considering PET in spring and *P. halepensis* LW and RW. Pointer year analysis showed that inhibitory climatic drivers of radial growth for both species were low precipitation from previous winter and current summer, associated with low temperatures during autumn. Our *P. halepensis* chronology showed a wider spatial outreach than that of *P. pinea* when compared to those from other Mediterranean countries. We conclude that current January temperatures and September precipitation are very important for *P. pinea* growth influencing both EW and LW growth whereas *P. halepensis* is mostly affected by the summer-early autumn climate conditions.

## LINKAGES OF PLANT MORPHOLOGICAL-FUNCTIONAL TRAITS TO SOIL PROPERTIES ON EXTENSIVELY USED GRASSLANDS OF POHORJE MOUNTAINS

*Tina Unuk*<sup>1</sup>, *Nataša Pipenbaher*<sup>2</sup>, *Sonja Škornik*<sup>2</sup>

<sup>1</sup> Cirkovce 1k, 2326 Cirkovce, Slovenia

<sup>2</sup> University of Maribor, Faculty of Natural science and Mathematics; Biology dept., Koroška 160, SI-2000 Maribor, Slovenia

E-mail: tina.unuk@gmail.com

**Keywords:** plant functional types, soil analysis, matgrass grasslands, species richness, multivariate analysis

In our research we used functional approach to determine relation between soil properties (chemical reaction – pH, phosphorus, potassium, nitrogen, calcium carbonate, C : N proportion, organic carbon and humus), species richness and *morphological-functional plant traits* (MFT) on extensively used oligotrophic grasslands of Rogla (Pohorje, NE Slovenia).

Our research was based on 60 vegetation relevés (plots) of extensively used grasslands. In this vegetation plots we recorded 60 plants species, for which we collected MFT. The research was divided into two parts; in the first part we used all 60 vegetation plots - 29 vegetation plots were from typical matgrass (*Nardus stricta*) grasslands and 31 vegetation plots were from ski trail Mašinžaga. Those areas/grasslands were compared, based on floristical composition, species richness and MFT. In the second part of the research we included only those 23 vegetation samples/plots, on which soil analyses were made.

We found that typical matgrass (*Nardus stricta*) grasslands and grassland Mašinžaga vary floristically as well as in species richness and occurrence of MFT. Those findings can be result of the variation of environmental conditions on those grasslands. Due to those results we decided to do further analyses of MFT and environmental variables. We analysed our data with two separate RDA analyses. Results of the first RDA analysis divide plant species into the three groups.

The first group is linked to the higher soil pH values and lower values for phosphorus, potassium and nitrogen. Those areas were found on deeper soil on ski trail Mašinžaga, where plant communities from association *Arrhenatherion* were developed. The second group of plant species can be found on areas with higher P, K and N, although they can not be absorbed because of low soil pH. Species are typical acidophilic plants (such as *Arnica montana*). The third group of species can be found on extreme areas – soils are acidic, very shallow and at the same time poor with nutrients. Here we can found pioneer species from genus of moss *Polytrichum*. MFT was also divided due to environmental data on those grasslands. On areas of ski trail Mašinžaga where are soils with higher pH and lower values of P, K and N, we found higher plants height, higher values of SLA, more tussock plants and plants with rosettes and life strategies CSR\_R and CSR\_C. On typical matgrass grasslands, where soil is acidic with higher values of P, K and N, we found more phanerophytes, geophytes and life strategies CSR\_S with higher values of LDMC.

Occurrence of small plant height on nutrient-rich soil can be explained with lower values of soil pH, because plants on such soil are not so efficient in absorbing nutrients. Similar explanation was offered for findings that in conditions where pH values are higher and at the same time values of P, K and N are lower, plants have higher values of SLA.

**Acknowledgements** I would like to thank my mentor doc. dr. Škornik and comentor doc. dr. Pipenbaher for all the support and help during the research.

Cornelissen JHC, Lavorel S, Garnier E, Diaz S, Buchmann N, Gurvich DE, Reich PB, ter Steege H, Morgan HD, van der Heijden MGA, Pausas JG,

- Poorter H (2003) A handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Australian Journal of Botany* 51: 335-380.
- García – Palacios P, Maestre FT, Milla R (2013) Community-aggregated plant traits interact with soil nutrient heterogeneity to determine ecosystem functioning. *Plant Soil* 364: 119-129.
- Kaligarič M, Škornik S (2009) Analiza travniške vegetacije in smernice za sanacijo tekaške steze na Rogli. Končno poročilo, Fakulteta za naravoslovje in matematiko.
- Laliberte E, Shipley B, Norton DA, Scott D (2012) Which plant traits determine abundance under long-term shifts in soil resource availability and grazing intensity? *Journal of Ecology* 100: 662-677.
- Roscher C, Schmid B, Buchmann N, Weigelt A, Schulze ED (2011) Legume species differ in the responses of their functional traits to plant diversity. *Oecologia* 165: 437-452.
- Schellberg J, Pontes LdaS (2012) Plant functional traits and nutrient gradients on grassland. *Grass and Forage Science* 67: 305–319.

## FINE ROOT DYNAMICS IN SLOVENIAN BEECH FORESTS IN RELATION TO SOIL TEMPERATURE AND WATER AVAILABILITY

*Peter Železnik<sup>1</sup>, Urša Vilhar<sup>1</sup>, Mike Starr<sup>2</sup>, Maarten de Groot<sup>1</sup>, Hojka Kraigher<sup>1</sup>*

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI – 1000 Ljubljana, Slovenia

<sup>2</sup> Department of Forest Sciences, University of Helsinki, P.O. Box 27, 00014 Helsinki, Finland

E-mail: [ursa.vilhar@gozdis.si](mailto:ursa.vilhar@gozdis.si)

**Keywords:** fine root ingrowth, fine root mortality, environmental factors, forest floor precipitation, evapotranspiration

### **Acknowledgements**

The study was part of a Ph.D. study, the 5th EU FP project Nature-Based Management of Beech in Europe (NAT-MAN 648 QLK-CT99-1349), ManFor C. BD project, titled “Managing forests for 649 multiple purposes: carbon, biodiversity and socio-economic wellbeing” (LIFE09ENV/IT/000078), EUFORINNO — “European Forest Research and Innovation” (Reg. Pot No. 315982), several projects within the Programme group “Forest biology, ecology and technology”, and finalized within the project “Carbon dynamics in forest soils and the rhizosphere” financed by the Ministry of Education, Science and Sport of the Republic of Slovenia.

Campbell JJ, Fine'r L, Messier C (1998) Fine-root production in small experimental gaps in successional mixed boreal forests. *J Veg Sci* 9: 537–542.

Fine'r L, Messier C, De Grandpre' L (1997) Fine-root dynamics in mixed boreal conifer—broad-leafed forest stands at different successional stages after fire. *Can J For Res* 27: 304–314. doi:10.1139/x96-170.

Fine'r L et al (2007) Variation in fine root biomass of three European tree species: Beech (*Fagus sylvatica* L.), Norway spruce (*Picea abies* L. Karst.), and Scots pine (*Pinus sylvestris* L.). *Plant Biosyst Int J Dealing Asp Plant Biol* 141(3): 394–405.

Fine'r L, Ohashi M, Noguchi K, Hirano Y (2011a) Factors causing variation in fine root biomass in forest ecosystems. *For Ecol Manage* 261: 265–277. doi:10.1016/j.foreco.2010.10.016.

Fine'r L, Ohashi M, Noguchi K, Hirano Y (2011b) Fine root production and turnover in forest ecosystems in relation to stand and environmental characteristics. *For Ecol Manage* 262: 2008–2023. doi:10.1016/j.foreco.2011.08.042.

Grebenc T, Christensen M, Vilhar U, Čater M, Mart'ín MP, Simončič P, Kraigher H (2009) Response of ectomycorrhizal community structure to gap opening in natural and managed temperate beech-dominated forests. *Can J For Res* 39: 1375–1386.

Kaspar TC, Bland WL (1992) Soil temperature and root growth. *Soil Sci* 154: 290–299.

## ROOT GROWTH DYNAMICS OF THREE BEECH (*Fagus sylvatica* L.) PROVENANCES

*Peter Železnik*<sup>1</sup>, *Marjana Westergren*<sup>1</sup>, *Gregor Božič*<sup>2</sup>, *Klemen Eler*<sup>2</sup>, *Marko Bajc*<sup>1</sup>, *Heljä-Sisko Helmisaari*<sup>3</sup>, *Aniko Horvath*<sup>4</sup>, *Hojka Kraigher*<sup>1</sup>

<sup>1</sup> Slovenian Forestry Institute, Večna pot 2, SI-1000 Ljubljana, Slovenia

<sup>2</sup> Biotechnical Faculty, University of Ljubljana, Jamnikarjeva 101, SI-1000 Ljubljana, Slovenia

<sup>3</sup> Department of Forest Sciences, University of Helsinki, P.O. Box 27, FI-00014 University of Helsinki, Finland

E-mail: peter.zeleznik@gozdis.si

**Keywords:** fine root turnover, longevity, root biomass, provenance adaptation

European beech (*Fagus sylvatica* L.) is commercially and ecologically important tree species in Central European forests but its intra-specific variability in drought and temperature tolerance might endanger its future distribution in Europe. Beech phenological and growth traits have been studied in large-scale international beech provenance trials, yet the growth and turnover of its fine roots (FR) has not been included among the observations. FR growth dynamics of three beech provenances in the international beech provenance trial Straza / Kamenski hrib, established in Slovenia in 1998, and from a natural beech regeneration site growing at its border, were studied from 2007 to 2010. We studied FR biomass using soil cores (SC), root production using ingrowth soil cores (IC), and root longevity using minirhizotrons (MR). There were no significant differences in total FR biomass (live and dead) among the tested provenances but live FR biomass differed between two of three tested provenances in the trial from seedlings in the natural regeneration site. Values of specific root length (SRL) in IC varied significantly among sampling periods, except for the natural regeneration. The turnover rates in IC were at the end of the experiment close to MR results. Median MR-based longevity of fine roots varied between 625 and 934 days. Survival curve of the slowest growing provenance (considering its aboveground characteristics) was significantly different from the other two, median longevity of later being higher. Death of FR, older than two years, occurred most likely in the winter. Our results suggest that there are significant differences in FR longevity among provenances, which might contribute to their adaptation to future environmental conditions. Furthermore, the calculated annual C investment into FR growth per ha differs up to twofold between provenances, contributing to different C dynamics of their future stands.

Alia R, Božič G, Gömöry D, Huber G, Rasztoivts E, von Wühlisch G (2010) The survival and performance of beech provenances over a Europe-wide gradient of climate. Paper presented at the COST E52 Final Meeting, Burgos, Spain, 4-6th May 2010.

Bardgett RD, van der Putten WH (2014) Belowground biodiversity and ecosystem functioning. *Nature* 515: 505-511 doi:10.1038/nature13855.

Mátyás C, Božič G, Gömöry D, Ivanković M, Rasztoivts E (2009a) Juvenile growth response of European beech (*Fagus sylvatica* L.) to sudden change of climatic environment in SE European trials. *iForest - Biogeosciences and Forestry* 2: 213-220 doi:10.3832/for0519-002.

Mátyás C, Božič G, Gömöry D, Ivanković M, Rasztoivts E (2009b) Transfer Analysis of Provenance Trials Reveals Macroclimatic Adaptedness of European Beech (*Fagus sylvatica* L.). *Acta Silvatica and Lignaria Hungarica* 5: 47-62.

McCormack ML et al. (2015) Redefining fine roots improves understanding of below-ground contributions to terrestrial biosphere processes. *New Phytologist*: n/a-n/a doi:10.1111/nph.13363.

Prislan P, Gričar J, de Luis M, Smith KT, Čufar K (2013) Phenological variation in xylem and phloem formation in *Fagus sylvatica* from two contrasting sites. *Agricultural and Forest Meteorology* 180:142-151 doi:http://dx.doi.org/10.1016/j.agrformet.2013.06.001.

Robson MT et al. (2010) The timing of leaf flush in European beech (*Fagus sylvatica* L.) saplings. Paper presented at the COST E52 Final Meeting, Burgos, Spain, 4-6th May 2010.

Štraus I, Mrak T, Ferlan M, Železnik P, Kraigher H (2014) Influence of soil temperature on growth traits of European beech seedlings. *Canadian Journal of Forest Research*: 245-250 doi:10.1139/cjfr-2014-0332.

Železnik P, Hrenko M, Then C, Koch N, Grebenc T, Levanič T, Kraigher H (2007) CASIROZ: Root parameters and types of ectomycorrhiza of young beech plants exposed to different ozone and light regimes. *Plant Biology* 9: 298-308.

Železnik P, Vilhar U, Starr M, de Groot M, Kraigher H (2015) Fine root dynamics in Slovenian beech forests in relation to soil temperature and water availability. *Trees - Structure and Function* 2015.

## INDEX OF AUTHORS

Name	Page	Name	Page	Name	Page
Ahmed I.	22, 33	Ceulemans R.	9, 31	Grebenc T.	37, 44, 55, 56, 58, 73
Akpinar C.	74	Çollaku N.	79	Gričar J.	14, 16, 66
Alberti G.	11	Constantin N.	15	Gutiérrez E.	12
Alexandrov A. H.	3	Cristian S.	15	Hafner P.	16
Alla A.	79	Cruz C.	27	Hansson K.	24
Andonovski V.	5	Čortan D.	6	Helmisaari H-S.	23, 24, 25, 83
Andrić I.	47	Čustović H.	57	Horemans J. A.	31
Anić M.	34	Darici C.	74	Horvath A.	83
Aravanopoulos F.	2, 5	De Boeck H.	7	Hrenko M.	52, 56
Aro L.	24	De Cinti B.	20	Hukić E.	57
Arriga N.	9	de Groot M.	82	Humar M.	71
Avramidou E. V.	5	De Groote T.	9	Hüttl R. F.	64
Baier R.	5	Deckmyn G.	31	Ionel P.	15
Bajc M.	38, 39, 54, 58, 71, 73, 83	Demirbas A.	74	Jentsch A.	7
Baldrian P.	21	Derome K.	30	Jukić B.	48
Ballian D.	5, 17, 48, 55	Dias T.	27	Kajba D.	5, 47, 48, 51
Baloh T.	37	Dighton J.	26	Kandemir G.	4
Barcza Z.	61	Dobor L.	31, 61	Karadelev M.	55
Barna M.	31	Domec J-C.	10	Karaman M.	54
Bashkim T.	49	Drekić M.	72	Katanić M.	58
Begerow D.	64	Drolc T.	43, 44, 45	Kavaliauskas D.	5
Bekiaroglou P.	62	El Kasmiooui O.	9	Kavčič A.	59
Benucci G. M. N.	55	Eler K.	11, 65, 66, 83	Kaya Z.	4
Berggren Kleja D.	24	Ernst D.	8	Kayler Z.	19
Berhongeray G.	9	Faruk B.	49	Kazana V.	70
Berki I.	50	Ferlan M.	11, 42, 66	Kern A.	34, 61
Bloor J.	7	Ferreira A.	51	King J. S.	10
Bojović D.	68	Finžgar D.	52	Kiourtsis F.	5, 62
Bosela M.	31	Fischer M.	10	Kisin B.	72
Božič G.	5, 51, 52, 56, 70, 83	Frank U.	8	Kizildag N.	74
Broeck A. V.	51	Freitas H.	32	Kobal M.	63
Broeckx L. S.	9	Fussi B.	2, 5	Kolozs L.	51
Brus R.	70	Galović V.	53, 54	Konnert M.	2, 5
Buters J. T. M.	8	Ganopoulos I. V.	5	Konrad H.	5
Campelo F.	32	Gessler A.	18, 19	Koprivica M.	72
Carvalho A.	32	Godbold D.	22, 33	Kovačević B.	58
Çengel B.	4	Gómez-Laurito J.	64	Kovič Dine M.	29
Cenkseven S.	74	Göransson H.	22, 33		
		Görres C.	9		
		Gray D. M.	26		

## INDEX OF AUTHORS

Name	Page	Name	Page	Name	Page
Kraigher B.	71	Münzenberger B.	64, 73	Stojanović D.	72
Kraigher H.	2, 5, 37, 43, 44, 45, 51, 52, 54, 55, 56, 58, 71, 73, 75, 82, 83	Nabais C.	32	Sucre E.	10
Kreft I.	28	Nieminen T. M.	30	Sulzbacher M.	55
Kremer A.	1	Nijs I.	7	Szabados L.	53
Kreyling J.	7	Njakou Djomo S.	9	Šeguljev B.	54
Kühndorf K.	64, 73	Nöjd P.	30	Šijačić-Nikolić M.	6
Kulmala L.	78	Nousiainen V.	25	Škornik S.	80
Kutlay A.	74	Öder S.	8	Štraus I.	73
Kutnar L.	65	Orlović S.	5, 53, 54, 58	Tayanç Y.	4
Lavrič M.	66	Ortas I.	74	Thaler N.	72
Lazarević J.	55, 68, 69	Ostrogović Sever M. Z.	34	Toromani E.	79
Leppälammii- Kujansuu J.	23, 24, 25	Pasho E.	79	Traidl-Hoffmann C.	8
Levanič T.	16, 17, 37, 41, 43, 44, 45	Peresotti A.	11	Tufekci S.	74
Linder S.	24	Perez I.	53	Ukonmaanaho L.	30
Lindroos A-J.	30	Perić O.	68	Ulrich A.	19
Lipar Ž.	45	Peteh M.	43, 44, 45	Unuk T.	80
Lukić T.	55	Pinuela Samaniego Y.	83	Vanbeveren S.	9
Luthar Z.	70	Pipenbaher N.	80	Velioğlu E.	4
Makita N.	25	Poljanšek S.	17	Veperdi G.	50
Malliarou E. S.	5	Premke K.	19	Verlinden M. S.	9
Marinšek A.	65	Ransijn J.	7	Veselič Ž.	5
Marjanović H.	34, 61	Rantaša B.	76	Vieira J.	32
Martin M. P.	55	Rasztovits E.	50	Vilhar U.	82
Martinović T.	71	Rebov I.	45	Vilotić D.	6
Martins-Loução M. A.	27	Robek R.	37, 38, 39	Vodnik D.	11, 66
Mašek A.	56	Rossi S.	32	von Rein I.	19
Matavuly M.	58	Rupel M.	77	von Thörne C.	8
Matović B.	72	Salemaa M.	24	von Wüehlich G.	48
Matteucci G.	20	Sarvani Ch.	62	Westergren M.	2, 5, 51, 83
Matyssek R.	36	Segers J.	9	Williamson J.	10
McCarroll D.	13	Sheppard L.	27	Zavadlav S.	41
McDowell N.	35	Simončić P.	11, 37, 57	Zeiter M.	7
Merilä P.	30	Simsek M.	74	Zenone T.	9
Mine V.	79	Sinjur I.	78	Železnik P.	43, 75, 82, 83
Móricz N.	50	Skudnik M.	16	Žlindra D.	39, 77
Mrak T.	40, 73	Smith A.	22	Žnidaršič N.	73
		Smith A. R.	33		
		Starr M.	30, 82		
		Stephens B.	30		

## List of Participants

Surname	First name	E-mail	Institute
Ahmed	Iftekhar Uddin	iftekhar.ahmed@boku.ac.at	Universität für Bodenkultur (BOKU), Austria
Alexandrov	Alexander	alexandrov_38@abv.bg	Forest Research Institute, Bulgaria
Andonovski	Vlatko	makmontana1@t-home.mk	Univ. Cyril & Methodius, Fac. of Forestry, FYRO Macedonia
Atanasova	Ralitsa	ralitsa.atanasova@ec.europa.eu	European Commission, EUFORINNO project officer
Bahovec	Marjana	marjana.westergren@gmail.com	Retired person, Slovenia
Bajc	Marko	marko.bajc@gozdis.si	Slovenian Forestry Institute, Slovenia
Balanč	Tadeja	tadeja.balanc@gmail.com	University of Ljubljana, Slovenia
Baldrian	Petr	baldrian@biomed.cas.cz	Institute of Microbiology of the ASCR, Czech Republic
Ballian	Dalibor	balliandalibor9@gmail.com	Univ. of Sarajevo, Faculty of Forestry, Bosnia and Herzegovina
Baloh	Tjaša	tjasa.baloh@gozdis.si	Slovenian Forestry Institute, Slovenia
Batič	Franc	franc.batic@bf.uni-lj.si	University of Ljubljana, Biotechnical Faculty, Slovenia
Bavcon	Jože	joze.bavcon@guest.arnes.si	University Botanic Gardens Ljubljana, Slovenia
Boeck	Hans	hans.deboeck@uantwerp.be	University of Antwerp, Belgium
Božič	Gregor	gregor.bozic@gozdis.si	Slovenian Forestry Institute, Slovenia
Ceulemans	Reinhart	reinhart.ceulemans@uantwerp.be	University of Antwerp, Centre of Excellence PLECO, Belgium
Čopič	Andreja	andreja.copic@gozdis.si	Slovenian Forestry Institute, Slovenia
Čortan	Dijana	dijanacortan@yahoo.com	University of Novi Sad, Faculty of Education, Serbia
Cruz Houghton	Cristina Maria	ccruz@fc.ul.pt	Uni. de Lisboa, Faculdade de Ciências, CE3C, Portugal
Dighton	John	dighton@camden.rutgers.edu	Rutgers University, USA
Drolc	Tina	tina.drolc@gozdis.si	Slovenian Forestry Institute, Slovenia
Eler	Klemen	klemen.eler@bf.uni-lj.si	University of Ljubljana, Biotechnical faculty, Slovenia
Ernst	Dieter	ernst@helmholtz-muenchen.de	Helmholtz Zentrum München, Germany
Ferlan	Mitja	mitja.ferlan@gozdis.si	Slovenian Forestry Institute, Slovenia
Filipič	Arijana	arijana_filipic@hotmail.com	University of Ljubljana, Biotechnical faculty, Slovenia
Finžgar	Domen	macjezivljenje@gmail.com	Slovenian Forestry Institute, Slovenia
Fortič	Ana	anaa.fortic@gmail.com	University of Ljubljana, Biotechnical faculty, Slovenia
Galovic	Vladislava	vladislava.galovic@gmail.com	Institute of Lowland Forestry and Environment, Serbia
Gessler	Arthur	arthur.gessler@wsl.ch	Swiss Federal Research institute WSL, Switzerland
Gioahin	Erika	erika.gioahin@gmail.com	University of Ljubljana, Biotechnical faculty, Slovenia
Göransson	Hans	hans.goeransson@boku.ac.at	University of Natural Resources and Life Sciences (BOKU), Austria
Grce	Dragomir	dragomirgrce@gmail.com	University of Ljubljana, Biotechnical faculty, Slovenia
Grebenc	Tine	tine.grebenc@gozdis.si	Slovenian Forestry Institute, Slovenia
Gričar	Jožica	jozica.gricar@gozdis.si	Slovenian Forestry Institute, Slovenia
Gutierrez Merino	Emilia	emgutierrez@ub.edu	University of Barcelona, Spain
Hafner	Polona	polona.hafner@gozdis.si	Slovenian Forestry Institute, Slovenia
Helmisaari	Heljä - Sisko	helja-sisko.helmisaari@helsinki.fi	University of Helsinki, Dep. of Forest Sciences, Finland
Herman-Planinšek	Marija	drevsnica@omorika.si	Omorika d.o.o., Slovenia
Horemans	Joanna	joannahoremans@uantwerpen.be	University of Antwerp, Belgium
Horvat	Eva	eva79horvat@gmail.com	Student of University of Ljubljana, Slovenia
Hrenko	Melita	melita.hrenko@gozdis.si	Slovenian Forestry Institute, Slovenia
Hukić	Emira	e.hukic@gmail.com	Faculty of Forestry, University of Sarajevo
Ivanković	Mladen	mladeni@sumins.hr	Croatian Forest Research Institute, Croatia
Japelj	Anže	anze.japelj@gozdis.si	Slovenian Forestry Institute, Slovenia
Jurše	Ana	ana.jurse@gov.si	

Surname	First name	E-mail	Institute
Kajba	Davorin	davorin.kajba@zg.t-com.hr	Faculty of Forestry University of Zagreb, Croatia
Kandemir	Gaye	gayekandemir@ogm.gov.tr	Forest Tree Seeds and Tree Breeding Research Institute Directorate, Turkey
Katanić	Marina	marinakatanic44@gmail.com	Institute of Lowland Forestry and Environment, Serbia
Kavčič	Andreja	andreja.kavcic@gozdis.si	Slovenian Forestry Institute, Slovenia
Kayler	Zachary	kayler@zalf.de	ZALF, Germany
Kern	Anikó	anikoc@nimbus.elte.hu	Eötvös Loránd University, Hungary
King	John	john_king@ncsu.edu	Department of Forestry, North Carolina State University, USA
Kiourtsis	Fotios	fkiourts@gmail.com	Decentralised Administration of Macedonia & Thrace General Directorate of Forests and Rural Affairs, Macedonia
Kljun	Ivan	ivankljun@hotmail.com	Student, Slovenia
Kobal	Milan	milan.kobal@bf.uni-lj.si	University of Ljubljana, Biotechnical faculty, Slovenia
Konnert	Monika	monika.konnert@asp.bayern.de	Bavarian Institution for Forest Seeding and Planting, Germany
Kos	Ivan	ivan.kos@bf.uni-lj.si	Univ. of Ljubljana, Biotechnical faculty, Dep. of Biology, Slovenia
Kovač	Marko	marko.kovac@gozdis.si	Slovenian Forestry Institute, Slovenia
Kovič Dine	Maša	masa.kovic-dine@pf.uni-lj.si	Faculty of Law, University of Ljubljana, Slovenia
Kraigher	Hojka	hojka.kraigher@gozdis.si	Slovenian Forestry Institute, Slovenia
Krajnc	Damjan	damjan.krajnc@um.si	Faculty of Chemistry and Chemical Engineering, Slovenia
Kreft	Ivan	ivan.kreft@guest.arnes.si	Slovenian Forestry Institute, Slovenia
Kremer	Antoine	antoine.kremer@pierreton.inra.fr	INRA, France
Kržišnik	Davor	davor.krzisnik@bf.uni-lj.si	Univerza v Ljubljani, Biotehniška fakulteta, Slovenia
Kuehdorf	Katja	Katja.Kuehdorf@zalf.de	Leibniz Centre for Agricultural Landscape Research (ZALF), Germany
Kuralt	Žan	zan.kuralt@gmail.com	Univ. of Ljubljana, Biotechnical faculty, Dep. of Biology, Slovenia
Kutnar	Lado	lado.kutnar@gozdis.si	Slovenian Forestry Institute, Slovenia
Lavrič	Martina	martina.lavric@gozdis.si	Slovenian Forestry Institute, Slovenia
Lazarević	Jelena	ena.lazarevic@gmail.com	University of Montenegro, Biotechnical faculty, Montenegro
Leppalammi - Kujansuu	Jaana	jaana.leppalammi-kujansuu@helsinki.fi	University of Helsinki, Finland
Levanič	Tom	tom.levanic@gozdis.si	Slovenian Forestry Institute, Slovenia
Lipar	Žiga	ziga.lipar@gozdis.si	Slovenian Forestry Institute, Slovenia
Luthar	Zlata	zlata.luthar@bf.uni-lj.si	University of Ljubljana, Biotechnical Faculty, Slovenia
Makita	Naoki	naoki.makita@helsinki.fi	University of Helsinki, Finland
Marjanović	Hrvoje	hrvojem@sumins.hr	Croatian Forest Research Institute, Croatia
Martinović	Tijana	tijana.martinovic@gozdis.si	Slovenian Forestry Institute, Slovenia
Surname	First name	E-mail	Institute
Matovic	Bratislav	bratislav.matovic@gmail.com	University of Novi Sad, Institute of Lowland Forestry and Environment, Serbia
Matteucci	Giorgio	giorgio.matteucci@cnr.it	National Research Council, Inst. of Agroenvironmental and Forest Biology, Italy
Matyssek	Rainer	matyssek@wzw.tum.de	Technische Universität München, Germany
McCarroll	Danny	d.mccarroll@swansea.ac.uk	Swansea University, United Kingdom
McDowell	Nate	mcdowell@lanl.gov	1Los Alamos National Lab, Los Alamos, USA
Mrak	Tanja	tanja.mrak@gozdis.si	Slovenian Forestry Institute, Slovenia
Müller	Daniel Alexander	Daniel.Mueller@asp.bayern.de	Bayerisches Amt für forstliche Saat- und Pflanzenzucht (ASP), Germany
Nagy	Laszlo	Inagy@erti.hu	NARIC Forest Research Institute, Hungary

Surname	First name	E-mail	Institute
Ortas	Ibrahim	iortas@cu.edu.tr	Cukuroava University, Turkey
Peteh	Maja	maja.peteh@gozdis.si	Forestry Library - Slovenian Forestry Institute & Univ. of Ljubljana, Biotechnical faculty, Dep. of Forestry and Forest Resources, Slovenia
Pinuela Samaniego	Yasmin	yasmin.pinuela@hotmail.com	Slovenian Forestry Institute, Slovenia
Planinšek	Vladimir	drevesnica@omorika.si	Omorika d.o.o., Slovenia
Poljanšek	Simon	simon.poljansek@gozdis.si	Slovenian Forestry Institute, Slovenia
Popa	Ionel	popaicas@gmail.com	National Research and Development Institute for Silviculture, Romania
Rantaša	Boris	boris.rantasa@gozdis.si	Slovenian Forestry Institute, Slovenia
Rasztovits	Ervin	rasztovits.ervin@emk.nyme.hu	University of West Hungary, Hungary
Ravnjak	Blanka	blanka.ravnjak@gmail.com	University Botanic Gardens Ljubljana, Slovenia
Rebov	Irena	irena.rebov@gozdis.si	Slovenian Forestry Institute, Slovenia
Robek	Robert	robert.robek@gozdis.si	Slovenian Forestry Institute, Slovenia
Rupel	Matej	matej.rupel@gozdis.si	Slovenian Forestry Institute, Slovenia
Sankovič	Iztok	iztok.sankovic@gozdis.si	Slovenian Forestry Institute, Slovenia
Senčar	Natalija	natalija.sencar@gozdis.si	Slovenian Forestry Institute, Slovenia
Simončič	Primož	primoz.simoncic@gozdis.si	Slovenian Forestry Institute, Slovenia
Sinjur	Iztok	iztok.sinjur@gozdis.si	Slovenian Forestry Institute, Slovenia
Špenko	Magda	magda.spenko@gozdis.si	Slovenian Forestry Institute, Slovenia
Stanković	Milena	milenastankovic60@gmail.com	Univ. of East Sarajevo, Fac. of Agriculture, Dep. of Forestry, Bosnia and Herzegovina
Starr	Mike	mike.starr@helsinki.fi	University of Helsinki, Finland
Stojnic	Srdjan	srdjan_stojnic@yahoo.com	Institute of Lowland Forestry and Environment, Serbia
Štraus	Ines	ines.straus@gozdis.si	Slovenian Forestry Institute, Slovenia
Thaci	Bashkim	bashkimt2002@yahoo.com	Ministry of Agriculture, Forestry and Rural Development, Kosovo
Toromani	Elvin	elvintoromani@gmail.com	Agricultural University of Tirana, Albania
Unuk	Tina	tina.unuk@gmail.com	Slovenian Forestry Institute, Slovenia
Urbanek Krajnc	Andreja	andreja.urbanek@um.si	Faculty of Agriculture and Life Science, Univ. of Maribor, Slovenia
Vieira	Joana	joana.vieira@uc.pt	University of Coimbra, Portugal
Vilhar	Urša	ursa.vilhar@gozdis.si	Slovenian Forestry Institute, Slovenia
Vodnik	Dominik	dominik.vodnik@bf.uni-lj.si	Biotechnical faculty, University of Ljubljana, Slovenia
Vukovič	Polona	polona.vukovic@gozdis.si	Slovenian Forestry Institute, Slovenia
Westergren	Marjana	marjana.westergren@gozdis.si	Slovenian Forestry Institute, Slovenia
Zavadlav	Saša	sasa.zavadlav@gozdis.si	Slovenian Forestry Institute, Slovenia
Železnik	Peter	peter.zeleznik@gozdis.si	Slovenian Forestry Institute, Slovenia
Žlahtič	Mojca	mojca.zlahtic@bf.uni-lj.si	Univerza v Ljubljani, Biotehniška fakulteta, Slovenia
Žlindra	Daniel	daniel.zlindra@gozdis.si	Slovenian Forestry Institute, Slovenia























































































