

# Upper Miocene ostracods from the Krško Basin, SE Slovenia

Miha Marinšek<sup>1</sup>, Valentina Hajek-Tadesse<sup>2</sup>, Marijan Poljak<sup>3</sup>, Tea Kolar-Jurkovšek<sup>1</sup> and Luka Gale<sup>1,4</sup>

<sup>1</sup> Geological Survey of Slovenia, Dimičeva ul. 14, 1000 Ljubljana, Slovenia; (miha.marinsek@geo-zs.si, tea.kolar-jurkovsek@geo-zs.si, luka.gale@geo-zs.si)

<sup>2</sup> Croatian Geological Survey, Sachsova 2, 10 000 Zagreb, Croatia; (tadesse@hgi-cgs.hr)

<sup>3</sup> Zgornje Gorenjce 27, 8333 Semič; (marijanpoljak7@gmail.com)

<sup>4</sup> University of Ljubljana, Faculty of Natural Sciences and Engineering, Department of Geology, Aškerčeva cesta 12, 1000 Ljubljana, Slovenia; (luka.gale@ntf.uni-lj.si)

doi: 10.4154/gc.2023.07



## Abstract

The purpose of this study was to identify the ostracod assemblage from a 43 m thick section of the Bizeljsko Formation, which constitutes the middle part of the Upper Miocene Posavje Group. The succession comprises sandstone, siltstone, and marlstone, deposited in a delta front environment. The determined ostracods belong to the families Cypridae, Cytheridae, Darwinulidae, and Loxoconchidae. In total, 30 species were identified. Additionally, 8 morphotypes were determined at the genus level. The most common genera are *Candona*, *Hemicytheria*, and *Cyprideis*. The assemblage belongs to the *Caspiocypris labiata* subzone from the upper Pannonian. The ostracod assemblage from the Krško Basin is similar in species composition to Pannonian ostracod assemblages from Croatia and Serbia.

## Article history:

Manuscript received January 10, 2023

Revised manuscript accepted June 29, 2023

Available online July 11, 2023

**Keywords:** Lake Pannon, Bizeljsko Formation, Pannonian, biostratigraphy, palaeoenvironment


## 1. INTRODUCTION

The Central Paratethys disintegrated towards the end of the middle Miocene, leaving in its wake the Late Miocene Lake Pannon (MAGYAR et al., 1999; PILLER et al., 2007). The brackish Lake Pannon lost its marine life, and the largely endemic ostracod and molluscan fauna of the lake is well-known from its fossil record. Research on the ostracod fauna is thus extremely important in determining the relative age, the palaeoecological conditions, and the palaeogeographic connections between individual sub-basins of Lake Pannon (KRSTIĆ, 1972, 1973; SOKAČ, 1981, 1989; JIŘIČEK & RIHA, 1991; RUNDIĆ, 1997; TER BORGH et al., 2013).


This paper presents the ostracod assemblage recovered from a 43 m thick succession of the Upper Miocene Bizeljsko Formation near the village of Bizeljsko, situated on the NE hilly rim of the Krško Basin in SE Slovenia (Figs. 1, 2). The succession is well exposed along a forest road, starting at coordinates Lat: 45°59'49.10", Lon: 15°41'01.23" (Figs. 3, 4).

The relative age of the studied section and palaeosalinity were determined, based on ostracods.

Finally, the determined ostracod assemblage was compared to assemblages of the same age from the Paratethys area. Data presented here serve as an important contribution to the understanding of the distribution of ostracods in Lake Pannon as the last known scientific paper on Miocene ostracods from Slovenia was published in 1989 (STEVANOVIĆ & ŠKERLJ, 1989). A short



**Figure 1.** Palaeogeographic map of the Central Paratethys (Pannonian age) with the approximate location of Slovenia (SB: Styrian Basin; MZB: Mura-Zala Basin) modified after KOVAC et al., 2017).



**Figure 2.** The present-day distribution of sediments and sedimentary rocks of the tectono-stratigraphic units of the Central Paratethys in eastern Slovenia. BL Balaton Line, CNL Celje-Nadžkaniža Line, CTZ Celje Tectonic Zone, DL Donat Line, MHZ Middle Hungarian Tectonic Zone, PAL Periadriatic Line, SF Sava Fault, ŠF, Šoštanj Fault, SB Senovo Basin, KB Krško Basin. For an explanation of the tectonostratigraphic units (A1, A2, B1, B2) see text. The star symbol marks the position of the sampled section. Modified after JELEN & RIFELJ (2002).


description of ostracods from the Krško Basin and the Bizeljsko section are given by VESEL-LUKIĆ (2012) and VESEL-LUKIĆ et al., (2013, 2014). There was also a Masters thesis done on the studied section by MARINŠEK (2020).

## 2. GEOLOGICAL AND STRATIGRAPHIC SETTING


Slovenia is located in the western part of Central Paratethys, where the Transtethyan corridor was theorised to be located in the Badenian (BARTOL, 2009). Sediments of the Central Paratethys in Slovenia are predominantly found in the eastern part of the country and extend westwards towards central Slovenia in a

series of synclines that once formed interconnected basins (Fig. 2). The NE part of Slovenia was connected to the “Styrian” and Mura-Zala basins (JELEN & RIFELJ, 2002) and most of the SE part was connected to the North Croatian Basin (JELEN et al., 1992; PAVELIĆ, 2005). The structural development of eastern Slovenia resulted in four tectono-stratigraphic units (TTU) (JELEN et al., 1992). From north to south, these are: TTU A1, TTU A2, TTU B1, and TTU B2 (Fig. 2).

Unit A1 covers the area north of the Periadriatic fault zone, namely the Northern Karavanke, Pohorje, and the Pannonian Plain of northeastern Slovenia. Unit A2 is a narrow area between



**Figure 3.** Position of the studied area. The exact position of the section is shown by a dashed rectangle in Figure 2B. The section starts at coordinates Lat: 45°59'49.10", Lon: 15°41'01.23".



**Figure 4.** Field pictures of the studied section. A – bottom of the marl in the middle of the section (view towards the south); B – a detailed picture of the marl section; C – transition from the marl to sand; D – sand section in the upper part of the outcrop (Author: Marina Vesel-Lukić, 2012)

the Periadriatic Fault Zone and the Don Tectonic Zone, which includes the Southern Karavanke and Haloze areas. Unit B1 stretches from the Julian Alps to the Sava Tectonic Zone and continues to the area of Varaždin in Croatia. Unit B2 covers the entire south-eastern part of Slovenia, including the Krško basin and thus also the area of Bizeljско. After deposition of the Neogene sediments, the area underwent compression, resulting in the formation of the Sava Folds, a W-E extending series of antiforms and synforms (PLACER, 1999a). The general facies succession of the Miocene sediments in Slovenia is composed of alternating marl/marlstone, sand/sandstone with cobblestone, and in some places, there are beds of lignite (PLENIČAR, 1954; CIGIT, 1958; VONČINA, 1966; PLENIČAR, 1968; PREMUR & MARINČIČ, 1971; KUŠČER, 1973; ARSOVSKI et al., 1975; MIOČ & ŽNIDARČIČ, 1978; KALOPER 1984; ANIČIĆ & JURIŠA 1985a, b; ŽNIDARČIČ & MIOČ, 1987; MARKOVIĆ & MIOČ, 1988; MIOČ & ŽNIDARČIČ, 1989; DJURASEK, 1995; PLACER, 1999b; ŠRAM et al., 2015). In Slovenia, the evolution of Lake Pannon is best recorded in the Krško and Senovo basins (Fig. 2). Pannonian calcareous marls conformably overlie Sarmatian beds; which are overlain by “ostracod marls”. In some parts of the Krško Basin the Pannonian lies unconformably on the Sarmatian beds (POLJAK et al., 2016) The Pannonian succession ends with quartz sand and marly-clayey intercalations with lignite on top. The entire Neogene succession can be interpreted as a transition from marine to brackish and finally freshwater conditions (PAVŠIČ & HORVAT, 2009; IVANČIČ, 2021). Important fossil groups that have been investigated in the Upper Miocene

sediments of eastern Slovenia are ostracods (PIERAU, 1958; SOKAČ, 1981; STEVANOVIĆ & ŠKERLIJ, 1985, 1989; HAJEK-TADESSE, 2007; VESEL-LUKIĆ, 2012; VESEL-LUKIĆ et al., 2013, 2014; MARINŠEK, 2020), and molluscs (MIKUŽ, 2005, 2014; MIKUŽ et al., 2015).

The Bizeljско area corresponds to the northern limb of the Krško synclinerium and is a part of the B2 TTU (PLACER, 1999a, b; POLJAK, 2017). The geological maps of the study area were produced by ANIČIĆ & JURIŠA (1985a, b), and more recently by POLJAK et al. (2017). A simplified geological map is shown in Figure 5. The stratigraphic succession in the study area is summarized after POLJAK (2017) (Fig. 6). The oldest deposits of the Central Paratethys in the research area belong to the Badenian–Sarmatian Laško Formation. It comprises shallow marine coralline algal-rich limestone, the “Lithothamnium limestone”, and the deeper marine “Laško marl”. The Laško Formation is, after another prominent unconformity, followed by marls and sands of the deltaic Pannonian Drnovo Formation, which in its upper part passes into the Bizeljско Formation. The latter comprises quartz sand, marl, and subordinate sandstone and gravel, deposited within a delta front environment. The uppermost Pannonian Raka Formation almost entirely consists of quartz sand, which was also deposited in a deltaic environment. The Drnovo, Bizeljско, and Raka Formations together comprise the Posavje Group, which represents the final remnants of the Central Paratethys. The youngest sediments in the area are represented by Quaternary gravel (POLJAK, 2017).




Figure 5. A simplified geological map by POLJAK et al. 2017, slightly modified.

The Bizeljsko Formation is best exposed in the eastern Krško Basin, especially in the Bizeljsko area. The formation was defined by POLJAK (2017). It was previously also researched by LAPAJNE (1976), ŠIKIĆ et al. (1979), SKABERNE (1990), ŠKERLJ (1990), TRAJANOVA (2006), HAJEK-TADESSE (2007), VESEL-LUKIĆ et al. (2013, 2014); MARINŠEK (2020) and POLJAK (2017). The formation consists mainly of marls and sands, which alternate over short distances. Beds of sand locally show traces of slumping and/or contain rip-up clasts of marl. It reaches its thickest point (up to 800 m) in the wider Bizeljsko area. The age was first determined based on bivalves (*Congeria croatica*, *C. rhomboidea*, *C. zagrabiensis*, *Lymnocardium majeri*, *L. riegeli*, *Paradacna abichi* and *Paradacna okrugici*, as well as ostracods (ŠIKIĆ et al., 1979; HAJEK-TADESSE, 2007; POLJAK, 2017). POLJAK (2017) interpreted the succession as a delta front deposit based on individual layers of sand showing signs of syndepositional slumping with frequent layers containing rip-up marl clasts. It represents the part of the Lake Pannon evolution where it began to shrink and transitioned locally from brackish to freshwater conditions. The Bizeljsko Formation is completely missing in the western part of the Krško Basin. Instead, the upper Pannonian deposits are represented there by a much thinner succession of marls and sandy marls of the Drnovo Formation (POLJAK, 2017).

### 3. MATERIAL AND METHODS

The section of the Bizeljsko Formation was logged and sampled with a total stratigraphic length of 43 m (Fig. 7). In total, 38 samples were collected and subsequently processed in the laboratory. Individual samples weighed between 1 and 2 kg. Each sample was quartered to a weight of 250g – 500g. Samples were dissolved in a 5% hydrogen peroxide solution; samples that contained a large number of ostracod valves filled with sediment were put in an ultrasonic bath for 30 seconds. The dissolved material was then wet sieved using mesh sizes of 0.500 mm, 0.250 mm, 0.125 mm, and 0.063 mm. The specimens were hand-picked under a ZEISS Stemi 2000-C binocular magnifier with a magnification of  $\times 50$ . Approximately 100 adult ostracod valves were

collected from each sample. Picking was nonselective to preserve the original proportions of the species within the assemblage. This was done by collecting valves either until we reached 100 valves or until the sample was looked through fully. Selected specimens were photographed on the JEOL JSM 6490LV scanning electron microscope at the Geological Survey of Slovenia. For determining the families of ostracods, the suprageneric classification scheme proposed by MOORE (1961) and HORNE et al. (2002) was used, as they are widely used classification schemes for ostracods. Samples are stored at the Geological Survey of Slovenia under field numbers GeoZS B-019 – B-057. The diversity was calculated using the Shannon-Wiener index (H) for each sample (SHANNON & WEAVER, 1949).

### 4. RESULTS

The total stratigraphic thickness of the section is 43 m (Fig. 7). A thin gravel layer at the base of the section is overlain by alternating sands and marls in 3 m thickness, with each sand and marl layer being approximately 20 to 30 cm thick (with a 1-meter-thick unlogged interval). The following 8-metre-thick sand has a sharp erosional base and clay intraclasts at the bottom part. The sand grades upward into marl, again overlain by a 10-metre-thick, dominantly sandy unit with an erosional base at 13 m. This unit contains thin clay and marl layers and even pebbly intervals at 17 m and between 20 and 21 m (Fig. 7). The sand is overlain by a 7-metre-thick clay with sand intercalations. The top of the clay is eroded by a channel sandstone at 30 metres, with pebbles above the erosion surface. This sandstone is 7 metres thick and contains a few thin marl layers. Marl predominates in the uppermost 6 m of the outcrop (Fig. 7). A total of 30 ostracod species were determined (Figs. 8-10 + Supplement 1). In addition, 8 morphotypes were determined to the genus level. Approximately 70 % of the collected valves were intact, and the rest were fragmented. Only adult specimens were included in the palaeontological determination. Diversity is strongly controlled by lithology: higher in marlstone, and lower in more sandy facies and claystone. The maximum value of diversity is 2.29 in sample B-024 at the 4 metre point of the section (taken from clay lithoclasts), but on aver-




Figure 6. A general stratigraphic column of the Miocene deposits within the Krško Basin (Modified after POLJAK et al., 2017).




Figure 7. Sedimentary log and stratigraphic distribution of ostracod species in the section. The "diversity" column shows values of the Shannon-Wiener index as a measure of species richness and evenness.

age the Shannon-Wiener diversity index values are between 1.0 and 1.5 (Fig. 7).

The most abundant specimens are from the genera *Cyprideis* and *Hemicytheria*. *Loxoconcha* is present in most of the section but represents only a small percentage of the total assemblage.

Of the species observed here, the most important ones for determining the age were: *Amplocypris dorsobrevis* SOKAČ, *Caspiocypris alta* (ZALÁNYI) and *Camptocypris* aff. *acronasuta* (LIVENTAL).

## 5. DISCUSSION

### 5.1. Depositional environment

The studied section begins with sandy facies which suggests a higher energy environment. Indicators for this are mud chips and interbedded gravel. The environment changes at 22 metres from the beginning of the succession where we see a thick sequence of clay with thin layers of sand. At the 30 metre mark, the facies change again, with a rise in energy with an erosional channel in the clay, in which sand was deposited. The section ends with a lower energy environment indicated by marl deposition. These shifts in diversity and lithological changes correspond to rapidly changing conditions and lateral migration of the delta (DALRYMPLE et al., 1992).

The ostracod genera *Cyprideis*, *Hemicytheria*, *Leptocythere* and *Loxoconcha* in the lowermost sand layer (0-2 m) indicate brackish conditions. Ostracod diversity is low in this sandy part. Moving upwards through the succession the ostracod diversity fluctuates significantly. Diversity is highest in the marl-dominated part of the section, which can be explained by low-energy sedimentation and the reduced influence of fresh water. The diversity of the ostracod community reaches its highest value in sample B-024, which was taken from clay lithoclasts (Fig. 6). The species *Cyprideis* ex. gr. *pannonica* from that sample is considered by the authors to be reworked from older beds. Further along the section, the diversity of the ostracod assemblage decreases drastically. The low values of diversity between samples B-035 and B-043 are due to the sudden intake of sand. The presence of *Cyprideis*, *Hemicytheria*, and *Loxoconcha* indicates that beds from the upper part of the sequence were deposited under brackish conditions.

Previous research of the Bizeljsko Formation suggested deposition in a delta-front environment (POLJAK, 2017), and this interpretation is accepted here. It is to be noted that POLJAK (2017) does not describe any progradation and/or cyclic deposition, which would be typical for deltaic deposition as described and studied in SZTANÓ et al. (2016), SEBE et al. (2020) and ŠPELIĆ et al. (2023). This studied outcrop is too short to log any cyclic deposition apart from alternating marl and sand. According to HAJEK-TADESSE (2007), the composition of the ostracod community of the Bizeljsko profile, the preservation and abundance of ostracod shells depend on the change in the salinity of the lake water and the sedimentation. In general, the same author (HAJEK-TADESSE, 2007) concluded that several changes in lake water salinity and ostracod fauna can be distinguished on the Bizeljsko profile. At the beginning of the late Pannonian, water salinity increases to mesohaline values, while in the uppermost samples of the late Pannonian, the ostracod fauna of the Bizeljsko profile takes on a more freshwater character and indicates deposition near fresh-water input and a near-shore environment as stated in POLJAK (2017). We agree with the paleoenvironmental interpretation of both HAJEK-TADESSE (2007) and POLJAK (2017).

### 5.2. Biostratigraphy

There are quite a few ostracod biostratigraphic zonal schemes available for the Pannonian stage (e.g., POKORNÝ, 1944; BRESTENSKÁ, 1961; KRSTIĆ, 1973, 1985; JIŘIČEK, 1975, 1985; SOKAČ 1972, 1989; JIŘIČEK & RIHA 1991; and RUNDIĆ, 1998, 2006). According to OLTEANU (2011), the ostracod assemblages are well-differentiated only between two units, the Lower Pannonian and the »Portaferrian« (with *Congerina rhomboidea*), and most of the known species cannot be considered reliable index fossils. The same author concludes that the biostratigraphy of the brackish-water facies, generally and especially of the Pannonian Basin, is also difficult to determine. A similar view was expressed by KOVÁCS et al. (2016). The same authors conclude that different ostracod biostratigraphic schemes have been erected in different parts of the Pannonian Basin System, but they usually reflect local characteristics and differ significantly from each other.

The four most commonly used schemes for biozonation are presented in Fig. 8. Here, the biozonation by SOKAČ (1989) was used. The great majority of the investigated section can be placed within the *Caspiocypris labiata* subzone (Fig. 8), which is the oldest subzone in the Upper Pannonian (previously known as upper Pontian) *Bakunella dorsoarcata* zone (SOKAČ, 1972; KRSTIĆ 1973; SOKAČ, 1989). The subzone is defined by *Caspiocypris labiata* (Zalányi), *Caspiocypris alta* (Zalányi) and *C. Caspiocypris pontica* Sokač. Although the species *Caspiocypris labiata* itself is absent, the defined subzone contains the typical *Caspiocypris* form, such as *Caspiocypris alta*. In addition, *Amplocypris* aff. *dorsobrevis* SOKAČ, *Camptocypris* aff. *acronasuta*, *Typhlocypris* cf. *centropunctata*, *Loxoconcha schweyerei* are relatively abundant in this zone which is characteristic for this subzone. The *Camptocypris flectimarginata*, which reaches its acme in the *Camptocypris flectimarginata* subzone, rarely occurs in the studied section. However, the younger *Pontiella truncata* subzone is excluded due to the absence of the index species, as well as the lack of the usual accompanying species e.g., *Pontiella paracuminata* and *Hastacandona loczyi* (JIŘIČEK, 1983; SOKAČ, 1989). The subzone was chosen out of all the others in the literature, due to the proximity of the localities where the subzone was defined, in agreement with the results of previous research undertaken around the studied area (SOKAČ, 1981; HAJEK-TADESSE, 2007). Ostracods determined here only correspond to the described ostracod assemblages of the *Caspiocypris labiata* subzone.

The lowermost 3.5 m of the section does not contain species diagnostic of the *Caspiocypris labiata* subzone, nor are there any species that would indicate the presence of the preceding, lower level of upper Pannonian *Hemicytheria prisca* SOKAČ biozone. The latter ostracod zone was defined by KRSTIĆ (1973) based on the nominate species, and the presence of the species *Pontiella ilica* KRSTIĆ, *Zalanyiella longissima* KRSTIĆ, *Camptocypris acuta* SOKAČ, *Bakunella anae* SPADI (SPADI et al., 2019), *Cyprideis salopeki* SOKAČ, and *C. parallela* KRSTIĆ. In the investigated section, the species *Camptocypris acuta* SOKAČ and *B. anae* SPADI occur in the lower part. The stratigraphic ranges of these species, however, extend to the upper Pannonian. The lowermost 3.5 m of the section is therefore not defined in age.

The species *Bakunella anae*, characteristic of the upper Pannonian (previously known as the lower and upper Pontian) (SOKAČ, 1972), was also observed in the sample B-024. The sample was collected from a marl lens within a sandstone bed, which is, as mentioned before, was interpreted as a clast eroded

Standard Chronostratigraphy				Central Paratethys Lake Pannon							
Geomagnetic polarity	Epoch	Age/Stage		OSTRACOD BIOZONATION							
				after Stevanović et al., 1990	after Neubauer et al., 2015	Krstić, 1973	Sokač, 1989	Jiříček & Riha, 1991	Rundić, 2006		
5	C3	Pliocene	Zanclean	Pannonian	Portaferrian	Portaferrian	<i>Bakunella dorsoarcuata</i>	<i>Camptocypria flectimarginata</i>	<i>Bakunella dorsoarcuata</i>	<i>Caspiolla acranasuta</i> - <i>Bakunella dorsoarcuata</i>	<i>Bakunella dorsoarcuata</i>
6											
7	C3A	Late Miocene	Pannonian	Pannonian	Novorossian	Transdanubian	<i>Hemicytheria prisca</i>	<i>Hemicytheria prisca</i>	<i>Caspiocypris labiata</i>	<i>Caspiolla balcanica</i> - <i>Caspiolla lobata</i>	<i>Hemicytheria josephinae</i>
8											
9	C3B	Pliocene	Zanclean	Pannonian	Novorossian	Transdanubian	Pannonian relicts	Pannonian relicts	Pannonian relicts	Pannonian relicts	Pannonian relicts
9											

Figure 8. Four selected biozonation schemes proposed by different authors (KRSTIĆ, 1973; SOKAČ, 1989; JIŘIČEK & RIHA, 1991; RUNDIĆ, 2006) with the stratigraphic position of the study site.

from the underlying beds. Resedimentation of the fossils is also possible for samples B-035, B-037, and B-048 (see Fig. 7).

According to JIŘIČEK & RIHA (1991), the equivalent ostracod zone for the *Caspiocypris labiata* subzone is the lower Pontian Ostracod Zone 20 “*Candona (C.) balcanica*- *Candona (C.) lobata*” in which there is the first occurrence of *Bakunella dorsoarcuata*. However, in the biozonation set by JIŘIČEK (1983), the species *Caspiocypris labiata* is placed in the upper Pannonian Zone E 1-2. The correlation of biozones is further complicated by the different compositions of ostracod assemblages in various parts of the Central Paratethys (see POKORNÝ, 1944; BRESTENSKÁ, 1961; KRSTIĆ, 1973, 1985; JIŘIČEK, 1975, 1985; RUNDIĆ, 1998, 2006). Also, there is much confusion related to the ostracod taxonomy, which needs a serious revision (see OLTEANU, 2011; SPADI et al., 2019).


### 5.3. Palaeosalinity levels

The Pannonian microfauna is dominated by brackish to freshwater ostracods which tolerate an increase in salinity (Figs. 8 and 10). This suggests that the composition of the Lake Pannon ostracod assemblages is strongly facies dependent, as was previously

demonstrated for the Pannonian mollusc fauna (MAGYAR et al., 1999). Many of the detected ostracod species may tolerate a considerable range of salinity. The number of freshwater species diminishes when salinity surpasses 3‰, and the number of true brackish water species in the range from 3-10‰ is not numerous but the number of specimens is large (MORKHOVEN, 1962).

The important genera for determining palaeosalinity levels in the logged part of the Bizeljско Formation are *Darwinula*, *Candonopsis*, *Pontoniella*, *Lineocypris*, *Camptocyprina*, *Caspiocypris*, *Typhlocypris*, *Bakunella*, *Hemicytheria*, *Leptocythere*, *Amplocypris*, and *Loxoconcha* (TUNOĞLU, 2003). *Candonopsis* and *Darwinula* are specific to freshwater environments. According to Meisch (2000), recent *Darwinula* tolerates an increase in salinity up to a maximum of 15‰, and *Candonopsis* from a maximum salt content of 5 ‰. Considering the poor preservation of their valves in sample B-024 and their probable re-deposition they were excluded from the environmental interpretation. Palaeosalinity was determined to be in the range of brackish conditions. The interpretation is based on the presence of the genus *Hemicytheria*, which was abundant in the section (MORKHOVEN, 1962; MEISCH, 2000; FRENZEL & BOOMER, 2005).






**Figure 9.** SEM pictures of Pannonian ostracods from the Bizeljsko section. 1. *Amplocypris subacuta* ZALÁNYI (LV, external view, sample B-024, fragmented valve); 2. *Amplocypris* cf. *dorsobrevis* SOKAČ (LV, external view, sample B-051, fragmented valve); 3. *Herpetocyprilla* cf. *pannonica* ZALÁNYI (LV, external view, sample B-048, fragmented valve); 4. *Bakunella anae* (Vekua) SPADI (RV, external view, sample B-024), alternatively, possible juvenile of *B. dorsoarcuta*; 5a., 5b. *Bakunella dorsoarcuta* ZALÁNYI (a. LV, external view, sample B-024; b. RV, internal view, sample B-024); 6. *Camptocypria acuta* SOKAČ (LV, internal view, sample B-027); 7. *Camptocypria* cf. *acronasuta* LIVENTAL (RV, internal view, sample B-027); 8. *Camptocypria flectimarginata* SOKAČ (LV, external view, sample B-031); 9. *Camptocypria*? sp. (LV, external view, sample B-030); 10. *Candonopsis* cf. *arcana* KRSTIĆ (LV, external view, sample B-024); 11. *Caspiocypris alta* ZALÁNYI (LV, internal view, sample B-031); 12. *Lineocypris reticulata* (MÉHES) (LV, external view, sample B-057); 13. *Lineocypris hodonensis* POKORNÝ (RV, external view, sample B-027); 14. *Lineocypris zagabiensis* SOKAČ (RV, external view, sample B-041).

#### 5.4. Comparison between the ostracod fauna from the Pannonian deposits of the Krško Basin and other parts of the Paratethys

The ostracod fauna from the Upper Pannonian deposits of the Krško Basin, which includes taxa that were determined both here

and in previously published papers (SOKAČ, 1981; STEVANOVIĆ & ŠKERLJ, 1989; HAJEK-TADESSE, 2007), was compared to the ostracod faunas of the same age reported from similar localities in the Central Paratethys. The results are presented in Supplement 2.




**Figure 10.** SEM pictures of Pannonian ostracods from the Bizeljsko section. 1. *Lineocypris* cf. *granulosa* ZALÁNYI (LV, external view, sample B-024); 2. *Pontoniella sagittosa* KRSTIĆ (RV, external view, sample B-024); 3. *Pontoniella* sp. 1 MANDELSTAM (LV, external view, sample B-024); 4. *Pontoniella* sp. 2 (LV, sample B-036); 5. *Typhlocypris* cf. *centropunctata* SUZIN (LV, external view, sample B-038); 6. *Typhlocypris* sp. (HÉJJAS) (LV, external view, sample B-040); 7. *Cyprideis macrostigma* KOLLMANN (a. RV, external view, b. LV, internal view, sample B-036); 8. *Cyprideis macrostigma spinosa* SOKAČ (a. LV, external view, b. LV, internal view, sample B-032); 9. *Cyprideis* ex gr. *pannonica* MÉHES (RV, external view, sample B-024); 10. *Cyprideis* ex gr. *pannonica* MÉHES (RV, external view, sample B-027); 11. *Cyprideis* ex gr. *pannonica* MÉHES (RV, external view, sample B-020); 12. *Cyprideis* sp. 1 (RV, external view, sample B-023); 13. *Cyprideis* sp. 2 (LV, external view, sample B-023).

In the comparison, from the literature, we only included Upper Pannonian ostracod species. The ostracod assemblage from the Krško Basin is most similar to that from Mt. Medvednica as the locality is very close by. Out of 73 species found in the Krško Basin, in total, Croatia has 43 of the same species from similar deposits. The resemblance with Croatia was expected as it belongs to the same system of supposed basins and the location of the outcrop and samples are only a few kilometres away from the comparison locations studied on Mt. Medvednica and reported in the literature. Ostracod assemblages from Romania and Serbia have, respectively, 29 and 33 species in common with Krško Basin. The Vienna Basin only has 13 species similar to those re-

corded at Bizeljko. Regarding Serbia and Romania, the number of similar species indicates a good connection with the Central Paratethys.

Papers involving the older Pannonian ostracods (e.g., VRSALJKO, 1999; KOVÁCS et al., 2016; CSOMA et al., 2021), as well as Pannonian ostracods from Austria /Vienna Basin (GROSS, 2004; GROSS et al., 2008; STAREK et al., 2010) are not included in Supplement 2. Apart from these localities, some ostracods are the same species as the ones that migrated from Paratethys to the Palaeo-Mediterranean. These are *Caspiocypris alta* and *Euxinocythere (M.) cf. praeabaquana* (GLIOZZI et al., 2012).



**Figure 11.** SEM pictures of Pannonian ostracods from the Bizeljko section. 1. *Cyprideis* sp. 3 (LV, external view, sample B-023); 2. *Hemicytheria croatica* SOKAČ (LV, external view, sample B-024); 3. *Hemicytheria cf. marginata* SOKAČ (LV, external view, sample B-024); 4. *Hemicytheria cf. prisca* SOKAČ (RV, external view, sample B-030); 5. *Hemicytheria dubokensis* KRSTIĆ (LV, juvenile, external view, sample B-026); 6. *Amnicythere naca* (MÉHES) (LV, external view, sample B-052); 7. *Amnicythere* sp. (LV, external view, sample B-020); 8. *Leptocythere?* sp. (LV, external view, sample B-051); 9. *Euxinocythere (Maeotocythere) cf. praeabaquana* (LIVENTAL) (RV, external view, sample B-033); 10. *Darwinula cf. stevensoni* BRADY & ROBERTSON (LV, external view, sample B-024) fragmented due to handling of the specimen; 11. *Loxoconcha schweyeri* SUZIN (LV, external view, sample B-024); 12. *Loxoconcha cf. spinosa* SOKAČ (LV, external view, sample B-024).

## 6. CONCLUSIONS

The Pannonian ostracods studied from the 43 m thick lithological succession (Bizeljско Fm., Krško basin) are relatively well preserved and diverse. The determined species delineate the late Pannonian *Caspiocypris labiata* subzone. Palaeosalinity levels, as indicated by the ostracods, show brackish water conditions. Comparison with the ostracod assemblages from the region shows the highest similarity in species composition to the fauna from Croatia, less so with faunas from Serbia and Romania. Within the Bizeljско ostracod assemblage, two species are recognised that migrated from Paratethys to the Palaeo-Mediterranean during the Lago-Mare event of the Messinian Salinity Crisis.

## ACKNOWLEDGMENT

This work is part of the ongoing PhD research financially supported by the Slovenian Research Agency (ARRS), within the program group Regional Geology (PI-0011). We thank Marija PETROVIĆ for the preparation of the samples.

## REFERENCES

- ANIČIĆ, B. & JURIŠA, M. (1985a): Osnovna geološka karta SFRJ, 1:100.000, list Rogatec [*Basic Geological Map of the SFRY, 1:100000, sheet Rogatec* – in Slovenian].– Zvezni geološki zavod, Beograd.
- ANIČIĆ, B. & JURIŠA, M. (1985b): Osnovna geološka karta SFRJ 1:100.000 – Tolmač za list Rogatec [*Basic Geological Map of SFRY 1:100000 – Explanatory book for the Rogatec sheet* – in Slovenian].– Zvezni geološki zavod, Beograd, 76 p.
- ARSOVSKI, M., STOJKOVIĆ, M., MIHAILOV, V. & PETROVSKI, D. (1975): Summary of the geological and seismological investigation of the nuclear power plant „Krško“ location.– Institute of earthquake engineering and engineering seismology, University Kiril and Metod, Skopje, 33 p.
- AVANIĆ, R., PAVELIĆ, D., MIKNIĆ, M., BAKRAČ, K., KOVAČIĆ, M., VRSALJKO, D., PIKJIA, M., BRKIĆ, M., BASCH, O. & SARKOTIĆ ŠLAT, M. (2022b): Formacija Medvedski Breg (MEB) [*Medvedski Breg Formation (MEB)* – in Croatian].– In: HALAMIĆ, J. & BELAK, M. (eds.): Tumač za Požešku goru. [*Explanatory book for Požeška Gora* – in Croatian]. Hrvatski geološki institut, Zagreb, 122–123.
- BARTOL, M. (2009): Middle Miocene calcareous nannoplankton of NE Slovenia (western Central Paratethys).– Paleontološki inštitut Ivana Rakovca ZRC SAZU, Ljubljana, 142 p. doi: 10.3986/9789612545598
- BASCH, O., BRKIĆ, M., GRIZELJ, A., KOVAČIĆ, M., MIKNIĆ, M., SARKOTIĆ ŠLAT, M. (2022): Formacija Nova Gradiška (NGR) [*Nova Gradiška Formation (NGR)* – in Croatian].– In: HALAMIĆ, J. & BELAK, M. (eds.): Tumač za Požešku goru [*Explanatory book for Požeška Gora* – in Croatian]. Hrvatski geološki institut, Zagreb, 124–127.
- BARTHA, I.R., BOTKA, D., CSOMA, V., KATONA, L.T., TÓTH, E., MAGYAR, I., SILYE, L. & SZTANÓ, O. (2021): From marginal outcrops to basin interior: a new perspective on the sedimentary evolution of the eastern Pannonian Basin.– Int. J. Earth Sci. (Geol Rundsch), 111, 335–357. doi: 10.1007/s00531-021-02117-6
- BRESTENSKÁ, E. (1961): Present knowledge and problems of the Pliocene of the West Carpathians.– Geol. Bratislava, Prace, Zošit, 60, 149–157.
- CIGIT, K. (1958): O geoloških razmerah filovske nafne strukture [*The geological settings of the Filovci oil structure* – in Slovenian].– Geologija, 4, 171–187.
- CSOMA, V., MAGYAR, I., SZUROMI-KOPECZ, A., SEBE, K., SZTANÓ, O., BUCZKÓ, K. & TÓTH, E. (2021): Pannonian (late Miocene) ostracod fauna from Pécs-Danitzpuszta in Southern Hungary.– Foldtani Kozlony, 151/3, 305–326. doi: 10.23928/foldt.kozl.2021.151.3.305
- CZICZER, I., MAGYAR, I., PIPÍK, R., BÖHME, M., ČORIĆ, S., BAKRAČ, K., SÜTŐSZENTAI, M., LANTOS, M., BABINSZKI, E. & MÜLLER, P. (2008): Life in the sublittoral zone of long-lived Lake Pannon: paleontological analysis of the Upper Miocene Szák Formation, Hungary.– Int. J. Earth Sci. (Geol Rundsch), 98, 1741–1766. doi: 10.1007/s00531-008-0322-3
- DALRYMPLE, R.W., ZAITLIN, B.A. & BOYD, R. (1992): Estuarine facies models: conceptual basis and stratigraphic implications.– J. Sediment. Petrol., 62/6, 1130–1146. doi: 10.1306/D4267A69-2B26-11D7-8648000102C1865D
- DJURASEK, S. (1995): Interpretacija refleksijskih seizmičnih profila na Krškem polju (KP-83-59, KP-84-59, KP-85-59, KP-86-59) [*Interpretation of reflection seismic profiles in Krško polje (KP-83-59, KP-84-59, KP-85-59, KP-86-59)* – in Slovenian].– Uprava RS za jedrsko varnost, Geološki zavod Slovenije, Ljubljana.
- FRENZEL, P. & BOOMER, I. (2005): The use of ostracods from marginal marine, brackish waters as bioindicators of modern and Quaternary environmental change.– Palaeogeogr. Palaeoclimatol., 225, 68–92. doi: 10.1016/j.palaeo.2004.02.051
- FLOROIU, A., STOICA, M., VASILIEV, I. & KRIJGSMAN, W. (2011): Maeotian/Pontian ostracods in the Badislava – Topolog area (South carpathian foredeep-Romania).– Geo-Eco-Marina, 17, 237–244.
- GLIOZZI, E., GROSSI, F., COSENTINO, D. & IADANZA, A. (2012): The late Messinian Lago-Mare biofacies in central Apennines: the ostracod perspective.– Rend. Online Soc. Geol. It., 23, 63–65.
- GROSS, M. (2004): Zur Ostracodenfauna (Crustacea), Paläoökologie und Stratigraphie der Tongrube Mataschen (Unter-Pannonium, Steirisches Becken, Österreich).– Joannea Geol. Paläontol., 5, 49–130.
- GROSS, M., MINATI, K., DANIELOPOL, D. & PILLER, W. (2008): Environmental changes and diversification of Cyprideis in the Late Miocene of the Styrian Basin (Lake Pannon, Austria).– Palaeobio. Palaeoenv., 88, 161–181. doi: 10.1007/BF03043987
- HAJEK-TADESSE, V. (2007): Mikropaleontološke analize uzoraka profila Bizeljско [*Micropaleontological analyzes of Bizeljско profile samples* – in Croatian].– Unpubl. report, Hrvatski geološki institut, 15 p.
- HORNE, D., COHEN, A. & MARTENS, K. (2002): Taxonomy, Morphology and Biology of Quaternary and Living Ostracoda.– In: HOLMES, J.A. & CHIVAS, A.R. (eds.): The Ostracoda: Applications in Quaternary Research. Geophysical Monograph Series. 131 p. doi: 10.1029/131GM02
- IVANČIĆ, K. (2021): Pregled sedimentacijskega razvoja in formacij na območju Slovenije v miocenu [*Overview of sedimentary development and formations in the area of Slovenia in the Miocene* – in Slovenian].– In: ROŽIČ, B. (ed.): Treatises, reports = 25th Meeting of Slovenian Geologists. Univerza v Ljubljani, Naravoslovnotehniška fakulteta, Oddelek za geologijo, 31–35.
- JELEN, B., ANIČIĆ, B., BREZIGAR, A., BUSER, S., CIMERMAN, F., DROBNE, K., MONOSTORI, M., KEDVES, M., PAVŠIĆ, J. & SKABERNE, D. (1992): Model of positional relationships for Upper Paleogene and Miocene strata in Slovenia.– In: MONTANARI, A., COCCIONI, R. & ODIN, G.S. (eds.): Interdisciplinary Geological Conference on the Miocene Epoch with Emphases on the Umbria-Marche Sequence. International Union of Geological Sciences, Subcommittee on Geochronology, Ancona, 71–72.
- JELEN, B. & RIFELJ, H. (2002): Stratigrafska zgradba terciarne tektonostratigrafske enote B1 v vzhodni Sloveniji. Stratigraphic structure of Tertiary tectonic stratigraphic unit B1 in eastern Slovenia.– Geologija, 45/1, 115–138.
- JIRIČEK, R. (1975): Biozonal division of the Upper Tertiary basins of the Eastern Alps and west Carpathians RCMNSVI congress, IUGS comm.– Stratigr., 11–147.
- JIRIČEK, R. (1983): Redefinition of the Oligocene and Neogene Ostracod Zonation of the Paratethys.– Knjivnička Zemljiho plynu a nafty, 4, 195–236.
- JIRIČEK, R. (1985): Die Ostracoden des Pannonien.– Chronostratigraphie und Neostratotypen. Miozän der Zentralen Paratethys, 7, 378–425.
- JIRIČEK, R. & RIHA, J. (1991): Correlation of Ostracod Zones in the Paratethys and Tethys. – Saito Ho-on Kai Special Publications (Proceedings of Shallow Tethys), 3, 435–457.
- KALOPER, D. (1984): Krško polje-Brežice, analogna obrada profila KP-83-59, KP-84-59, KP-85-59, KP-86-59 [*Krško field - Brežice, analog processing of profile KP-83-59, KP-84-59, KP-85-59, KP-86-59* – in Slovenian].– Geofizika, 3 p., Zagreb.
- KRSTIĆ, N. (1972): Rod *Candona* (Ostracoda) iz kongerijskih slojeva južnog dela Panskog basena) [*Genus Candona (Ostracoda) from Congeria beds of southern Pannonian Basin* – in Serbian, with an English Summary].– Serbian Academy of Science and Arts, vol. CDL, Sec. Nat. Math. Sci., 39, 145 p., XXX Plates, Belgrade.
- KRSTIĆ, N. (1973): Biostratigrafija kongerijskih slojeva okoline Beograda na osnovu ostrakoda. Biostratigraphy of congerian layers around Belgrade based on ostracods.– Posebno izd. Inst. geol. geof. istr. ispitivanja nuklearnih i drugih mineralnih surovina, 4, 5–158.
- KRSTIĆ, N. (1985): Ostracoden im Pannonien der Umgebung von Belgrad.– In: PAPP, A., JÁMBOR, Á. & STEININGER, F.F. (eds.): Chronostratigraphie und Neostratotypen, Miozän der Zentralen Paratethys 7, Pannonien. Akadémiai Kiadó, Budapest, 103–144.
- KOVÁCS, E., MAGYAR, I., SZTANÓ, O. & PIPÍK, R. (2016): Pannonian ostracods from the southwestern Transylvanian basin.– Geol. Croat., 69/2, 213–229. doi: 10.4154/GC.2016.16
- KOVAC, M., HUDÁČKOVÁ, N., HALÁSOVÁ, E., KOVÁČOVÁ, M., HOLCOVÁ, K., OSZCZYPKO-CLOWES, M., BÁLDI, K., LESS, G., NAGYMAROSY, A., RUMAN, A., KLUČIAR, T. & JAMRICH, M. (2017): The Central Paratethys palaeoceanography: a water circulation model based on microfossil proxies, climate, and changes of depositional environment.– Acta Geol. Slovaca, 9/2, 75–114.
- KUŠČER, D. (1973): Summary of reports on preliminary geologic site investigations for the Krško Nuclear Power Plant.– Geološki zavod Ljubljana.
- LAPAJNE, V. (1976): Kremenov pesek Bizeljско [*Flint sand Bizeljско* – in Slovenian].– Unpubl. report, Ljubljana, Geološki zavod, 40 p.
- LAZAREV, S., DE LEEUW, A., STOICA, M., MANDIĆ, O., VAN BAAK, C.G.C., VASILIEV, I. & KRIJGSMAN, W. (2020): From Khersonian drying to Pontian “flooding”: late Miocene stratigraphy and palaeoenvironmental evolution of the Dacian Basin (Eastern Paratethys).– Glob. Planet. Change, 192, 03224. doi: 10.1016/j.gloplacha.2020.103224

- MAGYAR, I., GEARY, D.H. & MÜLLER, P. (1999): Palaeogeographic evolution of the Late Miocene Lake Pannon in Central Europe.– *Palaeogeogr. Palaeoclimatol., 147*, 151–167. doi: 10.1016/S0031-0182(98)00155-2
- MAGYAR, I., SZTANÓ, O., SEBE, K., KATONA, L., CSOMA, V., GÖRÖG, Á., TÓTH, E., KORECZ, A., ŠUJAN, M., BRAUCHER, R., RUSZKICZAY-RÜDIGER, Z., KOROKNAI, B., WÖRUM, G., SANT, K., KELDER, N. & KRJIGSMAN, W. (2019): Towards a high-resolution chronostratigraphy and geochronology for the Pannonian Stage: Significance of the Paks cores (Central Pannonian Basin).– *Földtani Közlöny*, 149, 351. doi: 10.23928/foldt.kozl.2019.149.4.351
- MARKOVIĆ, S. & MIOČ, P. (1988): Osnovna geološka karta SFRJ 1:100.000 (List Nadkaniža) [*Basic Geological Map of SFRJ 1:100000 L 33-58 (Sheet Nadkaniža)*] – in Slovenian]. – Zvezni geološki zavod Beograd.
- MARINŠEK, M. (2020): Ostrakodna združba zgornjemiocenskih plasti na Bizeljskem [*Ostracod fauna from upper Miocene beds from Bizeljsko*] – in Slovenian, Abstract in English]. – Master's Thesis. Univerza v Ljubljani. 108 p.
- MEISCH, C. (2000): Freshwater Ostracoda of Western and Central Europe.– Spektrum, Akademischer Verlag (Gustav Fischer), Heidelberg, 1st edition, 522 p.
- MIKUŽ, V. (2005): Pannonian molluscs from surroundings of Čanje near Sevnica, Slovenia.– *Geologija*, 48/2, 225–243. doi: 10.5474/geologija.2005.019
- MIKUŽ, V. (2014): Some rare fossils from Slovenske gorice, Slovenia.– *Geologija*, 57/2, 155–166. doi: 10.5474/geologija.2014.013
- MIKUŽ, V., KRIŽNAR, M. & CAF, N. (2015): Pannonian molluscs from site Osek-2 in Slovenske gorice, Slovenia.– *Geologija*, 58/1, 71–80. doi: 10.5474/geologija.2015.006
- MIOČ, P. & ŽNIDARČIČ, M. (1978): Osnovna geološka karta SFRJ 1:100.000 L 33-55 (List Slovenj Gradec) [*Basic Geological Map of SFRJ 1:100,000 L 33-55 (List = Sheet Slovenj Gradec)*] – in Slovenian]. – Zvezni geološki zavod, Beograd, 74 p.
- MIOČ, P. & ŽNIDARČIČ, M. (1989): Osnovna geološka karta SFRJ 1:100.000 (Tolmač Maribor in Leibnitz) [*Basic Geological Map of SFRJ 1:100,000 (Explanatory book Maribor and Leibnitz)*] – in Slovenian]. – Zvezni geološki zavod, Beograd, 60 p.
- MOORE, R.C. ed. (1961): Treatise on invertebrate paleontology. Part Q, Arthropoda. 3, Crustacea, Ostracoda.– Boulder; Lawrence: Geological Society of America; University of Kansas.
- MORKHOVEN, VAN F.P.C.M. (1962): Post-Paleozoic Ostracoda, their morphology, taxonomy and economic use, Part 2.– Elsevier, New York, 478 p.
- OLTEANU, R. (2011): Atlas of the Pannonian and Pontian Ostracods from the Eastern Area of the Pannonian Basin.– *Geo-Eco-Marina*, 17, 135–177. doi: 10.5281/zenodo.45062
- PAVELIĆ, D. (2005): Cyclicity in the evolution of the Neogene North Croatian Basin (Pannonian Basin System).– In: MABESOONE, J.M. & NEUMANN, V.H. (eds.): *Cyclic Development of Sedimentary Basins. Developments in Sedimentology*, vol. 57. Elsevier, Amsterdam, 273–284. doi: 10.1016/S0070-4571(05)80011-1
- PAVŠIČ, J. & HORVAT, A. (2009): Eocen, oligocen in miocen v osrednji in vzhodni Sloveniji, Eocen in oligocen [*The Eocene, Oligocene and Miocene in Central and Eastern Slovenia, Eocene and Oligocene*] – in Slovenian and English].– In: PLENIČAR, M., OGORLEČ, B. & NOVAK, M. (eds.): *Geologija Slovenije [The Geology of Slovenia]* – in Slovenian and English], Geološki zavod Slovenije, 373–390.
- PIERAU, H. (1958): Zur Stratigraphie Und Tektonik Jungtertiärer Ablagerungen Im Nordwestlichen Krško Polje in Slovenien [*On the stratigraphy and tectonics of Late Tertiary deposits in the northwestern Krško Polje in Slovenia*] – in German].– *Geologija*, 4, 111–148.
- PILLER, W.E., HARZHAUSER, M. & MANDIĆ, O. (2007): Miocene Central Paratethys stratigraphy – current status and future directions.– *Stratigraphy*, 4, 2/3, 151–168.
- PLENIČAR, M. (1954): Obmurska naftna nahajališča [*Oil fields of the Mura River area*] – in Slovenian]. – *Geologija*, 2, 36–93.
- PLENIČAR, M. (1968): Osnovna geološka karta SFRJ (List Goričko) [*Basic Geological Map of SFRJ 1:100.000 L 33-45 (Sheet Goričko)*] – in Slovenian]. – Zvezni geološki zavod, Beograd.
- PLACER, L. (1999a): Prispevek k makrotektonski rajonizaciji mejnega ozemlja med Južnimi Alpami in Zunanji Dinaridi [*Contribution to the macrotectonic regionalization of the borderland between the Southern Alps and the Outer Dinarides*] – in Slovenian]. – *Geologija*, 41/1, 223–243.
- PLACER, L. (1999b): Strukturni pomen Posavskih gub [*Structural significance of the Posavina folds*] – in Slovenian, with an English Abstract]. – *Geologija*, 41/1, 191–221.
- POKORNÝ, V. (1944): La microstratigraphie du Pannonien entre Hodonin et Mikulčice (Moravie meridionale, Tchécoslovaquie) [*The Pannonian microstratigraphy between Hodonin and Mikulčice (South Moravia, Czechoslovakia)*] – in French].– *Bull. Inter. Acad. tcheque. Sci.*, 23, 1–25.
- POLJAK, M. (2017): Geološka karta vzhodnega dela Krške kotline 1:25.000, Tolmač [*Geological map of the eastern part of the Krško basin 1:25.000: explanatory booklet*] – in Slovenian]. – Geološki zavod Slovenije, Ljubljana, 109 p.
- POLJAK, M., RIŽNAR, I., DEMŠAR, M., NOVAK, M. & TOMAN, M. (2017): Geološka karta vzhodnega dela Krške kotline 1: 25.000 [*Geological map of the eastern part of the Krško basin 1:25.000*]. – Geološki zavod Slovenije, Ljubljana.
- PREMRU, U. & MARINČIČ, S. (1971): Tektonska zgradba jugovzhodne Slovenije in severozahodne Hrvaške, Študija za projekt nuklearne elektrarne v Krškem [*Tectonic structure of southeastern Slovenia and northwestern Croatia, Study for the nuclear power plant project in Krško*] – in Slovenian]. – Geološki zavod Ljubljana, 24 p.
- RUNDIĆ, L. (1997): Biostratigrafija kaspibrakičnog neogena Kolubarskog basena [*Biostratigraphy of the Caspibrackish Neogene of the Kolubara Basin*] – in Serbian, with an English Abstract]. – Unpubl. PhD Thesis, Faculty of Mining and Geology, University of Belgrade, 128 p.
- RUNDIĆ, L. (1998): The carapace of fossil ostracodes: paleoenvironmental indicators.– *Ann. Géol. Pénins. Balk.*, 62, 165–177.
- RUNDIĆ, L. (2006): Late Miocene ostracodes of Serbia: Morphologic and paleoenvironmental considerations.– *Ann. Géol. Pénins. Balk.*, 67, 89–100. doi: 10.2298/GAB-P0667089R
- RUNDIĆ, L., GANIĆ, M., KNEŽEVIĆ, S. & SOLIMAN, A. (2011): Upper Miocene Pannonian sediments from Belgrade (Serbia): new evidence and paleoenvironmental considerations.– *Geol. Carpath.*, 62/3, 267–278. doi: 10.2478/v10096-011-0021-z
- RUNDIĆ, L., GANIĆ, M., KNEŽEVIĆ, S., RADIVOJEVIĆ, D. & RADONJIĆ, M. (2019): Stratigraphic implications of the Mio-Pliocene geodynamics in the area of Mt. Avala: new evidence from Torlak Hill and Beli Potok (Belgrade, Serbia).– *Geol. Croat.*, 72/2, 109–128. doi: 10.4154/gc.2019.11
- SEBE, K., KOVAČIĆ, M., MAGYAR, I., KRIZMANIĆ, K., ŠPELIĆ, M., BIGUNAC, D., SÜTŐ-SZENTAI, M., KOVÁCS, A., SZUROMI-KORECZ, A., BAKRAČ, K., HAJEK-TADESSE, V., TROSKOT-ČORBIĆ, T. & SZTANÓ, O. (2020): Correlation of upper Miocene-Pliocene Lake Pannon deposits across the Drava Basin, Croatia and Hungary.– *Geol. Croat.*, 73/3, 177–195. doi: 10.4154/gc.2020.12
- SHANNON, C.E. & WEAVER, W. (1949): *The mathematical theory of communication*.– The University of Illinois Press, Urbana, 117 p.
- SKABERNE, D. (1990): Makroskopski opis jedrovanih odsekov vrtnice DRN-1 (Drnovo na Krškem polju) [*Macroscopic description of cored sections of well DRN-1 (Drnovo na Krško polje)*] – in Slovenian]. – In: KRANJČIĆ et al. (eds.): *Končno poročilo o geoloških raziskavah na Krškem polju za potrebe podzemnega skladiščenja plina – Vrtnica DRN-1/89 [Final report on geological research in the Krško polje for the needs of underground gas storage – Well DRN-1/89]* – in Slovenian]. Ljubljana, Geološki zavod Slovenije, 19 p.
- SOKAČ, A. (1972): Pannonian and Pontian ostracode fauna of Medvednica.– *Pal. Jugoslav.*, 11, 1–140.
- SOKAČ, A. (1981): Pannonian and Pontian ostracode fauna of the South-eastern slopes of Orlica and Krško polje.– *Geol. vjesnik*, 33, 95–99.
- SOKAČ, A. (1989): Pontian Ostracod fauna in the Pannonian basin.– In: MALEZ, M. & STEVANOVIĆ, P. (eds.): *Chronostratigraphie und Neostatotypen, Pliozan, Pontien*, bd. VIII. Jugosl. Akad. D. Wissen. Und Kunste, 672–700.
- SPADI, M., GLIOZZI, E., BOOMER, I., STOICA, M. & ATHERSUCH J. (2019): Taxonomic harmonization of Neogene and Quaternary candonid genera (Crustacea, Ostracoda) of the Paratethys.– *J. Syst. Palaeontol.*, 17/19, 1665–1698. doi: 10.1080/14772019.2018.1545708
- STAREK, D., PIPÍK, R. & HAGAROVÁ, I. (2010): Meiofauna, trace metals, TOC, sedimentology, and oxygen availability in the Late Miocene sublittoral deposits of Lake Pannon.– *Facies*, 56, 369–384. doi: 10.1007/s10347-009-0208-2
- STEVANOVIĆ, P. & ŠKERLJ, Ž. (1985): Prilog biostratigrafiji panonsko-pontskih sedimentata u okolini Videm-Krškog [*Annex to the biostratigraphy of the Pannonian-Pontic sediments in the vicinity of Videm-Krško*] – in Croatian]. – *Razprave IV*, 281–304.
- STEVANOVIĆ P. & ŠKERLJ, Ž. (1989): The Pontian sediments of Slovenia.– In: MALEZ, M. & STEVANOVIĆ, P. (eds.): *Chronostratigraphie und Neostatotypen, Pliozan, Pontien*, bd. VIII. Jugosl. Akad. D. Wissen. Und Kunste, 153–179.
- SZTANÓ, O., KOVÁČ, M., MAGYAR, I., ŠUJAN, M., FODOR, L., UHRIN, A., RYBÁR, S., CSILLAG, G. & TÖKÉS, L. (2016): Late Miocene sedimentary record of the Danube/Kisalföld Basin: interregional correlation of depositional systems, stratigraphy and structural evolution.– *Geol. Carpath.*, 67/6, 525–542. doi: 10.1515/geoca-2016-0033
- SZUROMI-KORECZ, A., SUTO-SZENTAI, M. & MAGYAR, I. (2004): Biostratigraphic revision of the HÓD-I well: Hungary's deepest borehole failed to reach the base of the Upper Miocene Pannonian stage.– *Geol. Carpath.*, 55/6, 475–485.
- ŠIKIĆ, K., BASCH, O. & ŠIMUNIĆ, A. (1979): Osnovna geološka karta SFRJ 1:100000, tumač za list Zagreb [*Basic Geological Map of the SFRJ 1:100000, Explanatory book for the Zagreb sheet*] – in Croatian]. – Geol. Zavod Zagreb, Savezni Geol. Zavod, Beograd, 81.
- ŠKERLJ, Ž. (1990): Mikropaleontološke analize in biostratografska interpretacija, Vrtnica DNR-1, (Interval 194–50m) [*Mikropaleontological analyzes and biostratigraphic interpretation, Well DNR-1, (Interval 194–50m)*] – in Slovenian]. – In: KRANJČIĆ et al. (eds.): *Končno poročilo o geoloških raziskavah na Krškem polju za potrebe podzemnega skladiščenja plina – Vrtnica DRN-1/89 [Final report on the geological research in the Krško basin of the need for underground gas storage – Well DRN-1/89]* – in Slovenian]. Unpubl. report, Ljubljana, GeoZS, 19 p.

- ŠPELIĆ, M., KOVÁCS, Á., SAFTIĆ, B. & SZTANÓ, O. (2023): Competition of deltaic feeder systems reflected by slope progradation: a high-resolution example from the Late Miocene-Pliocene, Drava Basin, Croatia.– *Int. J. Earth Sci. (Geol. Rundsch)*, 112, 1023–1041. doi: 10.1007/s00531-023-02290-w
- ŠRAM, D., RMAN, N., RIŽNAR, I. & LAPANJE, A. (2015): The three-dimensional regional geological model of the Mura-Zala Basin, northeastern Slovenia.– *Geologija*, 58/2, 139–154. doi: 10.5474/geologija.2015.011
- TER BORGH, M., VASILIEV, I., STOICA, M., KNEŽEVIĆ, S., MATENCO, L., KRJGSMAN, W., RUNDIĆ, L. & CLOETINGH, S. (2013): The isolation of the Pannonian basin (Central Paratethys): New constraints from magnetostratigraphy and biostratigraphy.– *Global. Planet. Change.*, 103, 99–118. doi: 10.1016/j.gloplacha.2012.10.001
- TRAJANOVA, M. (2006): Granulometrijska in litološka sestava vzorcev proda iz lokacije Vrbina (Sv. Urh in Libna) [*Granulometric and lithological composition of gravel samples from the Vrbina location (Sv. Urh and Libna)* – in Slovenian].– Unpubl. report, Geološki zavod Slovenije, Ljubljana, 9 p.
- TUNOĞLU, C. (2003): Systematics and biostratigraphy of the Pontian Candonidae (Ostracoda) from the eastern Black Sea region (Northern Turkey).– *Geol. Carpath.*, 54/1, 21–40.
- VESEL-LUKIČ, M. (2012): Panonijska in pontijska ostrakodna favna Krške kotline (vzhodna Slovenija, Centralna Paratetida) [*Pannonian and Pontic ostracod fauna of the Krška basin (eastern Slovenia, Central Paratetida)* – in Slovenian].– In: ATANACKOV, J., JAMŠEK RUPNIK, P. & MARKIČ, M. (eds.): Abstract book. Geološki Zavod Slovenije, Ljubljana, p. 21.
- VESEL-LUKIČ, M., HAJEK-TADESSE, V. & POLJAK, M. (2013): Late Miocene ostracoda from Bizeljsko section (eastern Slovenia).– In: *Il Naturalista siciliano Palermo: Tipolitografia luxograph s.r.l.*, 413–414.
- VESEL-LUKIČ, M., HAJEK-TADESSE, V. & POLJAK, M. (2014): Paleoeкологија згорњемiocenskih ostrakodov profila Bizeljsko [*Palaeoecology of Upper Miocene Ostracods from Bizeljsko section* – in Slovenian].– In: ROŽIČ, B., VERBOVŠEK, T. & VRABEC, M. (eds.) (2014): Povzetki in ekskurzije [*Abstracts and field trips* – in Slovenian]. Univerza v Ljubljani, Naravoslovnotehniška fakulteta, Ljubljana, 73–74.
- VONČINA, Z. (1966): Strukturni i litofacijelni odnos Lendavske formacije u Mursko sobotskoj zoni [*Structural and lithofacial relations of Lendava Formation in the Murska Sobota zone* – in Serbo-Croatian].– INA Naftaplin Zagreb, Archive GeoZS, 37 p.
- VRSALJKO, D. (1999): The Pannonian Palaeoecology and Biostratigraphy of Molluscs from Kostanjek – Medvednica Mt., Croatia.– *Geol. Croat.*, 52/1, 9–27.
- ŽNIDARČIČ, M. & MIOČ, P. (1987): Osnovna geološka karta SFRJ 1:100.000 L 33-56 in L 33-44, list Maribor in Leibnitz [*Basic Geological Map of SFRJ 1:100000 L 33-56 and L 33-44, sheets Maribor and Leibnitz* – in Slovenian].– Zvezni geološki zavod, Beograd.

**Appendix:****Supplement 1.** Systematic list of determined ostracod species in this study.

- Class Ostracoda LATREILLE, 1802  
 Subclass Podocopa SARS, 1866  
 Order Podocopida SARS, 1866  
**Suborder Cypridocopina BAIRD, 1845**  
**Superfamily Cypridoidea BAIRD, 1845**  
**Family Cyprididae BAIRD, 1845**  
*Amplocypris* ZALÁNYI, 1944  
*Amplocypris subacuta* ZALÁNYI, 1944  
*Amplocypris* cf. *dorsobrevis* SOKAČ, 1972  
**Subfamily Herpetocypridinae BRONSTEIN, 1947**  
*Herpetocyprilla* DADAY, 1909  
*Herpetocyprilla* cf. *pannonica* ZALÁNYI, 1959  
**Family Candonidae KAUFMANN, 1900**  
**Subfamily Candoninae KAUFMANN, 1900**  
*Bakunella* SCHNEIDER, 1958  
*Bakunella anae* (VEKUA) SPADI, 2019  
*Bakunella dorsoarcuta* ZALÁNYI, 1929  
*Camptocypria* ZALÁNYI, 1959  
*Camptocypria acuta* SOKAČ, 1972  
*Camptocypria* cf. *acronasuta* (LIVENTAL, 1929)  
*Camptocypria flectimarginata* SOKAČ, 1967  
*Camptocypria?* sp.  
*Candonopsis* VAVRA, 1891  
*Candonopsis* cf. *arcana* KRSTIĆ, 1968  
*Caspiocypris* MANDELSTAM in SCHNEIDER et al., 1956  
*Caspiocypris alta* (ZALÁNYI, 1929)  
*Lineocypris* ZALÁNYI, 1929  
*Lineocypris reticulata* (MÉHES, 1907)  
*Lineocypris hodonensis* POKORNÝ, 1952  
*Lineocypris zagrabiensis* SOKAČ, 1972  
*Lineocypris* cf. *granulosa* ZALÁNYI, 1959  
*Pontoniella* MANDELSTAM in LUEBIMOVA et al., 1960  
*Pontoniella sagittosa* KRSTIĆ, 1968  
*Pontoniella* spp.
- Typhlocypris* VEJDOVSKÝ, 1882 (sensu NAMIOTKO et al. 2014)  
*Typhlocypris* cf. *centropunctata* (SUZIN, 1956)  
*Typhlocypris* sp.
- Superfamily Cytheroidea BAIRD, 1850**  
**Family Cytherideidae SARS, 1925**  
**Subfamily Cytherideinae SARS, 1925**  
*Cyprideis* JONES, 1857  
*Cyprideis macrostigma* KOLLMANN, 1958  
*Cyprideis macrostigma spinosa* SOKAČ, 1972  
*Cyprideis* ex. gr. *pannonica* (MÉHES, 1908)  
*Cyprideis* spp.
- Family Hemicytheridae PURI, 1953**  
**Subfamily Hemicytherinae PURI, 1953**  
*Hemicytheria* POKORNÝ, 1955  
*Hemicytheria croatica* SOKAČ, 1963  
*Hemicytheria* cf. *marginata* SOKAČ, 1972  
*Hemicytheria* cf. *prisca* SOKAČ, 1972  
*Hemicytheria dubokensis* KRSTIĆ, 1963
- Family Leptocytheridae SARS, 1925**  
*Amniccythere* DEVOTO, 1965  
*Amniccythere naca* (MÉHES, 1909)  
*Amniccythere* sp.  
*Leptocythere* SARS, 1925  
*Leptocythere* sp.  
*Euxinocythere (Maeotocythere)* STANCHEVA, 1968  
*Euxinocythere (M.)* cf. *praebaquana* (LIVENTAL, 1956)
- Family Loxoconchidae SARS, 1925**  
*Loxoconcha* SARS, 1866  
*Loxoconcha schweyeri* SUZIN, 1956  
*Loxoconcha* cf. *spinosa* SOKAČ, 1972
- Superfamily Darwinuloidea BRADY & ROBERTSON, 1885**  
**Family Darwinulidae BRADY & ROBERTSON, 1885**  
*Darwinula* BRADY & ROBERTSON, 1885  
*Darwinula* cf. *stevensoni* BRADY & ROBERTSON, 1870

