

LUNG SCAN INTERPRETATION – COMPARISON OF DIFFERENT CRITERIA

Budihna N¹, Milčinski M¹, Šuštaršič J², Grmek M¹, Grošelc C¹, Porenta M¹

Abstract – Authors reevaluated lung perfusion scans performed for suspected pulmonary embolism in 1988 and 1989 using new criteria proposed by international prospective study for pulmonary embolism diagnosis (PIOPED (5)).

The difference between the first reading and the reevaluation was found in 200 (25,35%) of patients examined by lung perfusion scanning using 99mTc-macro-aggregated albumin (99mTc-MAA). The probability of lung embolism increased in 185 (92,5%) of patients where the disagreement between the first and the second reading was found. The degree of discordance is low in the patients under 30 years of age (9,59%) and in the interpretation of normal lung perfusion scan (2,14%).

Ventilation lung scanning by 99mTc-diethylenetriaminepentaacetate aerosol (99mTc – DTPA), which was performed after perfusion lung scanning in 59 patients, influenced the diagnosis in 14 (23,72 %) of patients by lowering the predicted probability of lung embolism as assessed by perfusion scanning alone.

The systematic two-observer approach to lung perfusion scan analysis using PIOPED criteria is more sensitive for the diagnosis of pulmonary embolism than single-observer method. Ventilation scanning helps to avoid overdiagnosis of lung embolism specially in intermediate and high probability of pulmonary embolism.

UDC: 616.24-005.7:539.163

Key words: pulmonary embolism–radionuclide imaging

Orig sci paper

Radiol gosl 1990; 24:245-8.

Introduction – Diagnosis of pulmonary embolism (PE) is difficult if only clinical and biochemical criteria are considered. Lung scintigraphy is one of the most important noninvasive imaging techniques. The interpretation of lung scans is liable to errors. As it has been proven with prospective studies, the systematic approach is most successful in the correct diagnosis of PE (1, 2, 3, 4). Since the systematic approach has not been accepted in our laboratory before, we compared the results of nonsystematic lung scan interpretation to the approach suggested by National Heart, Lung and Blood Institute in USA (5).

Patients and methods – Lung perfusion scanning was performed in 789 patients (pts) because of clinically suspected pulmonary embolism (PE) in the period of two years (from 1988 to 1989). The pts were from 10 to 90 years old, 448 (57,5%) were females.

Lung perfusion scanning was performed with a large view field gamma camera after intravenous application of 99mTc-human serum albumin macroaggregates (99mTc-MAA). During the injection the pts were supine, but the scans were taken in the sitting position whenever there were

no serious contraindications. The scans were performed mostly in four projections: anterior, posterior, left and right posterior oblique. Few pts had scans taken only in anterior and both anterior oblique projections.

Ventilation lung scanning by 99mTc-diethylenetriaminepentaacetate aerosol (99mTc-DTPA) was performed after perfusion lung scanning in 59 patients in the same positions as perfusion scanning.

At the first diagnostic reading the scans were evaluated by several nuclear medicine physicians (sometimes they were less experienced) without strict criteria for lung scan interpretation. When ventilation lung scanning was not available, a recent thorax radiograph was compared to the lung scan serving as an estimation of the lung ventilation in the pts.

On reevaluation PIOPED criteria were used (Table 1). These are essentially modified Biello's criteria, known as PIOPED (prospective study of pulmonary embolism diagnosis (5)). According to them, the scans were assessed by two observers as normal, concordant with low, intermediate or high probability of lung embolism.

Table 1 – PLOPED criteria for diagnostic reading of lung perfusion scans

PROBABILITY OF PE	PIOPED CRITERIA
NORMAL	NORMAL PERFUSION
LOW	(1) SMALL Q DEFECTS REGARDLESS OF NUMBER, V OR CXR FINDINGS (2) Q DEFECT SUBSTANTIALLY SMALLER THAN CXR DEFECT (V IRRELEVANT) (3) V-Q MATCH IN $\leq 50\%$ ONE LUNG OR $\leq 75\%$ OF ONE LUNG ZONE CXR NORMAL OR NEARLY NORMAL (4) SINGLE MODERATE Q WITH NORMAL CXR (V IRRELEVANT) (5) NONSEGMENTAL Q DEFECT
INTERMEDIATE	(1) ABNORMALITY THAT IS NOT DEFINED BY EITHER »HIGH« OR »LOW«
HIGH	(1) TWO OR MORE LARGE Q. V AND CXR NORMAL (2) TWO OR MORE LARGE Q IN WHICH Q IS SUBSTANTIALLY LARGER THAN EITHER MATCHING V OR CXR (3) TWO OR MORE MODERATE Q AND ONE LARGE Q. V AND CXR NORMAL (4) FOUR OR MORE MODERATE Q. V AND CXR NORMAL

Legend: Q = perfusion, V = ventilation, CXR = chest X ray (chest roentgenogram)

Table 2 – The degree of discordance between the first and the second interpretation of lung perfusion of lung perfusion scans according to the age of the patients

AGE (YEARS)	No OF PTS	No OF DISCORDANT INTERPRETATIONS	% OF DISCORDANCE
≤ 29	73	7	9.59
30 - 39	83	22	26.51
40 - 49	119	28	23.54
50 - 59	136	37	27.21
60 - 69	189	50	26.46
≥ 70	189	56	29.63
ALL	789	200	25.35

Table 3 – The degree of discordance between the first and the second interpretation of lung perfusion scans according to the scintigraphic probability of pulmonary embolism

SCAN RESULT	No OF PTS	No OF DISCORDANT INTERPRETATIONS	PER CENT DISCORDANCE
NORMAL SCAN	140	3	2.14 %
LOW PROBABILITY OF PE	213	35	16.43 %
INTERMEDIATE PROBABILITY OF PE	181	92	50.83 %
HIGH PROBABILITY OF PE	255	70	27.45 %
ALL	789	200	25.35 %

Results – The disagreement between the first interpretation of lung perfusion scans and the reevaluation by two observers was found in 200 (25,35%) of 789 patients. The probability of lung embolism increased in 185 (92,5%) of patients where the disagreement between the first and the second reading was found. The degree of discordance is low in the patients under 30 years of age (9,59% (Table 2)) and in the interpretation of normal lung perfusion scan (2,14%), (Table 3).

Ventilation lung scanning by ^{99m}Tc -diethylenetriaminopentaacetate aerosol (^{99m}Tc -DTPA), which was performed after perfusion lung scanning in 59 patients, influenced the diagnosis in 14 (23,72%) of patients by lowering the predicted probability of lung embolism as assessed by perfusion scanning alone.

Discussion – Disagreement in repeated readings of diagnostic pictures was first studied in radiology. According to Smith (7), following double reading of 300 consecutive radiologic examinations, diagnostic disagreements occur in about one-third of all film readings and about two-thirds of all errors are found by modified dual-reading techniques.

In our study, lung scintigrams performed in last two years were reevaluated by two experienced nuclear medicine physicians using PLOPED criteria and compared to the results of random approach of several individual, sometimes less experienced, nuclear medicine physicians who were not strictly using special criteria for lung scan interpretation. The percent of discordance between the first and the second reading of the scans in our study was comparable to the results published by Smith. An important difference was found in some of the patients in whom repeated reading increased the probability of lung embolism from low to intermediate or from intermediate to high. In the latter group, the disagreement was partly due to illdefining of larger than segmental defects composed of several juxtaposed segments thus overlaying the anatomic borders. Further, problems have arisen as already noted by Sullivan (6), in the definition of the size of the subsegmental defects. The multiobserver approach is of great value in these cases.

In our pts the ventilation scans mostly decreased the probability of lung embolism estimated on the basis of perfusion scans only. Unfortunately they were not performed frequently enough in all patients.

We conclude that the systematic applying of PLOPED criteria, especially with multiobserver approach, surpasses the nonsystematic scan analysis particularly in the borderline scans. Substantial experience is necessary for interpreters. Ventilation scan is peremptory in the cases of intermediate and high probability of lung embolism estimated on the base of perfusion scan, to avoid the scintigraphic overdiagnosis of high probability of lung embolism.

Povzetek

PONOVNA OCENA SCINTIGRAMOV PLJUČ BOLNIKOV S SUMOM NA PLJUČNO EMBOLIJO

Avtorji poročajo o ponovni oceni scintigramov pljuč pri 789 bolnikih, ki so jih preiskovali v letih 1988 in 1989 zaradi kliničnega suma na pljučno embolijo. Pri vseh bolnikih je bila izvedena perfuzijska scintigrafija pljuč z ^{99m}Tc -albuminskim makroagregatom (^{99m}Tc -MAA). Ventilacijsko scintigrafijo pljuč z aerosolom (^{99m}Tc -diethylenetriaminopentaacetata (^{99m}Tc -DTPA) so izvedli po perfuzijski scintigrafiji 59 bolnikom.

Pri ponovnem ocenjevanju scintigramov avtorji uporabljajo merila priporočena v mednarodni prospektivni študiji za diagnostiko pljučnih embolizmov (PLOPED (5)).

Pri 200 (25,35%) bolnikih od 789 se razlikujejo rezultati prvega in drugega odčitavanja perfuzijskih scintigramov pljuč. Stopnja neskladja je najmanjša pri bolnikih mlajših od 30 let (9,59%) in pri normalnih scintigramih pljuč (2,14%). Z uporabo kriterijev PLOPED se pri 185 (92,5%) bolnikih od 200 poveča verjetnost pljučnih embolizmov.

Pri 14 (23,72%) bolnikih od 59 rezultat dodatne ventilacijske scintigrafije pljuč zmanjša verjetnost obstoja pljučnih embolizmov.

Interpretacija scintigramov pljuč z dvema odčitovalcema, ki uporabljata merila PLOPED, poveča občutljivost (in s tem možnost lažno patoloških rezultatov) perfuzijske scintigrafije v diagnostiki pljučnih embolizmov v primerjavi s tehniko z enim odčitovalcem. Ventilacijska scintigrafija pljuč, ki sledi perfuzijski scintigrafiji, zmanjša število lažno patoloških rezultatov perfuzijske scintigrafije, posebno tistih, ki kažejo srednjo in visoko verjetnost pljučnih embolizmov.

References

- McNeil BJ, Holman BL, Adelstein SJ. The scintigraphic definition of pulmonary embolism. *JAMA* 1974; 227: 753-6.
- Veal HW, Sirota PS, Nelp WB. Ventilation perfusion scanning for pulmonary embolism: refinement of predictive value through Bayesian analysis. *AJR* 1985; 145: 967-72.
- Biello DR, Mattar AG, McKnight RC, Siegel BA. Ventilation perfusion studies in suspected pulmonary embolism. *AJR* 1979; 133: 1033-7.
- Hull RD, Hirsh J, Carter CJ, Raskob GE, GILL GJ, Jay RM, Leclerc JR, David M, Coates G. Diagnostic

value of ventilation perfusion lung scanning in patients with suspected pulmonary embolism. Chest 1985; 88: 819-21.

5. Sostman HD, Rapoport S, Gottschalk A, Green-span RH. Imaging of pulmonary embolism. Invest. Radiol. 1986; 21: 443-53.

6. Sullivan DC, Coleman RE, Mills SR, Ravin CE, Hedlund LW. Lung scan interpretation: effect of diffe-

rent observers and different criteria. Radiology 1983; 149: 803-7.

7. Smith MJ: Errors in diagnostic radiology on the basis of complacency. Amer J Roentgenol Rad Ther Nucl Med 1965; 94: 689-703.

Author's address: Dr Nataša Budihna, Univerzitetna klinika za nuklearno medicino, Zaloška 7, Ljubljana

SANOLABOR

Izvozno uvozno podjetje
za promet z medicinskimi instrumenti, aparati, zdravili,
opremo za bolnice, laboratorije in lekarne

LJUBLJANA, Cigaletova 9
Telefon: (061) 317-355
Telex: 31-668 sanlab yu
Telefax: 325-395

Predstavništvo:
ZAGREB, Šulekova 12
Telefon: (041) 233-369
Telefax: (041) 228-298

40 let z vami!

Iz prodajnega programa trgovine na debelo nudimo široko izbiro domačega in uvoženega blaga po konkurenčnih cenah in sicer:

- rentgenske filme in kemikalije proizvajalcev »Fotokemikà«, »KODAK«, »3 M – TRIMAX«, »AGFA GEVAERT« in drugih,
- medicinske instrumente, specialno medicinsko in sanitetno blago, potrošno blago za enkratno uporabo,
- medicinske aparate in rezervne dele zanje ter bolniško in drugo opremo,
- tekstilne izdelke, konfekcijo in obutev za potrebe zdravstvenih, proizvodnih in varstvenih organizacij,
- laboratorijske aparate, opremo, laboratorijsko steklovino, reagente, kemikalije in pribor,
- aparate, instrumente in potrošno blago za zobozdravstvo

V prodajalni na Cigaletovi 9 v Ljubljani prodajamo izdelke iz asortimana trgovine na debelo, s posebnim poudarkom na blagu za zobozdravstvo, nego bolnikov, ortopedskih pripomočkah in ostalemu blagu za široko potrošnjo.