

Proceedings of 9th Socratic Lectures 2024



University of Ljubljana



Repository

Scanning electron microscope images of *Phaeodactylum tricornutum* culture

Bedina Zavec Apolonija¹, Božič Darja^{2,3}, Hočevar Matej⁴, Igljč Aleš^{3,5}, Jeran Marko^{2,3}, Kralj-Igljč Veronika^{2*}, Romolo Anna²

¹National Institute of Chemistry, ²University of Ljubljana, Faculty of Health Sciences, Laboratory of Clinical Biophysics, ³University of Ljubljana, Faculty of Electrical Engineering, Laboratory of Physics,

⁴Institute of Metals and Technology, ⁵University of Ljubljana, Faculty of Medicine, Laboratory of Clinical Biophysics, Ljubljana, Slovenia

* Corresponding author: Veronika Kralj-Igljč, veronika.kralj-iglic@zf.uni-lj.si

Citation: Bedina Zavec A, Božič D, Hočevar M, Igljč A, Jeran M, Kralj-Igljč V, Romolo A. Scanning electron microscope images of *Phaeodactylum tricornutum* culture. Proceedings of Socratic Lectures. 2024, 9, 192-327.

<https://doi.org/10.55295/PSL.2024.D12>

Publisher's Note: UL ZF stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC0) license

Abstract: Scanning electron microscope images of small cellular particles isolated from conditioned media of microalgae *Phaeodactylum tricornutum* are presented. Each image is supplemented by description of the preparation of the sample and the data on the imaging technique and equipment.

The data curators of the repository are Veronika Kralj-Igljč and Anna Romolo. More data on experiments with microalgae small cellular particles can be found in (Adamo et al., 2021), (Picciotto et al., 2021) and (Božič et al., 2022).

Acknowledgements: This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 801338 and ARRS projects P1-0391, P2-0232, P3-0388, J1-9162, J2-4447, J2-4427, L3-2621, J3-3066, IO-0006 (A) and National Research, Development and Innovation Office (Hungary), grant number SNN 138407.

Keywords: Extracellular vesicles, Extracellular particles, Nanoalgosomes, Exosomes

References

1. Adamo G, Fierli D, Romancino DP, et al. Nanoalgosomes: Introducing extracellular vesicles produced by microalgae. *J Extracell Vesicles*. 2021;10(6):e12081. doi:10.1002/jev2.12081
2. Picciotto S, Barone ME, Fierli D, et al. Isolation of extracellular vesicles from microalgae: towards the production of sustainable and natural nanocarriers of bioactive compounds. *Biomater Sci*. 2021;9(8):2917-2930. doi:10.1039/d0bm01696a
3. Božič D, Hočevar M, Jeran M et al. Ultrastructure and stability of cellular nanoparticles isolated from *Phaeodactylum tricornutum* and *Dunaliella tertiolecta* conditioned media. *Open Res Europe* 2022, 2:121 (<https://doi.org/10.12688/openreseurope.14896.1>)




Figure *Phaeodactylum tricornutum* culture F2 SEM 1.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 μmol/m²s) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO₄ for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).

From: <https://zenodo.org/record/6908895>. Image:6
DOI 10.5281/zenodo.6908895.



Figure *Phaeodactylum tricornutum* culture in F2 SEM 2.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 μmol/m²s) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO₄ for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).

From: <https://zenodo.org/record/6908895.Image:7>

DOI 10.5281/zenodo.6908895.



Figure *Phaeodactylum tricornerutum* culture F2 SEM 3.

Cultivation of the algae

Culture of *Phaeodactylum tricornerutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).

From Božič et al., 2022,
<https://doi.org/10.12688/openreseurope.14896.1>.




Figure *Phaeodactylum tricornutum* culture F2 SEM 4.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 5.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 6.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture F2 SEM 7.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture F2 SEM 8.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornerutum* culture F2 SEM 9.

Cultivation of the algae

Culture of *Phaeodactylum tricornerutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 10.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture in F2 SEM 11.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornerutum* culture in F2 SEM 12.

Cultivation of the algae

Culture of *Phaeodactylum tricornerutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 13.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 14.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 15.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 16.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 17.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornerutum* culture F2 SEM 18.

Cultivation of the algae

Culture of *Phaeodactylum tricornerutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 19.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 20.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture in F2 SEM 21.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)17. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 22.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 23.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 24.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture F2 SEM 25.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).


Figure *Phaeodactylum tricornutum* culture F2 SEM 26.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).






Figure *Phaeodactylum tricornutum* culture in F2 SEM 27.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA)¹⁷. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 28.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture F2 SEM 29.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Guillard's (F/2) Marine Water Enrichment Solution (ref. G0154, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 30.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricorneratum* culture BG11 SEM 31.

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture BG11 SEM 32.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 33.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 34.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 35.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 36.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 37.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).

Figure *Phaeodactylum tricornutum* culture BG11 SEM 38.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).






Figure *Phaeodactylum tricornutum* culture BG11 SEM 39.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornerutum* culture BG11 SEM 40.

Cultivation of the algae

Culture of *Phaeodactylum tricornerutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 41.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricorneratum* culture BG11 SEM 42.

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricorneratum* culture BG11 SEM 43.

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 44.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG1 SEM 45.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG1 SEM 46.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture BG11 SEM 47.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with BG11 Broth (ref. 73816, Sigma Aldrich, Switzerland). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornerutum* culture LB SEM 48.

Cultivation of the algae

Culture of *Phaeodactylum tricornerutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture LB SEM 49.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with Lennox LB broth (ref. L3022, Sigma Aldrich, USA) broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture LB SEM 50.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM LB 51.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA) medium. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM LB 52.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM LB 53.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA) medium. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornerutum* culture SEM LB 54.

Cultivation of the algae

Culture of *Phaeodactylum tricornerutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM LB 55.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA) medium. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).

Figure *Phaeodactylum tricornutum* culture SEM LB 56.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM LB 57.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA) medium. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM LB 58.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM LB 59.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA) medium. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM LB 60.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM LB 61.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA) medium. Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM LB 62.

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricorneratum* culture SEM 63.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornerutum* culture SEM 64.
SEM feodaktikum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornerutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 65.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021 -
10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 66.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021 -
10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 67.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021 -
10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricorneratum* culture SEM 68.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 69.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021 -
10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 70.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 71.
SEM feodaktikum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornerum* culture SEM 72.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornerum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 73.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 74.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 75.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 76.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 77.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 78.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021 - 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 79.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 80.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 81.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 82.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 83.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 84.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 85.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 86.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 87.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 88.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).


Figure *Phaeodactylum tricornutum* culture SEM 89.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).






Figure *Phaeodactylum tricorneratum* culture SEM 90.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 91.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 92.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 93.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 94.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 95.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 96.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 97.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricorneratum* culture SEM 98.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 99.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 100.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 101.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021- 10 veliko krogel dolgo na plosci

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown in a respirometer (Echo, Slovenia) in 0.5-L borosilicate bottles, at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 102.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 103.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 104.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 μmol/m²s) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO₄ for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 105.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 106.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 107.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 108.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 109.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 110.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojiska

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 111.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricorneratum* culture SEM 112.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle, with aeration of 0.2 L/min.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 113.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 114.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 115.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 116.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 117.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojiska

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 118.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 119.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



IMT SEI 15.0kV X15,000 1μm WD 10.0mm

Figure *Phaeodactylum tricorneratum* culture SEM 120.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 μmol/m²s) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO₄ for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 121.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojiska

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 122.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 123.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 124.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojiska

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 125.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 126.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 127.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 128.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 129.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).

Figure *Phaeodactylum tricornutum* culture SEM 130.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).



Figure *Phaeodactylum tricornutum* culture SEM 131.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021-11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 132.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojjsca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 133.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricornutum* culture SEM 134.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojisca

Cultivation of the algae

Culture of *Phaeodactylum tricornutum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).




Figure *Phaeodactylum tricorneratum* culture SEM 135.
SEM feodaktilum netretiran #2ovoid/Darja-21-10-2021
- 11 ovalne celice s trdnega gojiska

Cultivation of the algae

Culture of *Phaeodactylum tricorneratum* CCAP 1052/1A from the Culture Collection of Algae and Protozoa (CCAP) of SAMS (Oban, Scotland) was grown in artificial seawater (Reef Crystals, Aquarium Systems, France). 22 g of salt was dissolved in one litre of distilled water, sterile filtered (0.2-micron cellulose filters, ref. 11107-47-CAN, Sartorius Stedim Biotech GmbH, Germany), autoclaved, and supplemented with LB Lennox LB broth (ref. L3022, Sigma Aldrich, USA). Culture was grown on agar at 20 °C and 20 % illumination (approximately 250 $\mu\text{mol}/\text{m}^2\text{s}$) with a 14-hour light / 10-hour dark cycle.

Scanning Electron Microscopy (SEM)

Samples were loaded onto 0.05-micron MCE filters (MF-Millipore™, ref. VMWP01300) and incubated in 2% OsO_4 for two hours. Then the osmium was removed, and the filter was taken out from the holder and further treated in a 24-well plate by changing the bath solution. After washing three times in distilled water, the samples were dehydrated in a graded series of ethanol (30%, 50%, 70%, 80%, 90%, absolute), treated with hexamethyldisilazane (30%, 50% mixtures with absolute ethanol, followed by pure hexamethyldisilazane), and air-dried. Samples were sputtered with Au/Pd (PECS Gatan 682) and examined with a JSM-6500F Field Emission Scanning Electron Microscope (JEOL Ltd., Tokyo, Japan).