



The truncated moment problem on curves $y = q(x)$ and $yx^\ell = 1$

A. Zalar

To cite this article: A. Zalar (2024) The truncated moment problem on curves $y = q(x)$ and $yx^\ell = 1$, Linear and Multilinear Algebra, 72:12, 1922-1966, DOI: [10.1080/03081087.2023.2212316](https://doi.org/10.1080/03081087.2023.2212316)

To link to this article: <https://doi.org/10.1080/03081087.2023.2212316>



© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 16 May 2023.



Submit your article to this journal [↗](#)



Article views: 698



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

The truncated moment problem on curves $y = q(x)$ and $yx^\ell = 1$

A. Zalar ^{a,b,c}

^aFaculty of Computer and Information Science, University of Ljubljana, Ljubljana, Slovenia; ^bFaculty of Mathematics and Physics, University of Ljubljana, Ljubljana, Slovenia; ^cInstitute of Mathematics, Physics and Mechanics, Ljubljana, Slovenia

ABSTRACT

In this paper, we study the bivariate truncated moment problem (TMP) on curves of the form $y = q(x)$, $q(x) \in \mathbb{R}[x]$, $\deg q \geq 3$ and $yx^\ell = 1$, $\ell \in \mathbb{N} \setminus \{1\}$. For even degree sequences, the solution based on the size of moment matrix extensions was first given by Fialkow [Fialkow L. Solution of the truncated moment problem with variety $y = x^3$. *Trans Amer Math Soc.* 2011;363:3133–3165.] using the truncated Riesz–Haviland theorem [Curto R, Fialkow L. An analogue of the Riesz–Haviland theorem for the truncated moment problem. *J Funct Anal.* 2008;255:2709–2731.] and a sum-of-squares representations for polynomials, strictly positive on such curves [Fialkow L. Solution of the truncated moment problem with variety $y = x^3$. *Trans Amer Math Soc.* 2011;363:3133–3165.; Stochel J. Solving the truncated moment problem solves the moment problem. *Glasgow J Math.* 2001;43:335–341.]. Namely, the upper bound on this size is quadratic in the degrees of the sequence and the polynomial determining a curve. We use a reduction to the univariate setting technique, introduced in [Zalar A. The truncated Hamburger moment problem with gaps in the index set. *Integral Equ Oper Theory.* 2021;93:36. doi: [10.1007/s00020-021-02628-6](https://doi.org/10.1007/s00020-021-02628-6); Zalar A. The truncated moment problem on the union of parallel lines. *Linear Algebra Appl.* 2022;649:186–239. doi: [10.1016/j.laa.2022.05.008](https://doi.org/10.1016/j.laa.2022.05.008); Zalar A. The strong truncated Hamburger moment problem with and without gaps. *J Math Anal Appl.* 2022;516:126563. doi: [10.1016/j.jmaa.2022.126563](https://doi.org/10.1016/j.jmaa.2022.126563).], and improve Fialkow’s bound to $\deg q - 1$ (resp. $\ell + 1$) for curves $y = q(x)$ (resp. $yx^\ell = 1$). This in turn gives analogous improvements of the degrees in the sum-of-squares representations referred to above. Moreover, we get the upper bounds on the number of atoms in the minimal representing measure, which are $k \deg q$ (resp. $k(\ell + 1)$) for curves $y = q(x)$ (resp. $yx^\ell = 1$) for even degree sequences, while for odd ones they are $k \deg q - \lceil \frac{\deg q}{2} \rceil$ (resp. $k(\ell + 1) - \lfloor \frac{\ell}{2} \rfloor + 1$) for curves $y = q(x)$ (resp. $yx^\ell = 1$). In the even case,

ARTICLE HISTORY

Received 20 December 2022
Accepted 18 April 2023

COMMUNICATED BY

L. Molnar

KEYWORDS

Truncated moment problems; K -moment problems; K -representing measure; minimal measure; moment matrix extensions; positivstellensatz; linear matrix inequality

AMS CLASSIFICATIONS

Primary 44A60; 47A57; 47A20; Secondary 15A04; 47N40

CONTACT A. Zalar  aljaz.zalar@fri.uni-lj.si.  Faculty of Computer and Information Science, University of Ljubljana, Večna pot 113, Ljubljana, Slovenia; Faculty of Mathematics and Physics, University of Ljubljana, Jadranska ulica, 19, Ljubljana, Slovenia; Institute of Mathematics, Physics and Mechanics, Jadranska ulica, 19, Ljubljana, Slovenia

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

these are counterparts to the result by Riener and Schweighofer [Riener C, Schweighofer M. Optimization approaches to quadrature: a new characterization of Gaussian quadrature on the line and quadrature with few nodes on plane algebraic curves, on the plane and in higher dimensions. J Complex. 2018;45:22–54., Corollary 7.8], which gives the same bound for odd degree sequences on all plane curves. In the odd case, their bound is slightly improved on the curves we study. Further on, we give another solution to the TMP on the curves studied based on the feasibility of a linear matrix inequality, corresponding to the univariate sequence obtained, and finally we solve concretely odd degree cases to the TMP on curves $y = x^\ell$, $\ell = 2, 3$, and add a new solvability condition to the even degree case on the curve $y = x^2$.

1. Introduction

Given a real two-dimensional sequence

$$\beta^{(d)} = \{\beta_{0,0}, \beta_{1,0}, \beta_{0,1}, \dots, \beta_{d,0}, \beta_{d-1,1}, \dots, \beta_{1,d-1}, \beta_{0,d}\}$$

of degree d and a closed subset K of \mathbb{R}^2 , the *truncated moment problem (K-TMP)* supported on K for $\beta^{(d)}$ asks to characterize the existence of a positive Borel measure μ on \mathbb{R}^2 with support in K , such that

$$\beta_{i,j} = \int_K x^i y^j d\mu \text{ for } i, j \in \mathbb{Z}_+, 0 \leq i + j \leq d. \tag{1}$$

If such a measure exists, we say that $\beta^{(d)}$ has a representing measure supported on K and μ is its *K-representing measure (K-rm)*.

Let $k = \lceil \frac{d}{2} \rceil$. In the degree-lexicographic order $1, X, Y, X^2, XY, Y^2, \dots, X^k, X^{k-1}Y, \dots, Y^k$ of rows and columns, the corresponding moment matrix to β is equal to

$$M_k = M_k(\beta) := \begin{pmatrix} M[0,0](\beta) & M[0,1](\beta) & \cdots & M[0,k](\beta) \\ M[1,0](\beta) & M[1,1](\beta) & \cdots & M[1,k](\beta) \\ \vdots & \vdots & \ddots & \vdots \\ M[k,0](\beta) & M[k,1](\beta) & \cdots & M[k,k](\beta) \end{pmatrix}, \tag{2}$$

where

$$M[i,j](\beta) := \begin{pmatrix} \beta_{i+j,0} & \beta_{i+j-1,1} & \beta_{i+j-2,2} & \cdots & \beta_{ij} \\ \beta_{i+j-1,1} & \beta_{i+j-2,2} & \beta_{i+j-3,3} & \cdots & \beta_{i-1,j+1} \\ \beta_{i+j-2,2} & \beta_{i+j-3,3} & \beta_{i+j-4,4} & \cdots & \beta_{i-2,j+2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \beta_{j,i} & \beta_{j-1,i+1} & \beta_{j-2,i+2} & \cdots & \beta_{0,i+j} \end{pmatrix}$$

and for odd d , the lower right corner $M[k,k]$ of $M_k(\beta)$ is undefined. Until the end of this section, we assume that M_k is fully determined, i.e. it corresponds to the even degree sequence $\beta^{(2k)}$. Let $\mathbb{R}[x, y]_k := \{p \in \mathbb{R}[x, y] : \deg p \leq k\}$ stand for the set of real polynomials in variables x, y of degree at most k , where for $p \neq 0$ the degree $\deg p$ stands for the

maximal sum $i + j$ over all monomials $x^i y^j$ appearing in p with a nonzero coefficient a_{ij} , while for $p \equiv 0$, $\deg p = 0$. For every $p(x, y) = \sum_{i,j} a_{ij} x^i y^j \in \mathbb{R}[x, y]_k$, we define its *evaluation* $p(X, Y)$ on the columns of the matrix M_k by replacing each capitalized monomial $X^i Y^j$ in $p(X, Y) = \sum_{i,j} a_{ij} X^i Y^j$ by the column of M_k , indexed by this monomial. Then $p(X, Y)$ is a vector from the linear span of the columns of M_k . If this vector is the zero one, i.e. all coordinates are equal to 0, then we say p is a *column relation* of M_k . Recall from [1], that β has an rm μ with the support $\text{supp}(\mu)$ being a subset of $\mathcal{Z}(p) := \{(x, y) \in \mathbb{R}^2 : p(x, y) = 0\}$ if and only if p is a column relation of M_k . We say that the matrix M_k is *recursively generated* (rg) if for $p, q, pq \in \mathbb{R}[x, y]_k$ such that p is a column relation of M_k , it follows that pq is also a column relation of M_k .

A *concrete solution* to the TMP is a set of necessary and sufficient conditions for the existence of a K -rm that can be tested in numerical examples. Among necessary conditions, M_k must be positive semidefinite (psd) and rg [1,2]. A crucial tool to tackle the TMP, discovered by Curto and Fialkow in 1996, was a *flat extension theorem (FET)* [1, Theorem 7.10] (see also [3, Theorem 2.19] and [4] for an alternative proof), which states that $\beta^{(2k)}$ admits a $(\text{rank } M_k)$ -atomic rm if and only if M_k is psd and admits a rank-preserving extension to a moment matrix M_{k+1} . Using the FET as the main tool the bivariate TMP has been concretely solved in the following cases: K is the variety defined by a polynomial $p(x, y) = 0$ with $\deg p \leq 2$ [5–8]; $K = \mathbb{R}^2$, $k = 2$ and M_2 is invertible [9], first solved nonconstructively in [10]; K is the variety $y = x^3$ [11]; M_k has a special feature such as *recursive determinateness* [12] or *extremality* [13]. Some special cases have also been solved in [14–16] based on the FET and in [17–25] using different approaches.

References to some classical work on the TMP are monographs [26–28], while for a recent development in the area we refer a reader to [29]. We also mention some variants of the TMPs, which attracted a recent research interest, such as versions of the infinite dimensional TMPs [30–32], the TMP on subspaces of polynomial algebra [33], the TMP for commutative \mathbb{R} -algebras [34], matrix and operator TMPs [35–43], etc.

In our previous work, we introduced a new approach to tackle the singular bivariate TMP, namely *a reduction to the univariate setting technique*. The idea is to use one of the column relations to transform the problem into the equivalent univariate TMP, where also negative moments of the measure could be present or not all moments between the lowest and highest degree ones are known. In the case all moments from degree 0 to the highest degree are known, the situation is well understood in terms of the existence and uniqueness of the rm and has been solved in full generality [44,45] for measures with support \mathbb{R} , $[a, \infty)$ or $[a, b] \subset \mathbb{R}$, $a, b \in \mathbb{R}$, $a < b$, as well as for even and odd degree sequences. In the presence of negative moments, we gave a solution along the lines of the classical case in [46], where we note that the existence of the solution even in the matrix case was already established by Simonov [47] (but the measure is not constructively obtained and the number of atoms in a minimal measure does not directly follow from this more general approach). Using these results, we presented [38,46,48,49] alternative solutions with shorter proofs compared to the original ones to the TMPs on the curves $xy = 0$, $y = x^3$, $y^2 = y$, $xy = 1$, but also obtained solutions to new cases, namely on the curve $y^2 = x^3$, on the union of three parallel lines and on $xy^2 = 1$.

The motivation for this paper was to use a reduction technique to the TMP on curves of the form $y = q(x)$ and $yq(x) = 1$, where $q \in \mathbb{R}[x]$. In [11, Section 6], Fialkow gave a

solution to the TMP on these curves for even degree sequences in terms of the bound on the degree m for which the existence of a positive extension M_m of M_k is equivalent to the existence of a rm. Namely, his bound is quadratic in k and $\deg q$. Using a reduction technique we are able to decrease his bound in the even degree case for all curves of the form $y = q(x)$, $\deg q \geq 3$, to $\deg q - 1$ and for curves of the form $yx^\ell = 1$, $\ell \in \mathbb{N} \setminus \{1\}$, to $\ell + 1$, which is our first main result. Moreover, the reduction technique also works in the odd degree case. A corollary to this improved bounds are also improvements of the sum-of-squares representations for polynomials, strictly positive on such curves, by decreasing the degrees of the polynomials in the representation. Our second main result are the upper bounds on the number of atoms in the minimal rm, which are for curves $y = q(x)$, $\deg q \geq 3$, equal to $k \deg q$ in the even and $k \deg q - \lceil \frac{\deg q}{2} \rceil$ in the odd case and for curves $yx^\ell = 1$, $\ell \geq 2$, equal to $k(\ell + 1)$ in the even and $k(\ell + 1) - \lfloor \frac{\ell}{2} \rfloor + 1$ in the odd case. In the even case these results are counterparts to the result of Riemer and Schweighofer [50, Corollary 7.8], who proved that for *all* plane curves, odd degree sequence has at most $k \deg q$ atoms in the minimal measure. For curves of the above form, we improve their bound slightly in the odd degree case. The third main result of the paper is another solution to the TMPs studied, which is based on the feasibility of a linear matrix inequality corresponding to the univariate sequence obtained. Moreover, we give concrete solutions to the odd degree TMPs on the curves $y = x^2$ and $y = x^3$ and an alternative solution to the even degree case on $y = x^2$ with a new solvability condition, which will be crucially needed in the solution of the TMP on the reducible curve $y(y - x^2) = 0$ in our forthcoming work.

1.1. Reader's guide

This paper is organized as follows. In Section 2, we fix some further notation and known results on the TMP, which will be used in the proofs of our results. In Section 3, we give two solutions to the K -TMP for $K = \{(x, y) \in \mathbb{R}^2 : y = q(x)\}$, $q \in \mathbb{R}[x]$, $\deg q \geq 3$, one based on the size of psd extensions of the moment matrix needed (see Theorems 3.1 and 3.2 for the even and odd degree cases, respectively) and the other one based on the feasibility question of a certain linear matrix inequality (see Theorem 3.8). Theorems 3.1 and 3.2 also give bounds on the number of atoms in a minimal K -rm. Moreover, Theorem 3.1 gives a Positivstellensatz on K as a corollary (see Corollary 3.3). Further on, we solve concretely the TMPs on the curve $y = x^2$ (see Theorems 3.4 and 3.7 for the even and odd degree cases, respectively) and on $y = x^3$ for the odd case (see Theorem 3.13). In Section 4, we give the corresponding results to the ones from Section 3 for curves $yx^\ell = 1$, $\ell \in \mathbb{N} \setminus \{1\}$. Theorems 4.1 and 4.2 are the counterparts of Theorems 3.1 and 3.2, respectively, Corollary 4.3 of Corollary 3.3 and Theorem 4.4 of Theorem 3.8.

2. Preliminaries

In this section, we fix some terminology, notation and present some tools needed in the proofs of our main results.

We write $\mathbb{R}^{n \times m}$ for the set of $n \times m$ real matrices. For a matrix M , we call the linear span of its columns a *column space* and denote it by $\mathcal{C}(M)$. The set of real symmetric matrices of size n will be denoted by S_n . For a matrix $A \in S_n$, the notation $A > 0$ (resp. $A \geq 0$) means A is positive definite (pd) (resp. positive semidefinite (psd)).

In the rest of this section, let $d \in \mathbb{N}$ and $\beta = \beta^{(d)} = \{\beta_{ij}\}_{i,j \in \mathbb{Z}_+, 0 \leq i+j \leq d}$ be a bivariate sequence of degree d .

2.1. Moment matrix

Let $k = \lceil \frac{d}{2} \rceil$ and $M_k = M_k(\beta)$ be the moment matrix of β (see (2)). Let Q_1, Q_2 be subsets of the set $\{X^i Y^j : i, j \in \mathbb{Z}_+, 0 \leq i + j \leq k\}$. We denote by $(M_k)|_{Q_1, Q_2}$ the submatrix of M_k consisting of the rows indexed by the elements of Q_1 and the columns indexed by the elements of Q_2 . In case $Q := Q_1 = Q_2$, we write $(M_k)|_Q = (M_k)|_{Q,Q}$ for short.

Remark 2.1: Whenever Q_1, Q_2 will be subsets of $\{x^i y^j : i, j \in \mathbb{Z}_+, 0 \leq i + j \leq k\}$ in the rest of the paper, in the notation $(M_k)|_{Q_1, Q_2}$ all monomials from Q_1, Q_2 are meant capitalized, i.e. $x^i y^j \mapsto X^i Y^j$.

2.2. Atomic measures

For $x \in \mathbb{R}^m$, δ_x stands for the Dirac measure supported on x . By a *finitely atomic positive measure* on \mathbb{R}^m , we mean a measure of the form $\mu = \sum_{j=0}^\ell \rho_j \delta_{x_j}$, where $\ell \in \mathbb{N}$, each $\rho_j > 0$ and each $x_j \in \mathbb{R}^m$. The points x_j are called *atoms* of the measure μ and the constants ρ_j the corresponding *densities*.

2.3. Riesz functional

The functional $L_\beta : \mathbb{R}[x, y]_{\leq d} \rightarrow \mathbb{R}$, defined by

$$L_\beta(p) := \sum_{\substack{i,j \in \mathbb{Z}_+, \\ 0 \leq i+j \leq d}} a_{i,j} \beta_{i,j}, \quad \text{where } p = \sum_{\substack{i,j \in \mathbb{Z}_+, \\ 0 \leq i+j \leq d}} a_{i,j} x^i y^j,$$

is called the *Riesz functional of the sequence* β .

2.4. Affine linear transformations

Let $K \subseteq \mathbb{R}^2$. The existence of a K -rm for β is invariant under invertible affine linear transformations (alts) of the form

$$\phi(x, y) = (\phi_1(x, y), \phi_2(x, y)) := (a + bx + cy, d + ex + fy), \quad (x, y) \in \mathbb{R}^2, \quad (3)$$

$a, b, c, d, e, f \in \mathbb{R}$ with $bf - ce \neq 0$ in the sense which we now explain. We denote by $\tilde{\beta}$ the two-dimensional sequence defined by

$$\tilde{\beta}_{i,j} = L_\beta (\phi_1(x, y)^i \cdot \phi_2(x, y)^j),$$

where L_β is the Riesz functional of β .

Proposition 2.1 ([7, Proposition 1.9]): Assume the notation above and let $d = 2k$.

- (1) $M_k(\beta)$ is psd if and only if $M_k(\tilde{\beta})$ is psd.

- (2) $\text{rank } M_k(\beta) = \text{rank } M_k(\tilde{\beta})$.
- (3) $M_k(\beta)$ is rg if and only if $M_k(\tilde{\beta})$ is rg.
- (4) β admits a r -atomic K -rm if and only if $\tilde{\beta}$ admits an r -atomic $\phi(K)$ -rm.

In case $d = 2k - 1$ is odd, the block $M[k, k]$ of $M_k(\beta)$ is undefined. We say that $M_k(\beta)$ is psd completable if there exists an extension $\beta^{(2k)}$ of β such that $M_k(\beta^{(2k)})$ is psd.

Proposition 2.2: Assume the notation above and let $d = 2k - 1, k \in \mathbb{N}$.

- (1) $M_k(\beta)$ is psd completable if and only if $M_k(\tilde{\beta})$ is psd completable.
- (2) Let $r \in \mathbb{N}$. There exists an extension $\beta^{(2k)}$ of β such that $\text{rank } M_k(\beta^{(2k)}) = r$ if and only if there exists an extension $\tilde{\beta}^{(2k)}$ of $\tilde{\beta}$ such that $\text{rank } M_k(\tilde{\beta}^{(2k)}) = r$.
- (3) Let $r \in \mathbb{N}$. There exists an extension $\beta^{(2k)}$ of β such that $M_k(\beta^{(2k)})$ is rg if and only if there exists an extension $\tilde{\beta}^{(2k)}$ of $\tilde{\beta}$ such that $M_k(\tilde{\beta}^{(2k)})$ is rg.
- (4) β admits a r -atomic K -rm if and only if $\tilde{\beta}$ admits an r -atomic $\phi(K)$ -rm.

Proof: Proposition 2.2 follows easily from Proposition 2.1 by defining the extension $\tilde{\beta}^{(2k)}$ of $\tilde{\beta}$ from the extension $\beta^{(2k)}$ of β using the same transformation ϕ together with the Riesz functional $L_{\beta^{(2k)}}$ of the extension. Similarly, for the other direction one uses ϕ^{-1} together with the Riesz functional $L_{\tilde{\beta}^{(2k)}}$ of the extension. For (4) we note that any r -atomic K -rm of the sequence β generates the extension $\beta^{(2k)}$ and then use (4) of Proposition 2.1. ■

2.5. Hankel matrices and univariate sequences

Let $k \in \mathbb{N}$. For $v = (v_0, \dots, v_{2k}) \in \mathbb{R}^{2k+1}$, we define the corresponding Hankel matrix as

$$A_v := (v_{i+j})_{i,j=0}^k = \begin{pmatrix} v_0 & v_1 & v_2 & \cdots & v_k \\ v_1 & v_2 & \ddots & \ddots & v_{k+1} \\ v_2 & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & v_{2k-1} \\ v_k & v_{k+1} & \cdots & v_{2k-1} & v_{2k} \end{pmatrix} \in S_{k+1}. \tag{4}$$

Let $\mathbf{v}_j := (v_{j+\ell})_{\ell=0}^k$ be the $(j + 1)$ -th column of $A_v, 0 \leq j \leq k$. In this notation, we have that

$$A_v = (\mathbf{v}_0 \quad \cdots \quad \mathbf{v}_k).$$

As in [44], the rank of v , denoted by $\text{rank } v$, is defined by

$$\text{rank } v = \begin{cases} k + 1, & \text{if } A_v \text{ is nonsingular,} \\ \min \{i: \mathbf{v}_i \in \text{span}\{\mathbf{v}_0, \dots, \mathbf{v}_{i-1}\}\}, & \text{if } A_v \text{ is singular.} \end{cases}$$

We denote

- the upper left-hand corner $(v_{i+j})_{i,j=0}^m \in S_{m+1}$ of A_v of size $m + 1$ by $A_v(m)$
- the lower right-hand corner $(v_{i+j})_{i,j=k-m}^k \in S_{m+1}$ of A_v of size $m + 1$ by $A_v[m]$.

We write

$$v^{(\text{rev})} := (v_{2k}, v_{2k-1}, \dots, v_0)$$

for the *reversed sequence* of v .

A sequence $v = (v_0, \dots, v_{2k})$ is called *positively recursively generated (prg)* if, denoting $r = \text{rank } v$, it holds that $A_v(r - 1) \succ 0$ and in case $r < k + 1$, also

$$v_j = \sum_{i=0}^{r-1} \varphi_i v_{j-r+i} \text{ for } j = r, \dots, 2k,$$

where

$$(\varphi_0 \ \dots \ \varphi_{r-1}) := A_v(r - 1)^{-1} (v_r \ \dots \ v_{2r-1})^T. \tag{5}$$

A sequence $v = (v_0, \dots, v_{2k})$ is called *negatively recursively generated (nrg)* if, denoting $r = \text{rank } v^{(\text{rev})}$, it holds that $A_v[r - 1] \succ 0$ and in case $r < k + 1$, also

$$v_{2k-r-j} = \sum_{i=0}^{r-1} \psi_i v_{2k-r+1-j+i} \text{ for } j = 0, \dots, 2k - r,$$

where

$$(\psi_0 \ \dots \ \psi_{r-1}) := A_v[r - 1]^{-1} (v_{2k-2r+1} \ \dots \ v_{2k-r})^T.$$

2.6. Univariate truncated moment problem

Given a real sequence

$$\gamma^{(k_1, k_2)} = (\gamma_{k_1}, \gamma_{k_1+1}, \dots, \gamma_{k_2-1}, \gamma_{k_2})$$

of degree (k_1, k_2) , $k_1, k_2 \in \mathbb{Z}$, $k_1 \leq k_2$, a subset K of \mathbb{R} , the *truncated moment problem supported on K for $\gamma^{(k_1, k_2)}$ ((K, k_1, k_2) -TMP)* asks to characterize the existence of a positive Borel measure μ on \mathbb{R} with support in K , such that

$$\gamma_i = \int_K x^i d\mu \text{ for } i \in \mathbb{Z}, k_1 \leq i \leq k_2. \tag{6}$$

If such a measure exists, we say that $\gamma^{(k_1, k_2)}$ has a *rm* supported on K and μ is its K -*rm*.

The $(\mathbb{R}, 0, k)$ -TMP with $k \in \mathbb{Z}_+$ is the usual *truncated Hamburger moment problem (THMP) of degree k* , which was solved in full generality in [44]. Algorithm 2.1 is a numerical procedure to determine the existence and extract an *rm*. It is an adaptation of [51, Algorithm 4.2], which is an algorithm to extract an *rm* in the multivariate TMP under the assumption that an *rm* exists, to the univariate setting with the addition of deciding whether a *rm* exists. For the latter, the solution from [44] is used.

Algorithm 2.1 Solution to the $(\mathbb{R}, 0, 2k)$ -TMP

Input: A univariate sequence $\gamma = (\gamma_0, \gamma_1, \dots, \gamma_{2k})$.

Output: A negative answer to the existence of an rm for γ or

the points $x_1, \dots, x_r \in \mathbb{R}$ and densities ρ_1, \dots, ρ_r such that $\mu = \sum_{j=1}^r \rho_j \delta_{x_j}$ is an \mathbb{R} -rm for γ .

Algorithm:

1. Try to compute Cholesky factorization VV^T of A_γ , where $V \in \mathbb{R}^{(k+1) \times r}$ is a lower triangular matrix. In case of a failure, an rm does not exist.

2.1 If $r < k + 1$, reduce V to a column echelon form U using Gaussian elimination on columns.

2.2 Else $r = k + 1$. Choose $\gamma_{2k+1} \in \mathbb{R}$ arbitrarily and solve a linear system $Vw = (\gamma_{k+1} \dots \gamma_{2k+1})^T$. Reduce $\tilde{V} = (V^T \ w)^T$ to a column echelon form U using Gaussian elimination on columns.

3.1. If $r < k + 1$ and U is of the form $U = (I_r \ B)^T$ for some $B \in \mathbb{R}^{r \times (k+1-r)}$ or $r = k + 1$, define N as an $\mathbb{R}^{r \times r}$ matrix consisting of rows $2, \dots, r + 1$ of U .

3.2. If $r < k + 1$ and U is not as in 3.1, then an rm for γ does not exist.

4. Compute the Schur decomposition $N = QTQ^T$ of N . The diagonal elements x_1, \dots, x_r of T are atoms in an rm for γ .

5. Solve the system $W\rho = (\gamma_0 \dots \gamma_{r-1})^T$, where W is the $r \times r$ Vandermonde matrix with i th row equal to $(x_j^{i-1})_{j=1, \dots, r}$. The coordinates $\rho_j, j = 1, \dots, r$, of ρ are the densities of an rm measure for γ .

Remark 2.2: (1) *Correctness of Algorithm 2.1:* Step 1 checks whether A_γ is psd. Step 2.1 only changes a basis of the column space of V . If Step 2.2 applies, then one can choose γ_{2k+1} arbitrarily and then compute γ_{2k+2} such that for the extended sequence $\tilde{\gamma} = (\gamma, \gamma_{2k+1}, \gamma_{2k+2})$ Step 2.1 applies, i.e. $A_{\tilde{\gamma}}$ is psd, $\text{rank } A_{\tilde{\gamma}} = k + 1 < k + 2$ and its Cholesky decomposition is equal to $\tilde{V}(\tilde{V})^T$. In the situation of Step 3.1, $A_\gamma(r)$ is pd and hence γ is prg. If Step 3.2 applies, the latter is not true. By [44, Theorem 3.9], an rm exists iff γ is prg. Since $U(v_{x_i}^{(0,r-1)}) = v_{x_i}^{(0,r-1)}$ for every atom x_i in an rm for γ , where $v_x^{(0,p)} = (1 \ x \dots \ x^p)^T$, it follows that $N(v_{x_i}^{(0,r-1)}) = x_i v_{x_i}^{(0,r-1)}$ and the atoms of an rm is the eigenvalues of N . These are computed in Step 4 and finally, a Vandermonde system in Step 5 determines the densities.

(2) *Uniqueness of an rm:* If $r < k + 1$ and Step 3.1 applies, then the rm for γ is unique. Otherwise, in case $r = k + 1$, there are infinitely many choices of $(k + 1)$ -atomic rms. Each choice of γ_{2k+1} gives a different one.

(3) *An adaptation of Algorithm 2.1 to solving $(\mathbb{R}, 0, 2k + 1)$ -TMP:* One has to do the following modifications:

(a) Let $\tilde{\gamma} = (\gamma_0, \gamma_1, \dots, \gamma_{2k+1})$ and γ as in Algorithm 2.1.

(b) If Step 3.1 applies with $r < k + 1$, one has to check whether the system $Vw = (\gamma_{k+1} \dots \gamma_{2k+1})^T$ is solvable. If not, an rm does not exist. Otherwise it does and compute it as in Algorithm 2.1.

(c) If Step 3.1 applies with $r = k + 1$, one does not choose γ_{2k+1} , since it is already given. Hence, the minimal $(k + 1)$ -atomic rm is unique.

The $(\mathbb{R} \setminus \{0\}, k_1, k_2)$ -TMP with $k_1, k_2 \in \mathbb{Z}, k_1 < 0 < k_2$ is the *strong truncated Hamburger moment problem (STHMP) of degree (k_1, k_2)* . For even k_1 and k_2 , the solution is [46, Theorem 3.1], but the technique in the proof can be extended to establish also the cases, where k_1, k_2 are not both even.

Let $\mathbb{R}[x^{-1}, x] = \{\sum_{i=r_1}^{r_2} a_i x^i : a_i \in \mathbb{R}, r_1, r_2 \in \mathbb{Z}, r_1 \leq r_2\}$ be the set of Laurent polynomials. For $k_1, k_2 \in \mathbb{Z}, k_1 \leq k_2$, we denote by $V_{(k_1, k_2)}$ a vector subspace in $\mathbb{R}[x^{-1}, x]$ generated by the set $\{x^{k_1}, x^{k_1+1}, \dots, x^{k_2}\}$. For a sequence $\gamma := \gamma^{(k_1, k_2)}$ the functional $L_\gamma : V_{(k_1, k_2)} \rightarrow \mathbb{R}$, defined by

$$L_\gamma(p) := \sum_{k_1 \leq i \leq k_2} a_i \gamma_i, \quad \text{where } p = \sum_{k_1 \leq i \leq k_2} a_i x^i,$$

is called the *Riesz functional of the sequence γ* .

Remark 2.3 (An adaptation of Algorithm 2.1 to solving $(\mathbb{R}, -2k_1, 2k_2)$ -TMP, $k_1, k_2 \in \mathbb{N}$): One has to do the following modifications (see [46, Theorem 3.1]):

- (1) The input is a sequence $\gamma = (-\gamma_{2k_1}, \gamma_{-2k_1+1}, \dots, 2k_2)$. The output is a negative answer to the existence of an rm for γ or the points $x_1, \dots, x_r \in \mathbb{R}$ and densities ρ_1, \dots, ρ_r such that $\mu = \sum_{j=1}^r \rho_j \delta_{x_j}$ is an \mathbb{R} -rm for γ .
- (2) One forms a sequence $\tilde{\gamma} = (\tilde{\gamma}_0, \dots, \tilde{\gamma}_{2(k_1+k_2)})$, where $\tilde{\gamma}_i = \gamma_{i-2k_1}, i = 0, 1, \dots, 2(k_1 + k_2)$, and do all computations from Algorithm 2.1 on $\tilde{\gamma}$.
- (3) If Step 2.2 applies, one chooses $\gamma_{2(k_1+k_2)+1}$ arbitrarily except for the number $v^T C^{-1} v$ in case C is invertible, where C is a submatrix of $A_{\tilde{\gamma}}$ consisting of rows $1, \dots, k_1 + k_2$ and columns $2, \dots, k_1 + k_2 + 1$, and $v = (\gamma_{-k_1+k_2+1} \ \dots \ \gamma_{-2k_2})^T$ is a vector.
- (4) If Step 3.1 applies with $r < k_1 + k_2 + 1$, one has to check whether $(r + 1)$ th column of $A_{\tilde{\gamma}}$ is in the span of columns $2, \dots, r$. If yes, then an rm does not exist. Otherwise it does. Equivalently, one can compute the Cholesky decomposition $V_1 V_1^T$ of the restriction of $A_{\tilde{\gamma}}$ to the principal submatrix on rows and columns $2, 3, \dots, r + 1$ and see whether V_1 has rank r or not.
- (5) If Step 3.1 applies with $r = k_1 + k_2 + 1$, then an rm exists.
- (6) The atoms are computed as in Step 4.
- (7) The densities in Step 5 are obtained by solving the system $W\rho = (\gamma_{-2k_1} \ \gamma_{-2k_1+1} \ \dots \ \gamma_{-2k_1+r-1})^T$, where W is the $r \times r$ Vandermonde matrix with i th row equal to $(x_j^{-2k_1+i-1})_{j=1, \dots, r}$. The coordinates $\rho_j, j = 1, \dots, r$, of ρ are the densities of an rm measure for γ .

3. The TMP on the curves $y = q(x)$

In this section, we study the K -TMP for K being a curve of the form $y = q(x), q \in \mathbb{R}[x]$. In Section 3.1, we first give a solution of the K -TMP, $\deg q \geq 3$, based on the size of positive semidefinite extensions of the moment matrix needed and also bound the number of atoms in the K -rm with the smallest number of atoms (see Theorem 3.1 for the even degree and Theorem 3.2 for the odd degree sequences). As a result, we obtain a sum-of-squares representation for polynomials, which are strictly positive on K (see Corollary 3.3). This improves bounds on the degrees in the previously known result [11, Proposition 6.3].

In Section 3.2, we apply the technique from the proofs of the results from Section 3.1 to give a concrete solution to the TMP on the curve $y = x^2$, which is an alternative solution to the one from [6] in the even case (see Theorem 3.4) and is new in the odd case (see Theorem 3.7). In Section 3.3, we give a solution to the K -TMP based on a feasibility of the corresponding linear matrix inequality (see Theorem 3.8). Finally, in Section 3.4 we concretely solve the TMP on the curve $y = x^3$ in the odd degree case (see Theorem 3.13).

3.1. Solution to the TMP in terms of psd extensions of M_k , bounds on the number of atoms in the minimal measure and a positivstellensatz

Theorem 3.1 (Even case): Let $K := \{(x, y) \in \mathbb{R}^2 : y = q(x)\}$, where $q \in \mathbb{R}[x]$ with $\deg q \geq 3$, and $\beta := \beta^{(2k)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$ with $k \geq \deg q$. The following statements are equivalent:

- (1) β has a K -representing measure.
- (2) β has a s -atomic K -representing measure for some s satisfying

$$\text{rank } M_k \leq s \leq k \deg q.$$

- (3) M_k satisfies $Y = q(X)$ and admits a positive semidefinite, recursively generated extension $M_{k+\deg q-2}$.
- (4) M_k satisfies $Y = q(X)$ and admits a positive semidefinite extension $M_{k+\deg q-1}$.

Theorem 3.2 (Odd case): Let $K := \{(x, y) \in \mathbb{R}^2 : y = q(x)\}$, where $q \in \mathbb{R}[x]$ with $\deg q \geq 3$, and $\beta^{(2k-1)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k-1}$ with $k \geq \deg q$. Then the following statements are equivalent:

- (1) β has a K -representing measure.
- (2) β has a s -atomic K -representing measure for some s satisfying

$$\text{rank } M_{k-1} \leq s \leq k \deg q - \left\lceil \frac{\deg q}{2} \right\rceil.$$

- (3) $\beta^{(2k-1)}$ can be extended to a sequence $\beta^{(2k)}$ such that M_k satisfies $Y = q(X)$ and admits a positive semidefinite, recursively generated extension $M_{k+\deg q-2}$.
- (4) $\beta^{(2k-1)}$ can be extended to a sequence $\beta^{(2k)}$ such that M_k satisfies $Y = q(X)$ and admits a positive semidefinite extension $M_{k+\deg q-1}$.

Remark 3.1: (1) Previous bounds on the size of extensions in (4) of Theorem 3.1: In [52], Curto and Fialkow studied polynomials $p \in \mathbb{R}[x, y]$ for which the existence of the $\mathcal{Z}(p)$ -rm is equivalent to the psd moment matrix extension of some size. In [11, Section 6], the author considered polynomials of the form $p(x, y) = y - q(x)$, where $q \in \mathbb{R}[x]$, and proved that a sequence of degree $2k$ admits a $\mathcal{Z}(p)$ -rm, if M_k admits a psd extension M_{k+r} , where $r = (2k + 1) \deg q - k$ [11, Propositions 6.1, 6.3]. The proof of this result relies on the truncated Riesz–Haviland theorem [52, Theorem 1.2] and a sum-of-squares representations for polynomials, strictly positive on $\mathcal{Z}(p)$ ([11, Proposition 6.3] and [53, Proposition 5.1]). Part (4) of Theorem 3.1

improves Fialkow’s result by decreasing the size of the extensions to $r = \deg q - 1$. We mention that this was known for the case of the curve $y = x^3$ [11, Corollary 5.3].

- (2) *Known bounds on the number of atoms in (2) of Theorems 3.1, 3.2:* In [50], the authors also studied odd degree sequences β , which are moments of a positive Borel measure supported on a plane curve $\mathcal{Z}(p)$, $p \in \mathbb{R}[x, y]$, and proved that every such sequence admits a $(k \deg p)$ -atomic $\mathcal{Z}(p)$ -rm [50, Corollary 7.6]. In the proof, they use their variant of Bézout’s theorem on the number of intersection points of two plane algebraic curves [50, Theorem 7.3]. Part (2) of Theorem 3.1 gives an analogue of [50, Corollary 7.6] for even degree sequences on curves $\mathcal{Z}(y - q(x))$, $\deg q \geq 3$, while part (2) of Theorem 3.2 improves [50, Corollary 7.6] for curves $\mathcal{Z}(y - q(x))$, $\deg q \geq 3$, by decreasing the upper bound on the number of atoms needed by $\lceil \frac{\deg q}{2} \rceil$.
- (3) *Theorem 3.1 in case $\deg q = 2$:* If $\deg q = 2$ in Theorem 3.1, then $y = q_2x^2 + q_1x + q_0 \in \mathbb{R}[x]$ with $q_2 \neq 0$ or equivalently $\frac{1}{q_2}y - q_1x - q_0 = x^2$. By applying an alt $\phi(x, y) = (x, \frac{1}{q_2}y - q_1x - q_0)$ to the sequence β we get a sequence $\tilde{\beta}$ with the moment matrix $M_k(\tilde{\beta})$ satisfying $Y = X^2$. So it is enough to observe the case of a parabola, which was concretely solved in [6] by the use of the FET. The technique used in the proof of Theorem 3.1 can be used to give an alternative proof of the solution from [6] and also obtain a new solvability condition (see Theorem 3.4). This condition will be essentially used in the solution of TMP on the cubic reducible curve $y(y - x^2) = 0$ in our forthcoming work, similarly as for the TMP on the union of three parallel lines [49, Theorem 4.2], where we needed such version of the solution to the TMP on the union of two parallel lines [49, Theorem 3.1]. The upper bound on the number of atoms in a minimal rm is $2k + 1$ and this is sharp (e.g. if M_k has only column relations coming from $Y = X^2$ by rg, then it is of rank $2k + 1$ and so every rm must have at least $2k + 1$ atoms). So the equivalence (1) \Leftrightarrow (2) of Theorem 3.1 does not extend to $\deg q = 2$. (The moment $\gamma_{k \deg q - 1}$ is not independent from β for $\deg q = 2$ as opposed to $\deg q > 2$ and hence in the last step of the proof below decreasing the number of atoms in the rm from $k \deg q + 1$ to $k \deg q$ cannot be done.) Also the equivalence (1) \Leftrightarrow (3) of Theorem 3.1 is not true for $\deg q = 2$, but we need to replace $k + \deg q - 2$ by $k + \deg q - 1$ in (3), because we do not get the information about $\gamma_{2k \deg q + 1} = \gamma_{4k + 1}$ and $\gamma_{2k \deg q + 2} = \gamma_{4k + 2}$ from $M_{k + \deg q - 2} = M_k$ for $\deg q = 2$ as opposed to $\deg q > 2$. However, the equivalence (1) \Leftrightarrow (4) still holds for $\deg q = 2$ with the argument given in Theorem 3.4.
- (4) *Theorem 3.1 in case $\deg q \leq 1$:* If $\deg q \leq 1$ in Theorem 3.1, then $q(x) = ax + by + c$, $a, b, c \in \mathbb{R}$. If $(a, b) \neq (0, 1)$, then the following statements are equivalent:
- (a) β has a K -rm.
 - (b) β has a s -atomic K -rm for some s satisfying

$$\text{rank } M_k \leq s \leq k + 1.$$

- (c) M_k satisfies $Y = aX + bY + c$, is psd and rg.

The equivalence (a) \Leftrightarrow (c) follows from [52, Proposition 3.11], while the equivalence (a) \Leftrightarrow (b) follows from the solution [44, Theorem 3.9] to the \mathbb{R} -TMP, which corresponds to $(M_k)|_{\{1, X, \dots, X^k\}}$. Namely, if the atoms x_1, \dots, x_m represent $\beta_{i,0}$, $i = 0, \dots, 2k$, then the atoms (x_i, y_i) , where $y_i = \frac{1}{1-b}(ax_i + c)$ will represent β if $b \neq 1$. If $b = 1$ and $a \neq 0$, then we change the roles of x and y in the argument above. If

$b = 1$ and $a = 0$, then $y = q(x)$ only makes sense if $c = 0$, but in this case there are no relations in the moment matrix and for $k > 2$ the solution to the TMP is not known (for $k = 2$ the solution is known [9,10]).

- (5) *Uniqueness and description of all solutions in Theorems 3.1 and 3.2:* These questions are nontrivial, being equivalent to the descriptions of psd completions of a partially defined Hankel matrix A_γ , where $\gamma \in \mathbb{R}^{2k\ell+1}$ is a partially defined univariate sequence, i.e. γ_t is defined by the formula (16) below. Namely, the original sequence β determines only those γ_t for which $t \bmod \ell + \lfloor \frac{t}{\ell} \rfloor \leq 2k$ (resp. $2k-1$) in the even (resp. odd) case. However, in a very special case $y = x^3$ the structure of the missing entries is simple enough (only γ_{6k-1} is missing) to answer these questions. The description is not explicitly stated in [48, Theorem 3.1], which solves $y = x^3$ using the univariate reduction technique, but one of the main steps of the proof is [48, Lemma 2.11], which actually describes all psd completions and among them there are one or two minimal ones (in terms of the rank). However, already for $y = x^4$ describing all psd completions concretely and among them minimal ones does not seem to be possible due to the structure of missing entries of A_γ , see Example 3.11.
- (6) *Complexity of checking conditions in (4) of Theorems 3.1 and 3.2:* Checking if $Y = q(X)$ is a column relation only requires checking whether the corresponding vector is in the kernel for M_k , while the existence of a psd extension $M_{k+\deg q-1}$ of M_k is a feasibility question of a semidefinite programme (SDP) in the variables $\beta_{i,j}$ with $2k < i + j \leq 2(k + \deg q - 1)$, $i, j \in \mathbb{Z}_+$, i.e. an SDP with matrices of size $\binom{k+\deg q+1}{2}$ in $\binom{2(k+\deg q)}{2} - \binom{2(k+1)}{2} = (\deg q - 1)(4k + 2 \deg q + 1)$ variables. The complexity of the SDP feasibility question is still unknown (see [54,55]), but for a fixed number of variables or size of matrices it has polynomial time complexity [56]. By Theorem 3.8, the feasibility question in (4) of Theorems 3.1 and 3.2 is equivalent to the feasibility question of a smaller SDP, i.e. the matrices are of size $k \deg q + 2$ and the number of variables is $\frac{1}{2}(\deg q - 2)(\deg q - 1) + 2$.

Remark 3.2 (Basic idea of the proof of the implication (3) \Rightarrow (2) of Theorem 3.1): The main steps are the following:

- (1) Due to the column relation $Y = q(X)$ satisfied by M_k and M_ℓ being rg, the column space of $M_{k+\ell-2}$ is spanned by the set \mathcal{B} of columns, indexed by monomials $Y^i X^j$, where $i = 0, \dots, k, j = 0, \dots, \deg q - 1$, and $i + j \leq k + \ell - 2$.
- (2) Writing $q(x) = \sum_{i=0}^{\ell} q_i x^i$, the point (1) implies, on the level of the sequence $\beta_{i,j}$, the following linear relations:

$$\beta_{i,j} = q_\ell \beta_{i+\ell,j-1} + q_{\ell-1} \beta_{i+\ell-1,j-1} + \dots + q_0 \beta_{i,j-1}, \tag{7}$$

where $i \in \mathbb{Z}_+, j \in \mathbb{N}$ and $i + j \leq 2(k + \ell - 2)$.

- (3) Using relations (7) one can show that the sequence

$$\beta_{i,j}, \quad i, j \in \mathbb{Z}_+, \quad i + j \leq 2(k + \ell - 2),$$

can be uniquely parametrized by

$$\gamma_i := \beta_{i,0}, \quad i = 0, \dots, 2k\ell + 2.$$

(See (16) and Claim 4 in the proof of Theorem 3.1 below.)

- (4) The next step is to note that the original sequence $\beta_{i,j}, i, j \in \mathbb{Z}_+, i + j \leq 2k$ has a K -rm if and only if the univariate sequence $\gamma_0, \gamma_1, \dots, \gamma_{2k\ell}$ has an \mathbb{R} -rm. (See Claim 5 in the proof of Theorem 3.1.)
- (5) Now one can use the solution to the \mathbb{R} -TMP due to Curto and Fialkow [44, Theorem 3.9] and derive the solution to the K -TMP. Namely, the sequence $\gamma^{(0,2k\ell)} = (\gamma_0, \gamma_1, \dots, \gamma_{2k\ell}) \in \mathbb{R}^{2k\ell+1}$ has a $(\text{rank } \gamma^{(0,2k\ell)})$ -atomic \mathbb{R} -rm iff a Hankel matrix $A_{\gamma^{(2k\ell+2)}}$ is psd for some extension $\gamma^{(2k\ell+2)} = (\gamma, \gamma_{2k\ell+1}, \gamma_{2k\ell+2}) \in \mathbb{R}^{2k\ell+3}$ of $\gamma^{(0,2k\ell)}$. Observing that the restriction $(M_{k+\ell-2})|_{\mathcal{B}}$ is equal to $PA_{\gamma^{(2k\ell+2)}}P^T$ for a certain matrix P (See Claim 6 in the proof of Theorem 3.1 below.), translates the condition of $A_{\gamma^{(2k\ell+2)}}$ being psd, to $(M_{k+\ell-2})|_{\mathcal{B}}$ (and hence $M_{k+\ell-2}$) being psd.
- (6) The last step is to decrease the number of atoms in the solution to the \mathbb{R} -TMP for $\gamma^{(0,2k\ell)}$ by 1 in case $A_{\gamma^{(0,2k\ell)}}$ has full rank, i.e. $\text{rank } A_{\gamma^{(0,2k\ell)}} = k\ell + 1$. This can be achieved in the following three steps:
 - (a) Applying an appropriate alt ϕ to β , one can assume that the coefficient $q_{\ell-1}$ of $q(x)$ is equal to 0. (See Claim 1 in the proof of Theorem 3.1.)
 - (b) Due to (a) none of the moments of the original sequence $\beta_{i,j}, i, j \in \mathbb{Z}_+, i + j \leq 2k$, depends on $\gamma_{2k\ell-1}$.
 - (c) One can replace $\gamma_{2k\ell-1}$ with $\tilde{\gamma}_{2k\ell-1}$ to obtain a sequence $\tilde{\gamma}^{(0,2k\ell)}$ with a $(k\ell)$ -atomic \mathbb{R} -rm.

Proof of Theorem 3.1.: Before starting a proof we do an alt ϕ which will be used in the proof of the implication (3) \Rightarrow (2) to justify in an easier way that the upper bound in (2) is $k \deg q$ instead of $k \deg q + 1$. We write $\ell := \deg q$ and let $q(x) = \sum_{i=0}^{\ell} q_i x^i$, where $q_{\ell} \neq 0$ and each $q_i \in \mathbb{R}$.

Claim 1: We may assume that $q_{\ell-1} = 0$.

Proof of Claim 1.: Defining $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $\phi(x, y) = (x + \frac{q_{\ell-1}}{\ell q_{\ell}}, y) =: (\tilde{x}, y)$, note that the relation $y = q(x)$ becomes

$$\begin{aligned}
 y &= q\left(\tilde{x} - \frac{q_{\ell-1}}{\ell q_{\ell}}\right) = \sum_{i=0}^{\ell} q_i \left(\tilde{x} - \frac{q_{\ell-1}}{\ell q_{\ell}}\right)^i \\
 &= q_{\ell} \tilde{x}^{\ell} + \underbrace{\left(-q_{\ell} \cdot \ell \frac{q_{\ell-1}}{\ell q_{\ell}} + q_{\ell-1}\right)}_{=0} \tilde{x}^{\ell-1} + \sum_{i=0}^{\ell-2} \tilde{q}_i \tilde{x}^i,
 \end{aligned}$$

for some $\tilde{q}_0, \dots, \tilde{q}_{\ell-2} \in \mathbb{R}$. Since the solution of the K -TMP is invariant under applying ϕ by Proposition 2.1, the conclusion of Claim 1 follows. ■

Now we start the proof of the theorem. The implications (1) \Rightarrow (4) and (2) \Rightarrow (1) are trivial. The implication (4) \Rightarrow (3) is [1, Theorem 3.14]. It remains to prove the implication

(3) \Rightarrow (2). Assume that M_k admits a psd, rg extension $M_{k+\ell-2}$. Let

$$\mathcal{B} = \left\{ 1, x, \dots, x^{\ell-1}, y, yx, \dots, yx^{\ell-1}, \dots, y^{k-1}, \dots, y^{k-1}x^{\ell-1}, y^k, y^k x \right\} \tag{8}$$

be a set of monomials and V a vector subspace in $\mathbb{R}[x, y]_{k+\ell-2}$ generated by the set \mathcal{B} . Since $M_{k+\ell-2}$ satisfies $X^i Y^j = X^i q(X)^j$ for every $i, j \in \mathbb{Z}_+$ such that $i + j \leq k + \ell - 2$, it follows that the columns from \mathcal{B} span $\mathcal{C}(M_{k+\ell-2})$. Let $p(x, y) = \sum_{i,j} p_{ij} x^i y^j \in V$ be a polynomial and \widehat{p} a vector of its coefficients ordered in the basis \mathcal{B} . Before we define a univariate polynomial $g_p(x)$ corresponding to $p(x, y)$, we prepare some computations. We have that

$$\begin{aligned} x^i (q(x))^j &= x^i \left(\sum_{0 \leq i_1, \dots, i_j \leq \ell} q_{i_1} q_{i_2} \cdots q_{i_j} x^{i_1 + \dots + i_j} \right) \\ &= \sum_{p=0}^{j\ell} \left(\sum_{\substack{0 \leq i_1, \dots, i_j \leq \ell, \\ i_1 + \dots + i_j = p}} q_{i_1} q_{i_2} \cdots q_{i_j} \right) x^{i+p} \\ &= \sum_{s=i}^{i+j\ell} q_{i,j,s} x^s, \end{aligned} \tag{9}$$

for all $i, j \in \mathbb{Z}_+$, where

$$q_{i,j,s} = \begin{cases} \sum_{\substack{0 \leq i_1, \dots, i_j \leq \ell, \\ i_1 + \dots + i_j = s-i}} q_{i_1} q_{i_2} \cdots q_{i_j}, & \text{if } i \leq s \leq i + j\ell, \\ 0, & \text{otherwise.} \end{cases} \tag{10}$$

Later on we will need the following observation about the numbers $q_{i,j,s}$.

Claim 2: Let $i_1, i_2, j_1, j_2, s \in \mathbb{Z}_+$. Then

$$q_{i_1+i_2, j_1+j_2, s} = \sum_{t=i_1}^s q_{i_1, j_1, t} q_{i_2, j_2, s-t}. \tag{11}$$

Proof of Claim 2.: We write $m_1 := i_1 + i_2$ and $m_2 := i_1 + i_2 + (j_1 + j_2)\ell$. We separate two cases: $s \in \{m_1, m_1 + 1, \dots, m_2\}$ and $s \notin \{m_1, m_1 + 1, \dots, m_2\}$.

Case 1: $s \in \{m_1, m_1 + 1, \dots, m_2\}$. We have that

$$\begin{aligned} q_{i_1+i_2, j_1+j_2, s} &= \sum_{\substack{0 \leq k_1, \dots, k_{j_1+j_2} \leq \ell, \\ k_1 + \dots + k_{j_1+j_2} = s-i_1-i_2}} q_{k_1} q_{k_2} \cdots q_{k_{j_1}} q_{k_{j_1+1}} \cdots q_{k_{j_1+j_2}} \\ &= \sum_{t=i_1}^s \sum_{\substack{0 \leq k_1, \dots, k_{j_1} \leq \ell, \\ k_1 + \dots + k_{j_1} = t-i_1}} \sum_{\substack{0 \leq k_{j_1+1}, \dots, k_{j_2} \leq \ell, \\ k_{j_1+1} + \dots + k_{j_2} = s-t-i_2}} q_{k_1} q_{k_2} \cdots q_{k_{j_1}} q_{k_{j_1+1}} \cdots q_{k_{j_1+j_2}} \end{aligned}$$

$$\begin{aligned}
 &=^3 \sum_{t=i_1}^s \left(\sum_{\substack{0 \leq k_1, \dots, k_{j_1} \leq \ell, \\ k_1 + \dots + k_{j_1} = t - i_1}} q_{k_1} q_{k_2} \cdots q_{k_{j_1}} \right) \left(\sum_{\substack{0 \leq k_{j_1+1}, \dots, k_{j_2} \leq \ell, \\ k_{j_1+1} + \dots + k_{j_2} = s - t - i_2}} q_{k_{j_1+1}} \cdots q_{k_{j_1+j_2}} \right) \\
 &=^4 \sum_{t=i_1}^s q_{i_1, j_1, t} q_{i_2, j_2, s-t},
 \end{aligned}$$

where the first equality follows by definition (10) of $q_{i_1+i_2, j_1+j_2, s}$, in the second we decomposed the sum into three sums, in the third we used independence of the inner two sums, while the last equality follows by definitions (10) of $q_{i_1, j_1, t}$ and $q_{i_2, j_2, s-t}$.

Case 2: $s \notin \{m_1, m_1 + 1, \dots, m_2\}$. For $s > m_2$ we have $q_{i_1+i_2, j_1+j_2, s} = 0$ and

$$\sum_{t=i_1}^s q_{i_1, j_1, t} q_{i_2, j_2, s-t} = \sum_{t=i_1}^{i_1+j_1\ell} q_{i_1, j_1, t} \underbrace{q_{i_2, j_2, s-t}}_{\substack{=0, \text{ since} \\ s-t > i_2+j_2\ell}} + \sum_{t=i_1+j_1\ell+1}^s \underbrace{q_{i_1, j_1, t}}_{\substack{=0, \text{ since} \\ t > i_1+j_1\ell}} q_{i_2, j_2, s-t} = 0,$$

which implies that (10) holds. Similarly, for $s < m_1$ we again have $q_{i_1+i_2, j_1+j_2, s} = 0$ and $\sum_{t=i_1}^s q_{i_1, j_1, t} q_{i_2, j_2, s-t} = 0$, since $q_{i_2, j_2, s-t} = 0$ for every t due to $s - t < i_2$. Also in this case (10) holds. ■

Now we define a univariate polynomial $g_p(x)$ corresponding to $p(x, y)$ by

$$g_p(x) := p(x, q(x)) = \sum_{i,j} p_{ij} \sum_{s=i}^{i+j\ell} q_{i,j,s} x^s =: \sum_{s=0}^{k\ell+1} g_{p,s} x^s \in \mathbb{R}[x]_{k\ell+1},$$

where we used (9) in the second equality.

Let \widehat{g}_p be its vector of coefficients in the basis

$$\mathcal{B}_1 = \{1, x, \dots, x^{k\ell+1}\}. \tag{12}$$

The following claim expresses \widehat{g}_p by \widehat{p} .

Claim 3: It holds that

$$\widehat{g}_p = P^T \widehat{p}, \tag{13}$$

where

$$P = \begin{pmatrix} I_\ell & 0 & \cdots & \cdots & 0 & 0 \\ P[1, 0] & P[1, 1] & 0 & & 0 & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots & \vdots \\ \vdots & & \ddots & \ddots & 0 & \vdots \\ P[k-1, 0] & P[k-1, 1] & \cdots & \cdots & P[k-1, k-1] & 0 \\ Q[0] & Q[1] & \cdots & \cdots & Q[k-1] & Q[k] \end{pmatrix} \in \mathbb{R}^{(k\ell+2) \times (k\ell+2)}$$

and

$$\begin{aligned}
 P[c, d] &= \begin{pmatrix} q_{0,c, dl} & q_{0,c, dl+1} & \cdots & q_{0,c, dl+l-1} \\ q_{1,c, dl} & q_{1,c, dl+1} & \cdots & q_{1,c, dl+l-1} \\ \vdots & \cdots & \cdots & \vdots \\ q_{\ell-1,c, dl} & q_{\ell-1,c, dl+1} & \cdots & q_{\ell-1,c, dl+l-1} \end{pmatrix} \in \mathbb{R}^{\ell \times \ell} \quad \text{for each } c, d, \\
 Q[d] &= \begin{pmatrix} q_{0,k, dl} & q_{0,k, dl+1} & \cdots & q_{0,k, dl+l-1} \\ q_{1,k, dl} & q_{1,k, dl+1} & \cdots & q_{1,k, dl+l-1} \end{pmatrix} \in \mathbb{R}^{2 \times \ell} \quad \text{for } d = 0, \dots, k-1, \\
 Q[k] &= \begin{pmatrix} (q_\ell)^k & 0 \\ q_{1,k, k\ell} & (q_\ell)^k \end{pmatrix} \in \mathbb{R}^{2 \times 2}.
 \end{aligned}$$

Proof of Claim 3.: We write $\vec{v}_{x,y}$ for the vector of monomials $x^i y^j$ from the basis \mathcal{B} (see (8)) and \vec{v}_x for the vector of monomials from the basis \mathcal{B}_1 (see (12)). We have that $p(x, y) = (\vec{v}_{x,y})^T \widehat{p}$ and $g_p(x) = (\vec{v}_x)^T \widehat{g}_p$. By (9) it follows that $\vec{v}_{x,y} = P\vec{v}_x$. Hence, the definition of g_p implies that $g_p(x) = (P\vec{v}_x)^T \widehat{p} = (\vec{v}_x)^T P^T \widehat{p}$. Thus $\widehat{g}_p = P^T \widehat{p}$, which proves Claim 3. ■

Note that

$$q_{i,j,i+j\ell} = (q_\ell)^j \neq 0 \tag{14}$$

and hence we can express $x^{i+j\ell}$ from (9) by the formula

$$x^{i+j\ell} = \frac{1}{(q_\ell)^j} \left(x^i (q(x))^j - \sum_{s=0}^{i+j\ell-1} q_{i,j,s} x^s \right). \tag{15}$$

We define two univariate sequences

$$\begin{aligned}
 \gamma &:= \gamma^{(0,2k\ell)} = (\gamma_0, \gamma_1, \dots, \gamma_{2k\ell}) \in \mathbb{R}^{2k\ell+1}, \\
 \tilde{\gamma} &:= \gamma^{(2k\ell+2)} = (\gamma, \gamma_{2k\ell+1}, \gamma_{2k\ell+2}) \in \mathbb{R}^{2k\ell+3},
 \end{aligned}$$

recursively for $t = 0, 1, \dots, 2k\ell + 2$ by the formula

$$\gamma_t = \frac{1}{(q_\ell)^{\lfloor \frac{t}{\ell} \rfloor}} \left(\beta_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor} - \sum_{s=0}^{t-1} q_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor, s} \cdot \gamma_s \right). \tag{16}$$

Note that $t \bmod \ell + \lfloor \frac{t}{\ell} \rfloor \leq \ell - 1 + 2k$ (here we used that $\ell \geq 3$ and thus $\lfloor \frac{t}{\ell} \rfloor \leq 2k$) and so $\beta_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor}$ is well-defined being an element of the matrix $M_{k+\ell-2}$ (since $2(k + \ell - 2) \geq \ell - 1 + 2k$ for $\ell \geq 3$).

Note that we defined γ_t by (16) only using $\beta_{i,j}$ with $i < \ell$. The following claim proves we could define γ_t using any i, j with $t = i + j\ell$.

Claim 4: Let $t \in \{0, 1, \dots, 2k\ell + 2\}$ and $t = i + j\ell$ with $i, j \in \mathbb{Z}_+, i + j \leq 2(k + \ell - 2)$. Then

$$\gamma_t = \frac{1}{(q_\ell)^j} \left(\beta_{i,j} - \sum_{s=0}^{t-1} q_{i,j,s} \cdot \gamma_s \right). \tag{17}$$

Proof of Claim 4.: If $i < \ell$, then (17) follows from definition (16) of γ_t . Let $i_0 \geq \ell$. Assume (17) is true for every $i = 0, 1, \dots, i_0 - 1$ and $j \in \mathbb{N}_0$ such that $i + j \leq 2(k + \ell - 2)$. We will prove it is true for i_0 and every $j \in \mathbb{N}_0$ such that $i_0 + j \leq 2(k + \ell - 2)$. We have that:

$$\begin{aligned} \beta_{i_0,j} &= \frac{1}{q_\ell} \left(\beta_{i_0-\ell,j+1} - \sum_{s=0}^{\ell-1} q_s \beta_{i_0-\ell+s,j} \right) \\ &= \frac{1}{q_\ell} \left(\sum_{s=i_0-\ell}^{i_0+j\ell} q_{i_0-\ell,j+1,s} \gamma_s - \sum_{s=0}^{\ell-1} q_s \left(\sum_{u=i_0-\ell+s}^{i_0+s+(j-1)\ell} q_{i_0-\ell+s,j,u} \gamma_u \right) \right) \end{aligned} \tag{18}$$

where in the first equality we used that $\beta^{(2(k+\ell-2))} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2(k+\ell-2)}$ is rg and $Y = q(X)$ is a column relation, and in the second we used the induction hypothesis together with the fact that $q_{i,j,s} = 0$ if $s < i$ and $q_{i_0-\ell,j+1,i_0+j\ell} = (q_\ell)^{j+1}$. Note that

$$\begin{aligned} \sum_{s=0}^{\ell-1} q_s \left(\sum_{u=i_0-\ell+s}^{i_0+s+(j-1)\ell} q_{i_0-\ell+s,j,u} \gamma_u \right) &= \sum_{s=0}^{\ell-1} q_s \left(\sum_{u=i_0-\ell+s}^{i_0+s+(j-1)\ell} q_{i_0-\ell,j,u-s} \gamma_u \right) \\ &= \sum_{s=0}^{\ell-1} \left(\sum_{u=i_0-\ell+s}^{i_0+s+(j-1)\ell} q_s q_{i_0-\ell,j,u-s} \gamma_u \right) \\ &= \sum_{u=i_0-\ell}^{i_0+j\ell-1} q_{i_0-\ell,j+1,u} \gamma_u - \sum_{u=i_0-\ell}^{i_0+j\ell-1} q_\ell q_{i_0-\ell,j,u-\ell} \gamma_u \end{aligned} \tag{19}$$

where in the first equality we used that $q_{i_0-\ell+s,j,u} = q_{i_0-\ell,j,u-s}$ by definition (10), in the second we moved q_s inside the inner sum, in the third we used that for a fixed u , as s runs from 0 to $\ell - 1$, the coefficients at γ_u run over all terms $q_{i_1} \cdots q_{i_{j+1}}$ such that $i_1 + \dots + i_{j+1} = u - i_0 + \ell$ except those where $q_{i_1} = q_\ell$. But if $q_{i_1} = q_\ell$, all terms $q_{i_2} \cdots q_{i_{j+1}}$ such that $i_2 + \dots + i_{j+1} = u - i_0$ sum up to $q_{i_0-\ell,j,u-\ell}$. Using (19) in (18) we get

$$\begin{aligned} \beta_{i_0,j} &= \frac{1}{q_\ell} \left(q_{i_0-\ell,j+1,i_0+j\ell} \gamma_{i_0+j\ell} + \sum_{u=i_0-\ell}^{i_0+j\ell-1} q_\ell q_{i_0-\ell,j,u} \gamma_u \right) \\ &= \frac{1}{q_\ell} \left((q_\ell)^{j+1} \gamma_{i_0+j\ell} + \sum_{u=i_0-\ell}^{i_0+j\ell-1} q_\ell q_{i_0-\ell,j,t} \gamma_u \right) \\ &= (q_\ell)^j \gamma_{i_0+j\ell} + \sum_{u=i_0-\ell}^{i_0+j\ell-1} q_{i_0-\ell,j,u} \gamma_u. \end{aligned} \tag{20}$$

where in the second equality we used that $q_{i_0-\ell,j+1,i_0+j\ell} = (q_\ell)^{j+1}$ and in the last we put $\frac{1}{q_\ell}$ inside the bracket. But (20) is (17) for $t = i_0 + j\ell$. ■

By the following claim solving the K -TMP for β is equivalent to solving the \mathbb{R} -TMP for γ .

Claim 5: Let $u \in \mathbb{N}$. A sequence γ admits a u -atomic \mathbb{R} -rm if and only if β admits a u -atomic K -rm.

Proof of Claim 5.: First we prove the implication (\Rightarrow) . Let x_1, \dots, x_u , be the atoms in the \mathbb{R} -rm for γ with the corresponding densities ρ_1, \dots, ρ_u . We will prove that the atoms $(x_1, q(x_1)), \dots, (x_u, q(x_u))$ with densities ρ_1, \dots, ρ_p are the K -rm for β . We use induction on the index i in $\beta_{i,j}$, where $i + j \leq 2k$. For $i < \ell$ and any j such that $i + j \leq 2k$ we have that

$$\begin{aligned} \beta_{i,j} &=^1 (q_\ell)^j \gamma_{i+j\ell} + \sum_{s=0}^{i+j\ell-1} q_{i,j,s} \gamma_s \\ &=^2 (q_\ell)^j \left(\sum_{p=0}^u \rho_p (x_p)^{i+j\ell} \right) + \sum_{s=0}^{i+j\ell-1} q_{i,j,s} \left(\sum_{p=0}^u \rho_p (x_p)^s \right) \\ &=^3 \sum_{p=0}^u \rho_p (q_\ell)^j (x_p)^{i+j\ell} + \sum_{p=0}^u \left(\rho_p \sum_{s=0}^{i+j\ell-1} q_{i,j,s} (x_p)^s \right) \\ &=^4 \sum_{p=0}^u \left(\rho_p \left((q_\ell)^j (x_p)^{i+j\ell} + \sum_{s=0}^{i+j\ell-1} q_{i,j,s} (x_p)^s \right) \right) \\ &=^5 \sum_{p=0}^u (\rho_p (x_p)^i (q(x_p))^j), \end{aligned}$$

where we used (16) with $t = i + j\ell$ in the first equality noticing that

$$i + j\ell = i + j + j(\ell - 1) \leq 2k + 2k(\ell - 1) \leq 2k\ell,$$

implying well-definedness of γ_s by s being bounded above by $2k\ell$, the definitions of ρ_p, x_p in the second equality, we interchanged the order of summation in the third and fourth equalities and in the last we used (15) for $x = x_p$. So the atoms $(x_1, q(x_1)), \dots, (x_u, q(x_u))$ with densities ρ_1, \dots, ρ_p indeed represent $\beta_{i,j}$ for $i < \ell$ and any j such that $i + j \leq 2k$. We now assume that this holds for all $i = 0, \dots, m$ and j such that $i + j \leq 2k$, where $m \geq \ell - 1$ and prove it for $i = m + 1$ and any $j \leq 2k - i$. We have that

$$\begin{aligned} \beta_{m+1,j} &=^1 \frac{1}{q_\ell} \left(\beta_{m+1-\ell,j+1} - \sum_{s=0}^{\ell-1} q_s \beta_{m+1-\ell+s,j} \right) \\ &=^2 \frac{1}{q_\ell} \left(\left(\sum_{p=0}^u \rho_p (x_p)^{m+1-\ell} (q(x_p))^{j+1} \right) - \sum_{s=0}^{\ell-1} q_s \left(\sum_{p=0}^u \rho_p (x_p)^{m+1-\ell+s} (q(x_p))^j \right) \right) \\ &=^3 \frac{1}{q_\ell} \sum_{p=0}^u \left(\rho_p \left((x_p)^{m+1-\ell} (q(x_p))^{j+1} - \sum_{s=0}^{\ell-1} q_s (x_p)^{m+1-\ell+s} (q(x_p))^j \right) \right) \end{aligned}$$

$$\begin{aligned}
 &=^4 \frac{1}{q^\ell} \sum_{p=0}^u \left(\rho_p(x_p)^{m+1-\ell} (q(x_p))^j \left(q(x_p) - \sum_{s=0}^{\ell-1} q_s(x_p)^s \right) \right) \\
 &=^5 \sum_{p=0}^u (\rho_p(x_p)^{m+1} (q(x_p))^j),
 \end{aligned}$$

where we used that β is rg in the first equality, the induction hypothesis in the second, in the third we interchanged the order of summation, factored out $(x_p)^{m+1-\ell} (q(x_p))^j$ in the fourth and in the last we used that $q(x) - \sum_{s=0}^{\ell-1} q_s x^s = q_\ell x^\ell$ by definition of q . This proves the implication (\Rightarrow) .

It remains to prove the implication (\Leftarrow) . Let $(x_1, q(x_1)), \dots, (x_u, q(x_u))$ be the atoms in the K -rm for β with the corresponding densities ρ_1, \dots, ρ_u . We will prove that the atoms (x_1, \dots, x_u) with densities ρ_1, \dots, ρ_p are the \mathbb{R} -rm for γ . We use induction on the index t in γ_t . For $t = 0$ the claim is trivial, since $\gamma_0 = \beta_{0,0} = \sum_{p=0}^u \rho_u$. We now assume that the claim holds for all $t-1$ with $0 \leq t-1 \leq 2k\ell - 1$ and prove it for t . We have that

$$\begin{aligned}
 \gamma_t &=^1 \frac{1}{(q_\ell)^{\lfloor \frac{t}{\ell} \rfloor}} \left(\beta_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor} - \sum_{s=0}^{t-1} q_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor, s} \cdot \gamma_s \right) \\
 &=^2 \frac{1}{(q_\ell)^{\lfloor \frac{t}{\ell} \rfloor}} \left(\sum_{p=0}^u \rho_p(x_p)^{t \bmod \ell} (q(x_p))^{\lfloor \frac{t}{\ell} \rfloor} - \sum_{s=0}^{t-1} q_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor, s} \cdot \left(\sum_{p=0}^u \rho_p(x_p)^s \right) \right) \\
 &=^3 \frac{1}{(q_\ell)^{\lfloor \frac{t}{\ell} \rfloor}} \sum_{p=0}^u \left(\rho_p \left((x_p)^{t \bmod \ell} (q(x_p))^{\lfloor \frac{t}{\ell} \rfloor} - \sum_{s=0}^{t-1} q_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor, s} \cdot (x_p)^s \right) \right) \\
 &=^4 \frac{1}{(q_\ell)^{\lfloor \frac{t}{\ell} \rfloor}} \sum_{p=0}^u \left(\rho_p q_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor, t \bmod \ell + \lfloor \frac{t}{\ell} \rfloor \ell} \cdot (x_p)^{t \bmod \ell + \lfloor \frac{t}{\ell} \rfloor \ell} \right) \\
 &=^5 \frac{1}{(q_\ell)^{\lfloor \frac{t}{\ell} \rfloor}} \sum_{p=0}^u \left(\rho_p (q_\ell)^{\lfloor \frac{t}{\ell} \rfloor} (x_p)^t \right) \\
 &=^6 \sum_{p=0}^u \rho_p(x_p)^t,
 \end{aligned}$$

where we used the definition (16) of γ_t in the first equality, the definitions of ρ_p, x_p and the induction hypothesis in the second equality, we interchanged the order of summation in the third equality, used (9) for $(i, j) = (t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor)$ in the fourth equality and the observation (14) for $(i, j) = (t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor)$ in the fifth equality. This proves the implication (\Leftarrow) . ■

Let $(M_{k+\ell-2})|_{\mathcal{B}}$ be the restriction of $M_{k+\ell-2}$ to the rows and columns indexed by monomials (capitalized) from \mathcal{B} . The following claim gives an explicit connection between $(M_{k+\ell-2})|_{\mathcal{B}}$ and the Hankel matrix $A_{\tilde{\gamma}}$ of the sequence $\tilde{\gamma}$.

Claim 6: We have that

$$(M_{k+\ell-2})|_{\mathcal{B}} = PA_{\tilde{\gamma}}P^T. \tag{21}$$

Proof of Claim 6.: Let $p(x, y) = \sum_{i,j} p_{ij}x^i y^j \in V$ and $r(x, y) = \sum_{i,j} r_{ij}x^i y^j \in V$ be polynomials from the vector subspace V and \widehat{p}, \widehat{r} vectors of their coefficients ordered in the basis \mathcal{B} (see (8)). Let $\widetilde{\beta} := \beta^{(2(k+\ell-2))}$. Then we have

$$\begin{aligned} (\widehat{r})^T ((M_{k+\ell-2})|_{\mathcal{B}}) \widehat{p} &=^1 L_{\widetilde{\beta}}(pr) = L_{\widetilde{\beta}} \left(\sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} r_{i_2 j_2} x^{i_1+i_2} y^{j_1+j_2} \right) \\ &=^2 \sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} r_{i_2 j_2} \beta_{i_1+i_2, j_1+j_2} \\ &=^3 \sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} r_{i_2 j_2} \left(\sum_{s=i_1+i_2}^{i_1+i_2+(j_1+j_2)\ell} q_{i_1+i_2, j_1+j_2, s} \gamma_s \right) \\ &=^4 \sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} r_{i_2 j_2} \left(\sum_{s=i_1+i_2}^{i_1+i_2+(j_1+j_2)\ell} \left(\sum_{t=i_1}^s q_{i_1, j_1, t} q_{i_2, j_2, s-t} \right) \gamma_s \right) \\ &=^5 \sum_{i_1, i_2, j_1, j_2} \left(\sum_{s=i_1+i_2}^{i_1+i_2+(j_1+j_2)\ell} \left(\sum_{t=i_1}^s p_{i_1 j_1} q_{i_1, j_1, t} r_{i_2 j_2} q_{i_2, j_2, s-t} \right) \gamma_s \right) \\ &=^6 L_{\widetilde{\gamma}} \left(\sum_{i_1, i_2, j_1, j_2} \left(\sum_{s=i_1+i_2}^{i_1+i_2+(j_1+j_2)\ell} \left(\sum_{t=i_1}^s p_{i_1 j_1} q_{i_1, j_1, t} r_{i_2 j_2} q_{i_2, j_2, s-t} \right) x^s \right) \right) \\ &=^7 L_{\widetilde{\gamma}} \left(\sum_{i_1, i_2, j_1, j_2} \left(\sum_{s=i_1+i_2}^{i_1+i_2+(j_1+j_2)\ell} \sum_{t=i_1}^s p_{i_1 j_1} q_{i_1, j_1, t} x^t \cdot r_{i_2 j_2} q_{i_2, j_2, s-t} x^{s-t} \right) \right) \\ &=^8 L_{\widetilde{\gamma}} \left(\sum_{i_1, i_2, j_1, j_2} \left(\sum_{t=i_1}^{i_1+j_1\ell} p_{i_1 j_1} q_{i_1, j_1, t} x^t \right) \left(\sum_{u=i_2}^{i_2+j_2\ell} r_{i_2 j_2} q_{i_2, j_2, u} x^u \right) \right) \\ &=^9 L_{\widetilde{\gamma}} \left(\left(\underbrace{\left(\sum_{i_1, j_1} \sum_{t=i_1}^{i_1+j_1\ell} p_{i_1 j_1} q_{i_1, j_1, t} x^t \right)}_{g_p(x)} \right) \left(\underbrace{\left(\sum_{i_2, j_2} \sum_{s=i_1+i_2}^{i_2+j_2\ell} r_{i_2 j_2} q_{i_2, j_2, s} x^s \right)}_{g_r(x)} \right) \right) \\ &=^{10} \widehat{g}_r^T A_{\widetilde{\gamma}} \widehat{g}_p = (P^T \widehat{r})^T A_{\widetilde{\gamma}} (P^T \widehat{p}) = \widehat{r}^T (PA_{\widetilde{\gamma}}P^T) \widehat{p}, \end{aligned}$$

where in the first line we used the correspondence between the moment matrix and the Riesz functional $L_{\widetilde{\beta}}$, the definition $L_{\widetilde{\beta}}$ in the second, Claim 4 in the third, Claim 2 in the fourth, we moved the factor $p_{i_1 j_1} r_{i_2 j_2}$ into the inner sum in the fifth, used the definition of $L_{\widetilde{\gamma}}$ in the sixth, split x^s into two parts and moved it into the inner sum in the seventh, decomposed a double sum into the product of two sums in the eight using that $q_{i_1, j_1, t}$ is

nonzero only for $t \leq i_1 + j_1\ell$ and $q_{i_2,j_1,u}$ is nonzero only for $u \leq i_2 + j_2\ell$, decomposed a sum into the product of two sums using independence of the factors in the ninth line, in the tenth we used the correspondence between $A_{\tilde{\gamma}}$ and the Riesz functional $L_{\tilde{\gamma}}$, where $\widehat{g}_p, \widehat{g}_r$ are the vectors of coefficients of g_p and g_r in the basis \mathcal{B}_1 (see (12)) and also Claim 3. Since p and q were arbitrary from V , this proves Claim 6. ■

Note that

$$P[c, c] = \begin{pmatrix} (q_\ell)^c & 0 & \cdots & \cdots & 0 \\ q_{1,c,cl} & (q_\ell)^c & 0 & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & 0 \\ q_{\ell-1,c,cl} & \cdots & \cdots & q_{\ell-1,c,cl+\ell-1} & q_\ell^c \end{pmatrix} \in \mathbb{R}^{\ell \times \ell} \quad \text{for } c = 1, \dots, k-1.$$

Since P is a lower triangular matrix with all nonzero diagonal entries, it is invertible. Claim 5 implies that

$$A_{\tilde{\gamma}} = P^{-1} \left((M_{k+\ell-2})|_{\mathcal{B}} \right) (P^{-1})^T. \tag{22}$$

Since $(M_{k+\ell-2})|_{\mathcal{B}}$ is psd, it follows from (22) that $A_{\tilde{\gamma}}$ is also psd. We separate two cases. Either $A_{\tilde{\gamma}}$ is pd or $A_{\tilde{\gamma}}$ is singular. In the first case in particular $A_{\tilde{\gamma}}(k\ell) = A_\gamma$ is pd, while in the second case $A_{\tilde{\gamma}}(k\ell)$ is psd and prg by [44, Theorem 2.6]. By [44, Theorem 3.9], γ admits a $(\text{rank } A_\gamma)$ -atomic \mathbb{R} -rm. Since $\text{rank } M_k \leq \text{rank } A_\gamma \leq k\ell + 1$, using Claim 4 the following holds:

(2') β has an s -atomic K -rm for some s satisfying

$$\text{rank } M_k \leq s \leq A_\gamma \leq k\ell + 1. \tag{23}$$

To obtain (2) of Theorem 3.1 we need to decrease the upper bound in (23) by 1. Note that the bound $k\ell + 1$ occurs only in the case A_γ is pd, which we assume in the rest of the proof. We denote by $\gamma(z)$ a sequence obtained from the sequence γ by replacing $\gamma_{2k\ell-1}$ with a variable z . The matrix $A_{\gamma(z)}$ is a partially pd matrix and by [48, Lemma 2.11] there exist two choices of z , which we denote by z^\pm , such that $A_{\gamma(z^\pm)}$ is psd and has rank $k\ell$. Since $\text{rank } A_{\gamma(z^\pm)}(k\ell - 1) = \text{rank } A_{\gamma(z^\pm)} = k\ell$, the sequence $\gamma(z^\pm)$ is prg and by [44, Theorem 3.9] it admits a $k\ell$ -atomic \mathbb{R} -rm. If none of the moments β_{i_j} of the sequence β depends on $\gamma_{2k\ell-1}$, the \mathbb{R} -rm for $\gamma(z^\pm)$ will generate a K -rm for β as in the proof of Claim 4. By (9), the only moment from β , which could depend on $\gamma_{2k\ell-1}$, is $\beta_{0,2k}$. Note that if $q_{0,2k,2k\ell-1} = 0$, then also $\beta_{0,2k}$ is independent from the value of $\gamma_{2k\ell-1}$. We have that

$$q_{0,2k,2k\ell-1} = \sum_{\substack{0 \leq i_1, \dots, i_{2k} \leq \ell, \\ i_1 + \dots + i_{2k} = 2k\ell - 1}} q_{i_1} q_{i_2} \dots q_{i_{2k}} = 2k(q_\ell)^{2k-1} q_{\ell-1},$$

where in the first equality we used the definition (10) of $q_{0,2k,2k\ell-1}$, while in the second we used the fact that $i_1 + \dots + i_{2k} = 2k\ell - 1$ could be fulfilled only if $2k-1$ indices i_j are ℓ and one is $\ell - 1$. So $q_{0,2k,2k\ell-1} = 0$ iff $q_{\ell-1} = 0$. But this is true by Claim 1 and concludes the proof of Theorem 3.1. ■

To prove Theorem 3.2 for $q(x) = x^\ell$, $\ell \geq 3$, only a little adaptation of the last part of the proof of Theorem 3.1 is needed, which we now explain.

Proof of Theorem 3.2.: The implications (1) \Rightarrow (4) and (2) \Rightarrow (1) are trivial. The implication (4) \Rightarrow (3) follows from [1, Theorem 3.14]. It remains to prove the implication (3) \Rightarrow (2). Following the proof of Theorem 3.1 everything remains the same until (2'). It remains to justify that the upper bound in (23) can be decreased to $m := k\ell - \lceil \frac{\ell}{2} \rceil$. If $\text{rank } A_\gamma \leq m$, then we are already done. From now on we assume that $r := \text{rank } A_\gamma > m$. Since γ admits a \mathbb{R} -rm, which we denote by μ , γ is prg and $\text{rank } \gamma = \text{rank } A_\gamma = \text{rank } A_\gamma(r-1)$ by [44, Theorem 3.9]. Hence, $A_\gamma(r-1)$ is pd and in particular also its submatrix $A_\gamma(m)$ is pd. We denote by $\gamma(z_1, \dots, z_\ell)$ a sequence obtained from the sequence γ by replacing $\gamma_{(2k-1)\ell+1}, \gamma_{(2k-1)\ell+2}, \dots, \gamma_{2k\ell}$ with variables z_1, \dots, z_ℓ . The sequence $\gamma^{(0, (2k-1)\ell)} := (\gamma_0, \dots, \gamma_{(2k-1)\ell})$ is represented by μ , being a subsequence of γ . If ℓ is even, then $(2k-1)\ell$ is also even and by [44, Theorem 3.9], $\gamma^{(0, (2k-1)\ell)}$ has a $(\text{rank } \gamma^{(0, (2k-1)\ell)})$ -atomic \mathbb{R} -rm. Otherwise ℓ is odd, $(2k-1)\ell$ is also odd and by [44, Theorem 3.1], $\gamma^{(0, (2k-1)\ell)}$ has a $(\text{rank } \gamma^{(0, (2k-1)\ell-1)})$ -atomic \mathbb{R} -rm, where $\gamma^{(0, (2k-1)\ell-1)} := (\gamma_0, \dots, \gamma_{(2k-1)\ell-1})$. We denote the measure obtained in this way by μ_1 and generate its moment sequence $\gamma(z_1, \dots, z_\ell)$, where z_1, \dots, z_ℓ are the moments of degrees $(2k-1)\ell+1, \dots, 2k\ell$. Hence, $\text{rank } A_{\gamma(z_1, \dots, z_\ell)}$ is equal to $\text{rank } A_{\gamma^{(0, (2k-1)\ell)}}$ for even ℓ and $\text{rank } A_{\gamma^{(0, (2k-1)\ell-1)}}$ for odd ℓ . Since $(2k-1)\ell = 2m$ for even ℓ and $(2k-1)\ell-1 = 2m$ for odd ℓ , μ_1 is m -atomic (since $A_\gamma(m) \succ 0$ by assumption). If none of the moments $\beta_{i,j}$ of the sequence $\beta^{(2k-1)}$ depends on $\gamma_{(2k-1)\ell+1}, \gamma_{(2k-1)\ell+2}, \dots, \gamma_{2k\ell}$, then μ_1 will generate a K -rm for $\beta^{(2k-1)}$ as in the proof of Claim 5 of Theorem 3.1. But by definition (16), none of the moments of $\beta^{(2k-1)}$ depends on $\gamma_{(2k-1)\ell+1}, \gamma_{(2k-1)\ell+3}, \dots, \gamma_{2k\ell}$, which concludes the proof of Theorem 3.2. ■

A corollary to Theorem 3.1 is an improvement of the bounds on the degrees of sums of squares in the Positivstellensatz [11, Corollary 6.3] for the curves of the form $y = q(x)$, $q \in \mathbb{R}[x]$, $\text{deg } q \geq 3$.

Corollary 3.3: *Let $K := \{(x, y) \in \mathbb{R}^2 : y = q(x)\}$, where $q \in \mathbb{R}[x]$ satisfies $\text{deg } q \geq 3$. Let $k \geq \text{deg } q$. If $r(x, y) \in \mathbb{R}[x, y]_{2k}$ is strictly positive on K , then r admits a decomposition*

$$r(x, y) = \sum_{i=1}^{\ell_1} f_i(x, y)^2 + (y - q(x)) \sum_{i=1}^{\ell_2} g_i(x, y)^2 - (y - q(x)) \sum_{i=1}^{\ell_3} h_i(x, y)^2,$$

where $\ell_1, \ell_2, \ell_3 \in \mathbb{Z}_+$, $f_i, g_i \in \mathbb{R}[x, y]$ and

$$\text{deg } f_i^2 \leq 2m, \text{ deg}((y - q(x))g_i^2) \leq 2m, \text{ deg}((y - q(x))h_i^2) \leq 2m$$

with $m = k + \text{deg } q - 1$.

Proof: By the equivalence (1) \Leftrightarrow (3) of Theorem 3.1, the set K has the property $(R_{k, \text{deg } q-2})$ in the notation of [52, p. 2713]. Now the result follows by [52, Theorem 1.5]. ■

Remark 3.3: The bound on m in Theorem 3.3 from [11, Corollary 5.4] is quadratic in k and $\text{deg } q$, namely $(2k + 1) \text{deg } q$.

3.2. Solution to the parabolic TMP

The following is a concrete solution to the parabolic TMP, first solved in [6]. We give an alternative proof together with a new solvability condition, i.e. (6) below, where the variety condition is removed.

Theorem 3.4 (Solution to the parabolic TMP, even case): *Let*

$$K := \{(x, y) \in \mathbb{R}^2 : y = q_2x^2 + q_1x + q_0\},$$

where $q_0, q_1, q_2 \in \mathbb{R}$ and $q_2 \neq 0$, be the parabola and $\beta := \beta^{(2k)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$, where $k \geq 2$. The following statements are equivalent:

- (1) β has a K -representing measure.
- (2) β has a $(\text{rank } M_k)$ -atomic K -representing measure.
- (3) M_k is positive semidefinite, recursively generated, satisfies the column relation $Y = q_2X^2 + q_1X + q_0$ and $\text{rank } M_k \leq \text{card } \mathcal{V}(\beta)$, where

$$\mathcal{V}(\beta) := \bigcap_{\substack{g \in \mathbb{R}[x,y]_{\leq k}, \\ g(X,Y)=0}} \mathcal{Z}(g).$$

- (4) M_k satisfies $Y = q_2X^2 + q_1X + q_0$ and admits a positive semidefinite, recursively generated extension M_{k+1} .
- (5) M_k satisfies $Y = q_2X^2 + q_1X + q_0$ and admits a positive semidefinite extension M_{k+1} .
- (6) M_k is positive semidefinite, the relations $\beta_{i,j+1} = q_2\beta_{i+2,j} + q_1\beta_{i+1,j} + q_0\beta_{i,j}$ hold for every $i, j \in \mathbb{Z}_+$ with $i + j \leq 2k - 2$ and, defining

$$\mathcal{B} = \{1, x, y, yx, \dots, y^{k-1}, y^{k-1}x, y^k\}, \tag{24}$$

one of the following statements holds:

- (a) $(M_k)|_{\mathcal{B} \setminus \{y^k\}}$ is positive definite.
- (b) $\text{rank}(M_k)|_{\mathcal{B} \setminus \{y^k\}} = \text{rank } M_k$.

Proof: By applying an alt $\phi(x, y) = (x, \frac{1}{q_2}y - q_1x - q_0)$ to the sequence β we get a sequence $\tilde{\beta}$ with the moment matrix $M_k(\tilde{\beta})$ satisfying $Y = X^2$. Using Proposition 2.1, each of the statements (1)–(6) holds for the original sequence β with the column relation $Y = q_2X^2 + q_1X + q_0$ iff it holds for $\tilde{\beta}$ with the column relation $Y = X^2$. So we may assume $(q_2, q_1, q_0) = (1, 0, 0)$. Let us start by proving the equivalences (1) \Leftrightarrow (2) \Leftrightarrow (6). By [57], (1) is equivalent to:

- (1') β has a s -atomic K -representing measure for some $s \in \mathbb{N}$.

Let

$$\gamma := \gamma^{(0,4k)} = (\gamma_0, \gamma_1, \dots, \gamma_{4k}) \in \mathbb{R}^{4k+1}, \tag{25}$$

where $\gamma_t = \beta_{t \bmod 2, \lfloor \frac{t}{2} \rfloor}$, which is a special case of definition (16) in the proof of Theorem 3.1. Claim 5 in the proof of Theorem 3.1 holds with the same proof also for

$q(x) = x^2$. Using Claim 5 and [44, Theorem 3.9] for γ , the equivalences (1') \Leftrightarrow (2) \Leftrightarrow (6) follow by noting that $A_\gamma = (M_k)|_{\mathcal{B}}$ and $A_\gamma(2k - 1) = (M_k)|_{\mathcal{B} \setminus \{Y^k\}}$.

The implications (2) \Rightarrow (4) and (4) \Rightarrow (5) are trivial. The implication (1) \Rightarrow (3) follows from the necessary conditions for the existence of a K -rm (the variety condition follows from [1, Proposition 3.1 and Corollary 3.7]).

Now we prove the implication (5) \Rightarrow (6). By [1, Theorem 3.14], it follows that M_k is rg. Defining the sequence $\tilde{\gamma} := \gamma^{(0,4k+2)} = (\gamma_0, \gamma_1, \dots, \gamma_{4k+1}, \gamma_{4k+2}) \in \mathbb{R}^{4k+3}$, where $\gamma_t = \beta_{t \bmod 2, \lfloor \frac{t}{2} \rfloor}$, which is a special case of definition (16) in the proof of Theorem 3.1, it follows by M_{k+1} being psd that in particular $(M_{k+1})|_{\mathcal{B} \cup \{y^{k+1}\}} = A_{\tilde{\gamma}}$ is also psd. If $A_\gamma(0,4k)$ is pd, then $(M_k)|_{\mathcal{B} \setminus \{y^k\}}$ is pd, which is (a) of Theorem 3.4. Otherwise $A_\gamma(0,4k)$ is singular and prg by [44, Theorem 2.6]. In particular, $\text{rank } A_\gamma = \text{rank } A_\gamma(2k - 1)$, which, by noting that $(M_k)|_{\mathcal{B} \setminus \{y^k\}} = A_\gamma(2k - 1)$, implies (b) of Theorem 3.4. This proves (5) \Rightarrow (6).

It remains to prove the implication (3) \Rightarrow (6). If $(M_k)|_{\mathcal{B} \setminus \{y^k\}}$ is pd, we are done. Otherwise $(M_k)|_{\mathcal{B} \setminus \{y^k\}}$ is not pd. We have to prove that in this case $\text{rank}(M_k)|_{\mathcal{B} \setminus \{y^k\}} = \text{rank } M_k$. We assume by contradiction that $\text{rank}(M_k)|_{\mathcal{B} \setminus \{y^k\}} < \text{rank } M_k$. Let γ be as in (25). The inequality $\text{rank}(M_k)|_{\mathcal{B} \setminus \{y^k\}} < \text{rank } M_k$ implies that $\text{rank } A_\gamma(2k - 1) < \text{rank } A_\gamma$. Let $r = \text{rank } \gamma$. Then, by [44, Theorem 2.6], $\text{rank } A_\gamma(2k - 1) = r$ and hence, [44, Theorems 3.9, 3.10] imply that $\gamma^{(0,4k-2)} = (\gamma_0, \dots, \gamma_{4k-2})$ has a unique r -atomic \mathbb{R} -rm with atoms x_1, \dots, x_r . Hence, $\text{rank } A_\gamma = r + 1$. Note that for every $g(x, y) \in \mathbb{R}[x, y]$, which is a column relation of M_k , it follows that $g(x, x^2) \in \mathbb{R}[x]$ is a column relation of A_γ (where columns of A_γ are $1, X, \dots, X^{2k}$). Since $(x, y) \in \mathcal{V}(\beta)$ and $(x, y') \in \mathcal{V}(\beta)$, implies that $y = y'$ (due to $y = x^2$ and $y' = x^2$), it follows that $\mathcal{V}(\beta) \subseteq \{(x_1, x_1^2), \dots, (x_r, x_r^2)\}$. (This is true, since the atoms of a finitely atomic measure always satisfy all column relations of the moment matrix. Moreover, the sets are equal, but we do not need this in the rest of the proof.) Hence, $|\mathcal{V}(\beta)| \leq r$. Since $\text{rank } A_\gamma = \text{rank } M_k$, this leads to a contradiction with the assumption $\text{rank } M_k \leq |\mathcal{V}(\beta)|$. ■

Remark 3.4: (1) *The proof of the implication (3) \Rightarrow (2) of Theorem 3.4 in [6]:* [6] considers 5 different cases according to the form of the relations between the columns of \mathcal{B} defined by (24). The most demanding cases, which both use the FET as the main tool in the construction of a flat extension M_{k+1} of M_k , are cases where there is only one relation and the column Y^k occurs nontrivially in it or if there is no relation present.

(2) *Complexity of checking conditions in the statements of Theorem 3.4:* Among conditions in (3) the most demanding is the variety condition $\text{rank } M_k \leq \text{card } \mathcal{V}(\beta)$, since it requires solving a system of polynomial equation, which can be numerically difficult and unstable. However, (6) of Theorem 3.4 requires less work and can be stably checked numerically. Namely, one has to check if the column relations $Y^i X^j = q(X)^i X^j, i + j \leq k, i \in \mathbb{N}$, hold, then try to compute the Cholesky decomposition VV^T of $(M_k)|_{\mathcal{B}}$, compute the column echelon form U of V and in case U is of the form $(I_r \ B)^T$ for some matrix B , then (a) or (b) holds and an rm exists. This follows by noting that $(M_k)|_{\mathcal{B}}$ corresponds to the Hankel matrix of a univariate sequence (see the proof above), for which the solution to the TMP is given in Algorithm 2.1.

The following example shows that the variety condition $\text{rank } M_k \leq \text{card } \mathcal{V}(\beta)$ from Theorem 3.4.(3) cannot be removed in contrast to the case of K being a circle [5, Theorem 2.1] or a union of two parallel lines [49, Theorem 3.1].

The *Mathematica* file with numerical computations can be found on the link https://github.com/ZalarA/TMP_quadratic_curves.

Example 3.5: Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 4}$ be a bivariate sequence of degree 4 with the moment matrix M_2 equal to

$$M_2 = \begin{matrix} & & 1 & X & Y & X^2 & XY & Y^2 \\ \begin{matrix} 1 \\ X \\ Y \\ X^2 \\ XY \\ Y^2 \end{matrix} & \left[\begin{array}{cccccc} 3 & 0 & 2 & 2 & 0 & 2 \\ 0 & 2 & 0 & 0 & 2 & 0 \\ 2 & 0 & 2 & 2 & 0 & 2 \\ 2 & 0 & 2 & 2 & 0 & 2 \\ 0 & 2 & 0 & 0 & 2 & 0 \\ 2 & 0 & 2 & 2 & 0 & 3 \end{array} \right. \end{matrix}.$$

M_2 is psd with the eigenvalues $\frac{1}{2}(9 + \sqrt{65}) \approx 8.53, 4, 1, \frac{1}{2}(9 - \sqrt{65}) \approx 0.47, 0, 0$, and the column relations $Y = X^2, XY = X$. Hence, M_2 is psd, rg and satisfies $Y = X^2$. The variety $\mathcal{V}(\beta)$ is equal to $\{(0, 0), (-1, 1), (1, 1)\}$. So $4 = \text{rank } M_2 > \text{card } \mathcal{V}(\beta) = 3$ and the variety condition is not satisfied. Thus β does not admit a representing measure supported on the parabola $y = x^2$. So M_k being psd, satisfying $Y = X^2$ and rg does not imply the variety condition and the existence of a representing measure.

Note that by Remark 3.4.(2) it is cheaper and more stable to check only that $Y = X^2$ is a column relation and then solve the TMP for γ , such that $A_\gamma = (M_2)|_{\mathcal{B}}$, using Algorithm 2.1. Since the case 3.2 applies, an rm does not exist.

The following example demonstrates the solution of [6, Example 1.6] in the univariate setting. The *Mathematica* file with numerical computations can be found on the link https://github.com/ZalarA/TMP_quadratic_curves.

Example 3.6: Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 6}$ be a bivariate sequence of degree 6 with the moment matrix M_3 equal to

$$M_3 = \begin{matrix} & & 1 & X & Y & X^2 & XY & Y^2 & X^3 & X^2Y & XY^2 & Y^3 \\ \begin{matrix} 1 \\ X \\ Y \\ X^2 \\ XY \\ Y^2 \\ X^3 \\ X^2Y \\ XY^2 \\ Y^3 \end{matrix} & \left[\begin{array}{cccccccccc} 1 & 0 & a & a & 0 & b & 0 & b & 0 & c \\ 0 & a & 0 & 0 & b & 0 & b & 0 & c & 0 \\ a & 0 & b & b & 0 & c & 0 & c & 0 & d \\ a & 0 & b & b & 0 & c & 0 & c & 0 & d \\ 0 & b & 0 & 0 & c & 0 & c & 0 & d & 0 \\ b & 0 & c & c & 0 & d & 0 & d & 0 & e \\ 0 & b & 0 & 0 & c & 0 & c & 0 & d & 0 \\ b & 0 & c & c & 0 & d & 0 & d & 0 & e \\ 0 & c & 0 & 0 & d & 0 & d & 0 & e & 0 \\ c & 0 & d & d & 0 & e & 0 & e & 0 & f \end{array} \right. \end{matrix},$$

with the inequalities $a > 0, b > a^2, c > \frac{b^2}{a}, d > \frac{b^3 - 2abc + c^2}{b - a^2}$, which ensure that $(M_2)|_{\{1, X, Y, XY, Y^2, X^2Y\}}$ is psd and $(M_2)|_{\{1, X, Y, XY, Y^2\}}$ is pd. Note that M_3 satisfies the column relations $Y = X^2$,

$XY = X^3$ and $Y^2 = X^2Y$. We introduce the univariate sequence

$$\gamma \in (1, 0, a, 0, b, 0, c, 0, d, 0, e, 0, f) \in \mathbb{R}^{13}$$

as in the proof of Theorem 3.4. We denote the rows and columns of A_γ by $1, X, \dots, X^6$. Since $(M_2)_{\{1, X, Y, XY, Y^2\}}$ is pd, it follows that $A_\gamma(4)$ is pd. For

$$e = (\mathbf{v}_\gamma(5, 4))^T (A_{(1,0,a,0,b,0,c,0,d)})^{-1} \mathbf{v}_\gamma(5, 4) = \frac{-c^3 + 2bcd - ad^2}{b^2 - ac},$$

we have that $A_\gamma(5) \succeq 0$ (e.g. using [58, Theorem 1] for $A_\gamma(5)$) and $X^5 \in \text{span}\{1, X, \dots, X^4\}$ in $A_\gamma(5)$, where the vector $\mathbf{v}_\gamma(5, 4) = (0 \ c \ 0 \ d \ 0)^T$ is the restriction of the column X^5 to the rows indexed by $1, X, X^2, X^3, X^4$. Hence, for γ to admit a \mathbb{R} -rm, $A_\gamma \succeq 0$ and $X^i \in \text{span}\{1, X, \dots, X^{i-1}\}$ for $i = 5, 6$ [44, Theorem 3.9]. Since $A_\gamma(5) \succeq 0$ and the last column of $A_\gamma(5)$ is a linear combination of the others, it only needs to hold by [58, Theorem 1], that

$$\begin{aligned} \mathbf{v}_\gamma(6, 5) \in \mathcal{C}(A_\gamma(5)) \quad \text{and} \quad f &= (\mathbf{v}_\gamma(6, 5))^T (A_\gamma(5))^\dagger \mathbf{v}_\gamma(6, 5) \\ &= \frac{-bc^4 - b^2c^2d - 2ac^3d - b^3d - b^3d^2 + 4abcd^2 - a^2d^2}{(b^2 - ac)^2}, \end{aligned}$$

where $\mathbf{v}_\gamma(6, 5)$ denotes the restriction of X^6 to the rows indexed by $1, \dots, X^5$ in A_γ and $(A_\gamma(5))^\dagger$ denotes the Moore–Penrose inverse of $A_\gamma(5)$. Using *Mathematica* we check that the equality $A_\gamma(5)(A_\gamma(5))^\dagger \mathbf{v}_\gamma(6, 5) = \mathbf{v}_\gamma(6, 5)$ holds, which implies that $\mathbf{v}_\gamma(6, 5) \in \mathcal{C}(A_\gamma(5))$ is true. By [44, Theorem 3.10], in this case the \mathbb{R} -rm is unique, 5-atomic and consists of the roots of the polynomial

$$\begin{aligned} p(x) &= (1 \ x \ x^2 \ x^3 \ x^4 \ x^5) (A_{(1,0,a,0,b,0,c,0,d)})^{-1} \mathbf{v}_\gamma(5, 4) \\ &= x \left(x^4 + \frac{ad - bc}{b^2 - ac} x^2 + \frac{c^2 - bd}{b^2 - ac} \right). \end{aligned}$$

Since $p(x)$ has roots $0, x_1, -x_1, x_2, -x_2$ and the atoms for the K -rm for β are $(0, 0), (x_1, x_1^2), (-x_1, x_1^2), (x_2, x_2^2), (-x_2, x_2^2)$.

The following theorem is a concrete solution to the parabolic TMP of odd degree, which is solved using the same technique as in the proof of Theorem 3.2, but here we get explicit conditions for the existence of the solution, similarly as in the even degree case.

Theorem 3.7 (Solution to the parabolic TMP, odd case): *Let $K := \{(x, y) \in \mathbb{R}^2 : y = x^2\}$ be the parabola and $\beta := \beta^{(2k-1)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k-1}$, where $k \geq 2$. Let $\gamma := (\gamma_0, \gamma_1, \dots, \gamma_{4k-2})$ be a sequence, defined by $\gamma_t := \beta_{t \bmod 2, \lfloor \frac{t}{2} \rfloor}$ for $t = 0, 1, \dots, 4k - 2$. The following statements are equivalent:*

- (1) β has a K -representing measure.
- (2) β has a $(\text{rank } \gamma)$ -atomic K -representing measure.

- (3) β can be extended to a sequence $\beta^{(2k)}$ such that M_k is psd, rg, has a column relation $Y = X^2$ and satisfies $\text{rank } M_k \leq \text{card } \mathcal{V}(\beta^{(2k)})$, where

$$\mathcal{V}(\beta^{(2k)}) := \bigcap_{\substack{g \in \mathbb{R}[x,y]_{\leq k}, \\ g(X,Y)=\mathbf{0} \text{ in } M_k}} \mathcal{Z}(g).$$

- (4) β can be extended to a sequence $\beta^{(2k+2)}$ such that M_{k+1} is psd and has a column relation $Y = X^2$.
 (5) The relations $\beta_{i,j+1} = \beta_{i+2,j}$ hold for every $i, j \in \mathbb{Z}_+$ with $i + j \leq 2k - 1$, $A_\gamma \geq 0$ and the sequence γ is positively recursively generated.
 (6) The relations $\beta_{i,j+1} = \beta_{i+2,j}$ hold for every $i, j \in \mathbb{Z}_+$ with $i + j \leq 2k - 1$ and defining $\beta_{i,2k-i} = \beta_{i \bmod 2, 2k-i + \lfloor \frac{i}{2} \rfloor}$ for $2 \leq i \leq 2k$, the moment matrix $(M_k)|_{\mathcal{B} \setminus \{y^k\}}$, where $\mathcal{B} = \{1, x, y, yx, \dots, y^{k-1}, y^{k-1}x, y^k\}$, is positive semidefinite and

$$(\beta_{0,k} \ \beta_{1,k} \ \beta_{0,k+1} \ \beta_{1,k+1} \ \cdots \ \beta_{0,2k-1})^T \in \mathcal{C}((M_k)|_{\mathcal{B} \setminus \{xy^{k-1}, y^k\}, \mathcal{B} \setminus \{y^k\}}). \quad (26)$$

Proof: The equivalences (1) \Leftrightarrow (3) \Leftrightarrow (4) follow by Theorem 3.4. By [57], (1) is equivalent to:

- (1') β has an s -atomic K -rm for some $s \in \mathbb{N}$.

Claim 5 of Theorem 3.1 holds with the same proof also for $q(x) = x^2$ and odd degree sequence (i.e. $i + j \leq 2k - 1$). Together with [44, Theorem 3.9], the equivalences (1') \Leftrightarrow (2) \Leftrightarrow (5) follow. Note that $(M_k)|_{\mathcal{B} \setminus \{y^k\}} = A_\gamma$. By [39, Theorem 2.7.5], γ is prg if and only if (26) holds. This establishes the equivalence (5) \Leftrightarrow (6). ■

- Remark 3.5:** (1) Note that $\text{rank } \gamma$ in Theorem 3.7 is at most $2k$ and it is $2k$ iff A_γ is positive definite.
 (2) Theorem 3.7 also solves the odd degree TMP on any curve of the form $y = q_2x^2 + q_1x + q_0$, where $q_0, q_1, q_2 \in \mathbb{R}$ and $q_2 \neq 0$. As in the proof of the even degree case one applies ϕ from the proof of Theorem 3.4 to β to come into the case $y = x^2$ and then use Theorem 3.7.

3.3. A solution to the TMP based on a feasibility of a linear matrix inequality

In this section, we give another alternative solution to the TMP on curves $y = q(x)$, where $q(x) \in \mathbb{R}[x]$ and $\text{deg } q \geq 3$, which is based on a feasibility of a linear matrix inequality associated to the univariate sequence γ , obtained from the original sequence β as in the proofs of the results of previous subsections. The feasibility question appears as a result of the fact that γ is not fully determined by β , but β admits a K -rm if and only if γ can be completed to a sequence admitting an \mathbb{R} -rm.

For $n \in \mathbb{N}$, we denote by $[n]$ the set of all nonnegative integers smaller or equal to n . Let the sets N_1, N_2 form a partition of $[n]$, i.e. $N_1, N_2 \subseteq [n]$, $N_1 \cup N_2 = [n]$ and $N_1 \cap N_2 = \emptyset$. Let $\Gamma_1 := (\gamma_t)_{t \in N_1}$ be a sequence of real numbers indexed by integers from N_1 and $\Gamma_2 :=$

$(\gamma_t)_{t \in N_2}$ a tuple of variables indexed by integers from N_2 . Let

$$F_{\Gamma_1}(\Gamma_2) : \mathbb{R}^{|N_2|} \rightarrow \mathbb{R}^{|N_1|+|N_2|} \tag{27}$$

be a function with the output a sequence $(\tilde{\gamma}_t)_{t \in [n]}$ where $\tilde{\gamma}_t = \begin{cases} \gamma_t, & \text{if } t \in N_1, \\ \boldsymbol{\gamma}_t, & \text{if } t \in N_2. \end{cases}$

In Theorem 3.8, the set N_1 will be the set of indices, for which the corresponding univariate sequence γ is determined by β , while the indices of the non-determined part will belong to N_2 . Since we can either get an odd or an even sequence, for which the solutions to the \mathbb{R} -TMP are slightly different, we separate two cases for $N_1 \cup N_2 = [n]$ (see (29)).

Theorem 3.8: Let $K := \{(x, y) \in \mathbb{R}^2 : y = q(x)\}$, where $q(x) = \sum_{i=0}^{\ell} q_i x^i \in \mathbb{R}[x]$, $\ell \geq 3$, $q_\ell \neq 0$, and

$$\beta := \beta^{(d)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq d},$$

where $\lceil \frac{d}{2} \rceil \geq \deg q$. For $i, j, s \in \mathbb{Z}_+$, such that $i + j \leq d$, define real numbers

$$q_{i,j,s} := \begin{cases} \sum_{\substack{0 \leq i_1, \dots, i_j \leq \ell, \\ i_1 + \dots + i_j = s-i}} q_{i_1} q_{i_2} \dots q_{i_j}, & \text{if } i \leq s \leq i + j\ell, \\ 0, & \text{otherwise.} \end{cases}$$

Let

$$N_1 := \left\{ t \in \mathbb{Z}_+ : t \bmod \ell + \left\lfloor \frac{t}{\ell} \right\rfloor \leq d \right\},$$

$$\gamma_t = \frac{1}{(q_\ell)^{\lfloor \frac{t}{\ell} \rfloor}} \left(\beta_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor} - \sum_{s=0}^{t-1} q_{t \bmod \ell, \lfloor \frac{t}{\ell} \rfloor, s} \cdot \gamma_s \right) \text{ for every } t \in N_1, \tag{28}$$

and $\Gamma_1 := (\gamma_t)_{t \in N_1}$. Let

$$[n] := \begin{cases} [d\ell + 2], & \text{if } d\ell \text{ is even,} \\ [d\ell + 1], & \text{if } d\ell \text{ is odd,} \end{cases} \tag{29}$$

$\Gamma_2 := (\boldsymbol{\gamma}_t)_{t \in N_2}$ be a tuple of variables with $N_2 = [n] \setminus N_1$ and $F_{\Gamma_1}(\Gamma_2)$ be defined as in (27). Then the following statements are equivalent:

- (1) β has a K -representing measure.
- (2) $\beta_{i,j} = \sum_{p=0}^{\ell} q_p \beta_{i+p,j-1}$ for every $i, j \in \mathbb{Z}_+$, such that $i + j \leq d - \ell + 1$ and there exists a tuple $\Gamma_2 = (\boldsymbol{\gamma}_t)_{t \in N_2} \in \mathbb{R}^{N_2}$ such that $A_{F_{\Gamma_1}(\Gamma_2)} \succeq 0$.

Proof: Observing the proof of Theorem 3.1 for a general $q(x)$ one can note that $F_{\Gamma_1}(\Gamma_2)$ corresponds to the sequence $\tilde{\gamma}$. The original sequence β determines only γ_t for $t \in N_1$ by (28), while for $t \in N_2$, $\boldsymbol{\gamma}_t$ are variables. By the proof of Theorem 3.1, β will have a K -rm iff it satisfies the rg relations coming from the column relation $Y = q(X)$ and there exists $\tilde{\gamma}$ such that $A_{\tilde{\gamma}} \succeq 0$. This proves Theorem 3.8 for even d .

Observing the proof of Theorem 3.2 in case d is odd one can notice that only $\gamma^{(0, d\ell)} = (\gamma_0, \dots, \gamma_{d\ell})$ needs to have a \mathbb{R} -rm to obtain a K -rm for β . In case $d\ell$ is even, this is by [44,

Theorem 3.9] equivalent to $A_{\gamma^{(0, d\ell+2)}} \geq 0$, where $\gamma^{(0, d\ell+2)} = (\gamma_0, \dots, \gamma_{d\ell}, \gamma_{d\ell+1}, \gamma_{d\ell+2})$ for some $\gamma_{d\ell+1}, \gamma_{d\ell+2}$. Since $\gamma^{(0, d\ell+2)}$ corresponds to the sequence $F_{\Gamma_1}(\Gamma_2)$, this proves Theorem 3.8 for even $d\ell$ with d being odd. If $d\ell$ is odd, then by [44, Theorem 3.1] it suffices that there is $\gamma_{d\ell+1}$ such that $A_{\gamma^{(0, d\ell+1)}} \geq 0$, where $\gamma^{(0, d\ell+1)} = (\gamma_0, \dots, \gamma_{d\ell}, \gamma_{d\ell+1})$, and this proves Theorem 3.8 for odd $d\ell$. ■

We will present the statement of Theorem 3.8 on a few examples. The following example is for the case $\deg q = 3$ and a sequence β of even degree.

Example 3.9: Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$ be a bivariate sequence of degree $2k, k \geq 3$, and $K := \{(x, y) \in \mathbb{R}^2 : y = x^3\}$. For the existence of a K -rm β must satisfy the relations $\beta_{i,j} = \beta_{i+3,j-1}$ for every $i, j \in \mathbb{Z}_+$ such that $i + j + 2 \leq 2k$. In the notation of Theorem 3.8 we have

$$q_{i,j,s} := \begin{cases} 1, & \text{if } s = i + 3j, \\ 0, & \text{otherwise,} \end{cases} \quad \text{for } i, j, s \in \mathbb{Z}_+, \text{ such that } i + j \leq 2k,$$

$$N_1 := \left\{ t \in \mathbb{Z}_+ : t \bmod 3 + \left\lfloor \frac{t}{3} \right\rfloor \leq 2k \right\} = \{t \in \mathbb{Z}_+ : t \leq 6k, t \neq 6k - 1\},$$

$$[n] := [6k + 2] \quad \text{and} \quad N_2 := \{6k - 1, 6k + 1, 6k + 2\}.$$

The formula (28) is equal to

$$\gamma_t = \beta_{t \bmod 3, \lfloor \frac{t}{3} \rfloor} \quad \text{for every } t \in N_1$$

and the function $F_{\Gamma_1} : \mathbb{R}^3 \rightarrow \mathbb{R}^{6k+3}$ is defined by

$$F_{\Gamma_1}(\Gamma_2) = F_{\Gamma_1}(\boldsymbol{\gamma}_{6k-1}, \boldsymbol{\gamma}_{6k+1}, \boldsymbol{\gamma}_{6k+2}) := (\gamma_0, \gamma_1, \dots, \gamma_{6k-2}, \boldsymbol{\gamma}_{6k-1}, \gamma_{6k}, \boldsymbol{\gamma}_{6k+1}, \boldsymbol{\gamma}_{6k+2}).$$

The matrix $A_{F_{\Gamma_1}(\Gamma_2)}$ is equal to

$$\begin{pmatrix} \gamma_0 & \gamma_1 & \gamma_2 & \gamma_3 & \cdots & \cdots & & & \gamma_{3k} & \gamma_{3k+1} \\ \gamma_1 & \gamma_2 & \gamma_3 & \ddots & & & & & & \vdots \\ \gamma_2 & \gamma_3 & \ddots & & & & & & & \gamma_{6k-2} \\ \gamma_3 & \ddots & & & & & & & & \boldsymbol{\gamma}_{6k-1} \\ \vdots & & & & & & & & & \gamma_{6k} \\ \gamma_{3k} & \cdots & \cdots & \cdots & \cdots & \gamma_{6k-2} & \boldsymbol{\gamma}_{6k-1} & \gamma_{6k} & \boldsymbol{\gamma}_{6k+1} & \boldsymbol{\gamma}_{6k+2} \\ \gamma_{3k+1} & \cdots & \cdots & \cdots & \gamma_{6k-2} & \boldsymbol{\gamma}_{6k-1} & \gamma_{6k} & \boldsymbol{\gamma}_{6k+1} & \boldsymbol{\gamma}_{6k+2} & \end{pmatrix},$$

The question of feasibility of $A_{F_{\Gamma_1}(\Gamma_2)} \geq 0$ can be answered analytically, since the structure of the missing entries is simple enough. Actually it is even easier to work with $A_{(\gamma_0, \dots, \boldsymbol{\gamma}_{6k-1}, \gamma_{6k})}$ and answer the feasibility question together with the condition from the solution of [44, Theorem 3.9] (see [48, Theorem 3.1]).

Remark 3.6: If $\deg q = 3$ in Theorem 3.1, then a polynomial is of the form $y = q_3x^3 + q_2x^2 + q_1x + q_0 \in \mathbb{R}[x]$, where $q_3 \neq 0$, and using alts it can be transformed to $y = x^3$.

Indeed, by first applying an alt as at the beginning of the proof of Claim 1 of Theorem 3.1, we can assume that $q_2 = 0$, i.e. the polynomial becomes $y = q_3x^3 + q_1x + q_0$. Now we apply an alt $(x, y) \mapsto (x, y - q_1x - q_0)$, followed by $(x, y) \mapsto (\sqrt[3]{q_3}x, y)$ and get a polynomial $y = x^3$.

The following example demonstrates the statement of Theorem 3.8 for the case $\deg q = 3$ and a sequence β of odd degree.

Example 3.10: Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k-1}$ be a bivariate sequence of degree $2k-1, k \geq 3$, and $K := \{(x, y) \in \mathbb{R}^2 : y = x^3\}$. For the existence of a K -rm β must satisfy the relations $\beta_{i,j} = \beta_{i+3,j-1}$ for every $i, j \in \mathbb{Z}_+$ such that $i + j + 2 \leq 2k - 1$. In the notation of Theorem 3.8, we have

$$q_{i,j,s} := \begin{cases} 1, & \text{if } s = i + 3j, \\ 0, & \text{otherwise.} \end{cases} \quad \text{for } i, j, s \in \mathbb{Z}_+, \text{ such that } i + j \leq 2k - 1,$$

$$N_1 := \left\{ t \in \mathbb{Z}_+ : t \bmod 3 + \left\lfloor \frac{t}{3} \right\rfloor \leq 2k - 1 \right\} = \{t \in \mathbb{Z}_+ : t \leq 6k - 3 \text{ and } t \neq 6k - 4\},$$

$$[n] = [6k - 2] \quad \text{and} \quad N_2 := \{6k - 4, 6k - 2\}.$$

The formula (28) is equal to

$$\gamma_t = \beta_{t \bmod 3, \lfloor \frac{t}{3} \rfloor} \quad \text{for every } t \in \mathbb{N},$$

and the function $F_{\Gamma_1} : \mathbb{R}^2 \rightarrow \mathbb{R}^{6k-1}$ is defined by

$$F_{\Gamma_1}(\Gamma_2) = F_{\Gamma_1}(\mathcal{Y}_{6k-4}, \mathcal{Y}_{6k-2}) := (\gamma_0, \gamma_1, \dots, \gamma_{6k-5}, \mathcal{Y}_{6k-4}, \gamma_{6k-3}, \mathcal{Y}_{6k-2}).$$

Since $6k - 3 = (2k - 1) \cdot 3$ is odd, only feasibility of the inequality $A_{F_{\Gamma_1}(\Gamma_2)} \geq 0$ is important for the existence of the rm for β , where $A_{F_{\Gamma_1}(\Gamma_2)}$ is equal to

$$\begin{pmatrix} \gamma_0 & \gamma_1 & \gamma_2 & \gamma_3 & \cdots & \cdots & \gamma_{3k-1} \\ \gamma_1 & \gamma_2 & \gamma_3 & \ddots & & & \vdots \\ \gamma_2 & \gamma_3 & \ddots & & & & \gamma_{6k-5} \\ \gamma_3 & \ddots & & & & \gamma_{6k-5} & \mathcal{Y}_{6k-4} \\ \vdots & & & & \gamma_{6k-5} & \mathcal{Y}_{6k-4} & \gamma_{6k-3} \\ \gamma_{3k-1} & \cdots & \cdots & \gamma_{6k-5} & \mathcal{Y}_{6k-4} & \gamma_{6k-3} & \mathcal{Y}_{6k-2} \end{pmatrix}.$$

This feasibility question can be answered analytically, since the structure of missing entries is simple enough. See Theorem 3.13.

The following example demonstrates the statement of Theorem 3.8 for the case $y = x^4$ and a sequence β of even degree.

Example 3.11: Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$ be a bivariate sequence of degree $2k, k \geq 4$, and $K := \{(x, y) \in \mathbb{R}^2 : y = x^4\}$. For the existence of a K -rm β must satisfy the relations $\beta_{i,j} =$

$$\begin{aligned}
 N_1 &:= \left\{ t \in \mathbb{Z}_+ : t \bmod 4 + \left\lfloor \frac{t}{4} \right\rfloor \leq 2k - 1 \right\} \\
 &= \{ t \in \mathbb{Z}_+ : t \leq 8k - 4 \text{ and } t \notin \{8k - 9, 8k - 6, 8k - 5\}\}, \\
 n &:= [8k - 2] \quad \text{and} \quad N_2 := \{8k - 9, 8k - 6, 8k - 5, 8k - 3, 8k - 2\}.
 \end{aligned}$$

The formula (28) is equal to

$$\gamma_t = \beta_{t \bmod t, \lfloor \frac{t}{4} \rfloor} \quad \text{for every } t \in \mathbb{N},$$

the function $F_{\Gamma_1} : \mathbb{R}^5 \rightarrow \mathbb{R}^{8k-1}$ is defined by

$$\begin{aligned}
 F_{\Gamma_1}(\Gamma_2) &= F(\gamma_{8k-9}, \gamma_{8k-6}, \gamma_{8k-5}, \gamma_{8k-3}, \gamma_{8k-2}) \\
 &:= (\gamma_0, \gamma_1, \dots, \gamma_{8k-10}, \gamma_{8k-9}, \gamma_{8k-8}, \gamma_{8k-7}, \gamma_{8k-6}, \gamma_{8k-5}, \gamma_{8k-4}, \gamma_{8k-3}, \gamma_{8k-2}),
 \end{aligned}$$

and the matrix $A_{F_{\Gamma_1}(\Gamma_2)}$ is equal to

$$\left(\begin{array}{cccccccc|cccc}
 \gamma_0 & \gamma_1 & \gamma_2 & \gamma_3 & \cdots & & & & \cdots & \gamma_{4k-2} & & \gamma_{4k-1} \\
 \gamma_1 & \gamma_2 & \gamma_3 & \ddots & & & & & & & & \vdots \\
 \gamma_2 & \gamma_3 & \ddots & & & & & & & & & \gamma_{8k-8} \\
 \gamma_3 & \ddots & & & & & & & & \gamma_{8k-10} & \gamma_{8k-9} & \gamma_{8k-8} & \gamma_{8k-7} \\
 \vdots & & & & & & & & & & & & \gamma_{8k-6} \\
 & & & & & & \gamma_{8k-10} & \gamma_{8k-9} & \gamma_{8k-8} & \gamma_{8k-7} & \gamma_{8k-6} & \gamma_{8k-5} & \gamma_{8k-4} \\
 & & & & & & \gamma_{8k-10} & \gamma_{8k-9} & \gamma_{8k-8} & \gamma_{8k-7} & \gamma_{8k-6} & \gamma_{8k-5} & \gamma_{8k-4} \\
 & & & & & & & & & & & & \gamma_{8k-3} \\
 \gamma_{4k-2} & & & & & & \gamma_{8k-10} & \gamma_{8k-9} & \gamma_{8k-8} & \gamma_{8k-7} & \gamma_{8k-6} & \gamma_{8k-5} & \gamma_{8k-4} \\
 \gamma_{4k-1} & \cdots & \gamma_{8k-9} & \gamma_{8k-8} & \gamma_{8k-7} & \gamma_{8k-6} & \gamma_{8k-5} & \gamma_{8k-4} & \gamma_{8k-3} & \gamma_{8k-2} & & & \gamma_{8k-2}
 \end{array} \right)$$

Since $8k - 4 = (2k - 1) \cdot 4$ is even, the problem has the same structure as in the even degree case (see Example 3.11).

3.4. A solution to the odd degree TMP on $y = x^3$

The following theorem is a concrete solution to the TMP of odd degree on the curve $y = x^3$, which can be solved using the same technique as odd cases of the TMP on $y = x^\ell$, $\ell \geq 3$. However, for $\ell = 3$ we get explicit conditions for the existence of the solution, similarly as in the even degree case [48, Theorem 3.1].

Theorem 3.13 (Solution to the TMP on $y = x^3$, odd case): *Let $K := \{(x, y) \in \mathbb{R}^2 : y = x^3\}$ and $\beta := \beta^{(2k-1)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k-1}$, where $k \geq 3$. Let $\gamma(z) := (\gamma_0, \gamma_1, \dots, \gamma_{6k-5}, \gamma_{6k-3})$ be a sequence, defined by $\gamma_t := \beta_{t \bmod 3, \lfloor \frac{t}{3} \rfloor}$ for $t = 0, 1, \dots, 6k - 5, 6k - 3$, and z is a variable. The following statements are equivalent:*

- (1) β has a K -representing measure.
- (2) β has a $(\text{rank } M_{k-1})$ -atomic or $(\text{rank } M_{k-1} + 1)$ -atomic K -representing measure.

- (3) The relations $\beta_{i,j+1} = \beta_{i+3,j}$ hold for every $i, j \in \mathbb{Z}_+$ with $i + j \leq 2k - 4$ and denoting $\mathcal{B} = \{1, x, x^2, y, yx, yx^2, \dots, y^{k-1}\}$, one of the following holds:
- (a) $(M_{k-1})|_{\mathcal{B}} > 0$.
 - (b) $(M_{k-1})|_{\mathcal{B}} \not\geq 0, (M_{k-1})|_{\mathcal{B}} \geq 0$, denoting $\gamma := (\gamma_0, \gamma_1, \dots, \gamma_{6k-6}), r := \text{rank } \gamma$ and

$$(\varphi_0 \quad \dots \quad \varphi_{r-1}) := A_{\gamma}(r - 1)^{-1} (\gamma_r \quad \dots \quad \gamma_{2r-1})^T, \tag{30}$$

it holds that

$$\gamma_{6k-u} = \sum_{i=0}^{r-1} \varphi_i \gamma_{6k-u-r+i} \quad \text{for } u = 3, 5, \tag{31}$$

where γ_{6k-4} is defined by (31) for $u = 4$.

Moreover, if a K -representing measure exists, then there does not exist a $(\text{rank } M_{k-1})$ -atomic one if and only if $(M_{k-1})|_{\mathcal{B}} > 0$ and γ_{6k-3} does not satisfy (31) for $u = 3$, where γ_{6k-4} is obtained by (31) for $u = 4$ and one uses (30) with $r = 3k-2$.

Proof: By Theorem 3.8, (1) is equivalent to the validity of the relations $\beta_{i,j+1} = \beta_{i+3,j}$ for every $i, j \in \mathbb{Z}_+$ with $i + j \leq 2k - 4$ and feasibility of $A_{F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})} \geq 0$, where the linear matrix function $A_{F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})}$ is as in Example 3.10. The latter is further equivalent to the existence of γ_{6k-4} and γ_{6k-2} such that $F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})$ has an \mathbb{R} -rm. Here we note that if $A_{F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})} \geq 0$ is such that $\text{rank } F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2}) < A_{F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})}$, then $\text{rank } F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2}) = A_{F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})} - 1$ by [44, Corollary 2.5]. Since γ_{6k-2} occurs only in the bottom right corner of $A_{F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})}$, we can replace it with $\tilde{\gamma}_{6k-2}$ such that $A_{F_{\Gamma_1}(\gamma_{6k-4}, \tilde{\gamma}_{6k-2})} \geq 0$ and $\text{rank } F_{\Gamma_1}(\gamma_{6k-4}, \tilde{\gamma}_{6k-2}) = A_{F_{\Gamma_1}(\gamma_{6k-4}, \tilde{\gamma}_{6k-2})}$, which by [44, Theorem 3.9] indeed implies the existence of an \mathbb{R} -rm. We have that $(M_{k-1})|_{\mathcal{B}} = A_{\gamma}$. If $(M_{k-1})|_{\mathcal{B}} > 0$, there exists γ_{6k-4} such that $A_{(\gamma_0, \gamma_1, \dots, \gamma_{6k-4})} > 0$ and by [44, Theorem 3.1], the sequence $(\gamma_0, \gamma_1, \dots, \gamma_{6k-3})$ has a $(3k - 1)$ -atomic \mathbb{R} -representing measure. Hence, one also finds γ_{6k-2} such that $F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})$ has an \mathbb{R} -rm. If $(M_{k-1})|_{\mathcal{B}} \geq 0$ and $(M_{k-1})|_{\mathcal{B}} \not\geq 0$, then by [44, Theorem 3.8], $(\gamma_0, \gamma_1, \dots, \gamma_{6k-5})$ has a unique \mathbb{R} -rm. This measure also represents γ_{6k-3} iff (31) for $u = 4$ and $u = 3$ holds. This establishes the equivalence (1) \Leftrightarrow (3). The equivalence of both with (2) follows by observing that $F_{\Gamma_1}(\gamma_{6k-4}, \gamma_{6k-2})$ admits a $(\text{rank } \gamma)$ -atomic or $(\text{rank } \gamma + 1)$ -atomic \mathbb{R} -rm. The first case happens iff (b) holds or (a) holds and γ_{6k-4} is obtained by (31) for $u = 4$, where one uses (30) with $r = 3k-2$, and γ_{6k-3} is obtained by (31) for $u = 3$. Since $\text{rank } A_{\gamma} = \text{rank } M_{k-1}$, the equivalence follows. ■

4. The TMP on the curves $yx^{\ell} = 1$

In this section, we study the K -TMP for K being a curve of the form $yx^{\ell} = 1, \ell \in \mathbb{N}, \ell \geq 2$. In Section 4.1, we first give a solution to the K -TMP, based on the size of positive semidefinite extensions of the moment matrix needed and also bound the number of atoms in the K -rm with the smallest number of atoms (see Theorem 4.1 for the even degree and Theorem 4.2 for the odd degree sequences). As a result we obtain a sum-of-squares representation for polynomials, which are strictly positive on K (see Corollary 4.3). This improves bounds in the previously known result [11, Proposition 6.4]. In Section 4.2, we

give a solution to the K -TMP, based on a feasibility of the corresponding linear matrix inequality (see Theorem 4.4).

4.1. Solution to the TMP in terms of psd extensions of M_k , bounds on the number of atoms in the minimal measure and a positivstellensatz

Theorem 4.1 (Even case): Let $K := \{(x, y) \in \mathbb{R}^2 : yx^\ell = 1\}$, where $\ell \in \mathbb{N} \setminus \{1\}$, and $\beta := \beta^{(2k)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$, where $k \geq \ell + 1$. The following statements are equivalent:

- (1) β has a K -representing measure.
- (2) β has a s -atomic K -representing measure for some s satisfying

$$\text{rank } M_k \leq s \leq k(\ell + 1).$$

- (3) M_k satisfies $YX^\ell = 1$ and admits a positive semidefinite, recursively generated extension $M_{k+\ell}$.
- (4) M_k satisfies $YX^\ell = 1$ and admits a positive semidefinite extension $M_{k+\ell+1}$.

Theorem 4.2 (Odd case): Let $K := \{(x, y) \in \mathbb{R}^2 : yx^\ell = 1\}$, where $\ell \in \mathbb{N} \setminus \{1\}$, and $\beta := \beta^{(2k-1)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k-1}$, where $k \geq \ell + 1$. The following statements are equivalent:

- (1) β has a K -representing measure.
- (2) β has a s -atomic K -representing measure for some s satisfying

$$\text{rank } M_k \leq s \leq k(\ell + 1) - \left\lfloor \frac{\ell}{2} \right\rfloor + 1.$$

- (3) $\beta^{(2k-1)}$ can be extended to a sequence $\beta^{(2k)}$ such that M_k satisfies $YX^\ell = 1$ and admits a positive semidefinite, recursively generated extension $M_{k+\ell}$.
- (4) $\beta^{(2k-1)}$ can be extended to a sequence $\beta^{(2k)}$ such that M_k satisfies $YX^\ell = 1$ and admits a positive semidefinite extension $M_{k+\ell+1}$.

Remark 4.1: (1) *Previous bounds on the size of extensions in (4) of Theorem 4.1:* In [11, Section 6], the author considered TMPs on $\mathcal{Z}(p)$ in terms of the size of psd extensions of the moment matrix also for polynomials of the form $p(x, y) = yq(x)$, where $q \in \mathbb{R}[x]$. Namely, by [11, Propositions 6.1, 6.4], a sequence of degree $2k$ admits a $\mathcal{Z}(p)$ -rm, if M_k admits a psd extensions M_{k+r} , where $r = (2k + 2)(2 + \text{deg } q) - (1 + \text{deg } q + k)$. The proof of this result relies on the truncated Riesz–Haviland theorem [52, Theorem 1.2] and a sum-of-squares representations for polynomials, strictly positive on $\mathcal{Z}(p)$ ([11, Proposition 6.4] and [53, Proposition 5.1]). Part (4) of Theorem 4.1 improves Fialow’s result in case $q(x) = x^\ell$, $\ell \geq 2$, by decreasing the size of the extensions to $r = \ell + 1$.

- (2) *Known bounds on the number of atoms in (2) of Theorems 4.1, 4.2:* Similarly as in Remark 3.1.(3.1), part (2) of Theorem 4.1 is a counterpart of [50, Corollary 7.6] for even degree sequences on curves $\mathcal{Z}(yx^\ell - 1)$, while part (2) of Theorem 4.2 improves [50, Corollary 7.6] for curves $\mathcal{Z}(yx^\ell - 1)$ by decreasing it for $\lfloor \frac{\ell}{2} \rfloor - 1$.
- (3) *Uniqueness and description of all solutions in Theorems 4.1 and 4.2:* The same comment as for Remark 3.1.(5) applies here. Beyond the case $yx^2 = 1$ (see Example 4.5) the

structure of the missing entries of A_γ from the proof of Theorem 4.1 is too complicated to have control over all psd completions and consequently over the uniqueness and the description of all solutions.

- (4) *Complexity of checking conditions in (4) of Theorems 4.1 and 4.2:* The same comment as for Remark 3.1.(6) applies here. The main complexity question is the SDP feasibility question, which is cheaper when dealt with on a univariate sequence $A_{\tilde{\gamma}}$ defined in the proof. A precise SDP is stated in Theorem 4.4.

Proof of Theorem 4.1.: The implications (1) \Rightarrow (4) and (2) \Rightarrow (1) are trivial. The implication (4) \Rightarrow (3) follows by [1, Theorem 3.14]. It remains to prove the implication (3) \Rightarrow (2). Assume that YX^ℓ is a column relation and M_k admits a psd, rg extension $M_{k+\ell}$. Let

$$\mathcal{B} = \left\{ y^{k+1}x^{\ell-1}, y^k, y^k x, \dots, y^k x^{\ell-1}, \dots, y, yx, \dots, yx^{\ell-1}, 1, x, \dots, x^{k+1} \right\} \quad (32)$$

be the set of monomials and V the vector subspace in $\mathbb{R}[x, y]_{k+\ell}$ generated by the set \mathcal{B} . Since $M_{k+\ell}$ satisfies

$$X^i Y^j = \begin{cases} X^{i \bmod \ell} Y^{j - \lfloor \frac{j}{\ell} \rfloor}, & \text{if } i, j \in \mathbb{Z}_+, i + j \leq k, j \geq \lfloor \frac{j}{\ell} \rfloor, \\ X^{i-j}, & \text{if } i, j \in \mathbb{Z}_+, i + j \leq k, j < \lfloor \frac{j}{\ell} \rfloor, \end{cases}$$

it follows that the columns from \mathcal{B} span $\mathcal{C}(M_{k+\ell})$. Let $p(x, y) = \sum_{i,j} p_{ij} x^i y^j \in V$ be a polynomial and \hat{p} a vector of its coefficients ordered in the basis \mathcal{B} . We define a univariate polynomial $q_p(x)$ corresponding to $p(x, y)$, by

$$g_p(x) := p(x, x^{-\ell}) = \sum_{i,j} p_{ij} x^{i-\ell j} =: \sum_{s=-k\ell-1}^{k+1} g_{p,s} x^s \in \mathbb{R}[x]_{k\ell+1}. \quad (33)$$

Let \hat{g}_p be its vector of coefficients in the basis

$$\mathcal{B}_1 = \{x^{-k\ell-1}, x^{-k\ell}, \dots, x^{k+1}\}. \quad (34)$$

The monomials $x^{i_1} y^{j_1}, x^{i_2} y^{j_2}$ from \mathcal{B} correspond to the same monomial x^s by the correspondence (33) iff $i_1 - \ell j_1 = i_2 - \ell j_2$, which is further equivalent to $i_1 = i_2$ and $j_1 = j_2$ (since i_1 and i_2 are at most $\ell - 1$ in \mathcal{B}). Therefore

$$\hat{g}_p = \hat{p}. \quad (35)$$

We define two univariate sequences

$$\begin{aligned} \gamma &:= \gamma^{(-2k\ell, 2k)} = (\gamma_{-2k\ell}, \gamma_{-2k\ell+1}, \dots, \gamma_0, \dots, \gamma_{2k}) \in \mathbb{R}^{2k(1+\ell)+1}, \\ \tilde{\gamma} &:= \gamma^{(-2k\ell-2, 2k+2)} = (\gamma_{-2k\ell-2}, \gamma_{-2k\ell-1}, \gamma, \gamma_{2k+1}, \gamma_{2k+2}) \in \mathbb{R}^{2k(1+\ell)+5}, \end{aligned}$$

by the formula

$$\gamma_t = \begin{cases} \beta_{t,0}, & \text{if } t \geq 0, \\ \beta_{t+\ell \lceil \frac{|t|}{\ell} \rceil, \lceil \frac{|t|}{\ell} \rceil}, & \text{if } t < 0. \end{cases} \tag{36}$$

Note that for $t < 0$ we have that $t + \ell \lceil \frac{|t|}{\ell} \rceil \leq \ell - 1$, $\lceil \frac{|t|}{\ell} \rceil \leq 2k + 1$ (since $\ell \geq 3$) and hence

$$t + \ell \left\lceil \frac{|t|}{\ell} \right\rceil + \left\lceil \frac{|t|}{\ell} \right\rceil \leq \ell - 1 + 2k + 1 = 2k + \ell.$$

Therefore $\beta_{t+\ell \lceil \frac{|t|}{\ell} \rceil, \lceil \frac{|t|}{\ell} \rceil}$ is well-defined being an element of the matrix $M_{k+\ell}$ (since $2k + \ell \geq 2k + 2\ell$).

By the following claim solving the K -TMP for β is equivalent to solving the $(\mathbb{R} \setminus \{0\})$ -TMP for γ .

Claim 7: Let $u \in \mathbb{N}$. A sequence γ admits a u -atomic $(\mathbb{R} \setminus \{0\})$ -rm if and only if β admits a u -atomic K -rm.

Proof of Claim 7.: First we prove the implication (\Rightarrow) . Let x_1, \dots, x_u , be the atoms in the $(\mathbb{R} \setminus \{0\})$ -rm for γ with the corresponding densities ρ_1, \dots, ρ_u . We will prove that the atoms $(x_1, (x_1)^{-\ell}), \dots, (x_u, (x_u)^{-\ell})$ with densities ρ_1, \dots, ρ_p are the K -rm for β . We separate two cases:

(1) $\lfloor \frac{i}{\ell} \rfloor \geq j$:

$$\beta_{i,j} = \beta_{i-\ell j,0} = \gamma_{i-\ell j} = \sum_{p=0}^u \rho_p(x_p)^{i-\ell j} = \sum_{p=0}^u \rho_p(x_p)^i ((x_p)^{-\ell})^j,$$

where we used the fact that β is rg in the first equality, (36) in the second equality, the definitions of ρ_p, x_p in the third equality and split $(x_p)^{i-\ell j}$ into two parts in the last equality.

(2) $\lfloor \frac{i}{\ell} \rfloor < j$:

$$\begin{aligned} \beta_{i,j} &= \beta_{i \bmod \ell, j - \lfloor \frac{i}{\ell} \rfloor} = \gamma_{-(j - \lfloor \frac{i}{\ell} \rfloor)\ell + i \bmod \ell} = \sum_{p=0}^u \rho_p(x_p)^{-(j - \lfloor \frac{i}{\ell} \rfloor)\ell + i \bmod \ell} \\ &= \sum_{p=0}^u \rho_p(x_p)^{\lfloor \frac{i}{\ell} \rfloor \ell + i \bmod \ell} ((x_p)^{-\ell})^j \\ &= \sum_{p=0}^u \rho_p(x_p)^i ((x_p)^{-\ell})^j, \end{aligned}$$

where we used the fact that β is rg in the first equality, (36) in the second equality, the definitions of ρ_p, x_p in the third equality, split the exponent at x_p into two parts in the fourth equality and used that $\lfloor \frac{i}{\ell} \rfloor \ell + i \bmod \ell = i$ in the last equality.

This proves the implication (\Rightarrow) .

It remains to prove the implication (\Leftarrow) . Let $(x_1, (x_1)^{-\ell}), \dots, (x_u, (x_u)^{-\ell})$ be the atoms in the K -rm for β with the corresponding densities ρ_1, \dots, ρ_u . We will prove that the atoms (x_1, \dots, x_u) with densities ρ_1, \dots, ρ_p are the $(\mathbb{R} \setminus \{0\})$ -rm for γ :

- For $t \geq 0$ we have that

$$\gamma_t = \beta_{t,0} = \sum_{p=0}^u \rho_p (x_p)^t,$$

where we use the definition (36) in the first equality and the definitions of ρ_p, x_p in the second.

- For $t < 0$ we have that

$$\gamma_t = \beta_{t+\lceil \frac{|t|}{\ell} \rceil, \lceil \frac{|t|}{\ell} \rceil} = \sum_{p=0}^u \rho_p (x_p)^{t+\lceil \frac{|t|}{\ell} \rceil} ((x_p)^{-\ell})^{\lceil \frac{|t|}{\ell} \rceil} = \sum_{p=0}^u \rho_p (x_p)^t,$$

where we use the definition (36) in the first equality and the definitions of ρ_p, x_p in the second.

This proves the implication (\Leftarrow) . ■

Let $(M_{k+\ell})|_{\mathcal{B}}$ be the restriction of $M_{k+\ell}$ to the rows and columns indexed by monomials (capitalized) from \mathcal{B} . The following claim gives an explicit connection between $(M_{k+\ell})|_{\mathcal{B}}$ and the Hankel matrix $A_{\tilde{\gamma}}$ of the sequence $\tilde{\gamma}$.

Claim 8: We have that

$$(M_{k+\ell})|_{\mathcal{B}} = A_{\tilde{\gamma}}. \tag{37}$$

Proof of Claim 8.: Let $p(x, y) = \sum_{i,j} p_{ij} x^i y^j \in V$ and $r(x, y) = \sum_{i,j} r_{ij} x^i y^j \in V$ be polynomials from the vector subspace V and \hat{p}, \hat{r} vectors of their coefficients ordered in the basis \mathcal{B} . Let $\tilde{\beta} := \beta^{(2(k+1)\ell+2(\ell-1))}$. Then we have

$$\begin{aligned} (\hat{r})^T ((M_{k+\ell})|_{\mathcal{B}}) \hat{p} &= {}^1 L_{\tilde{\beta}}(pr) = L_{\tilde{\beta}} \left(\sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} r_{i_2 j_2} x^{i_1+i_2} y^{j_1+j_2} \right) \\ &= {}^2 \sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} r_{i_2 j_2} \beta_{i_1+i_2, j_1+j_2} \\ &= {}^3 \sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} r_{i_2 j_2} \gamma_{i_1+i_2-(j_1+j_2)\ell} \\ &= {}^4 L_{\tilde{\gamma}} \left(\sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} r_{i_2 j_2} x^{i_1+i_2-(j_1+j_2)\ell} \right) \\ &= {}^5 L_{\tilde{\gamma}} \left(\sum_{i_1, i_2, j_1, j_2} p_{i_1 j_1} x^{i_1-j_1\ell} \cdot r_{i_2 j_2} x^{i_2-j_2\ell} \right) \end{aligned}$$

$$\begin{aligned}
 &=^6 L_{\tilde{\gamma}} \left(\left(\underbrace{\sum_{i_1, j_1} p_{i_1 j_1} x^{i_1 - j_1 \ell}}_{g_p(x)} \right) \left(\underbrace{\sum_{i_2, j_2} r_{i_2 j_2} x^{i_2 - j_2 \ell}}_{g_r(x)} \right) \right) \\
 &=^7 \widehat{g}_r^T A_{\tilde{\gamma}} \widehat{g}_p = \widehat{r}^T A_{\tilde{\gamma}} \widehat{p},
 \end{aligned}$$

where in the first line we used the correspondence between the moment matrix and the Riesz functional $L_{\tilde{\beta}}$, the definition $L_{\tilde{\beta}}$ in the second, (36) and the fact that β is rg in the third (rg is needed if $i_1 + i_2 \geq \ell$), the definition of $L_{\tilde{\gamma}}$ in the fourth, decomposed the exponent of x into two parts in the fifth, decomposed a sum into the product of two sums in the sixth, in the seventh we used the correspondence between $A_{\tilde{\gamma}}$ and the Riesz functional $L_{\tilde{\gamma}}$, where $\widehat{g}_p, \widehat{g}_r$ are the vectors of coefficients of g_p and g_r in the basis \mathcal{B}_1 (see (34)) and (35). Since p and q were arbitrary from V , this proves Claim 8. ■

Since $(M_{k+\ell})|_{\mathcal{B}}$ is psd, it follows from (37) that $A_{\tilde{\gamma}}$ is psd. We separate two cases. Either A_{γ} is pd or A_{γ} is psd, singular, prg by [44, Theorem 2.6] and nrg by [46, Proposition 2.1.(5)]. By [46, Theorem 3.1], γ admits a $(\text{rank } A_{\gamma})$ -atomic $(\mathbb{R} \setminus \{0\})$ -rm. Since $\text{rank } M_k \leq \text{rank } A_{\gamma} \leq k(\ell + 1) + 1$, using Claim 8 the following holds:

(2') β has an s -atomic K -rm for some s satisfying

$$\text{rank } M_k \leq s \leq \text{rank } A_{\gamma} \leq k(\ell + 1) + 1. \tag{38}$$

To obtain (2) of Theorem 4.1, we need to decrease the upper bound in (38) by 1. Note that the bound $k(\ell + 1) + 1$ occurs only in the case A_{γ} is pd. We denote by $\gamma(z)$ a sequence obtained from the sequence γ by replacing $\gamma_{-2k\ell+1}$ with a variable z . The matrix $A_{\gamma(z)}$ is a partially positive definite matrix and by [48, Lemma 2.11] there exist two choices of z , which we denote by z^{\pm} , such that $A_{\gamma(z^{\pm})}$ is psd and has rank $k(\ell + 1)$. Using [46, Proposition 2.5] for the reversed sequence $(\gamma(z^{\pm}))^{(\text{rev})}$ of $\gamma(z^{\pm})$, we see that at least one of $(\gamma(z^{\pm}))^{(\text{rev})}$ admits a $(\mathbb{R} \setminus \{0\})$ -rm. Hence, at least one of $\gamma(z^{\pm})$ is prg and nrg, and admits a $k(\ell + 1)$ -atomic $(\mathbb{R} \setminus \{0\})$ -rm. If none of the moments $\beta_{i,j}$ of the sequence β depends on $\gamma_{-2k\ell+1}$, the $(\mathbb{R} \setminus \{0\})$ -rm for $(\gamma(z^{\pm}))^{(\text{rev})}$ will generate the K -rm for β as in the proof of Claim 8. But by definition (36), there is indeed no moment from β , which depends on $\gamma_{-2k\ell+1}$ (since we need to represent moments of degree at least $-2k\ell$ and at most $2k$, while $\gamma_{-2k\ell+1}$ corresponds to $\beta_{\ell-1, 2k}$ in some extension of β), which concludes the proof of Theorem 4.1. ■

To prove Theorem 4.2 only a little adaptation of the last part of the proof of Theorem 4.1 is needed, which we now explain.

Proof of Theorem 4.2.: The implications (1) \Rightarrow (4) and (2) \Rightarrow (1) are trivial. The implication (4) \Rightarrow (3) follows from [1, Theorem 3.14]. It remains to prove the implication (3) \Rightarrow (2). Following the proof of Theorem 4.1 everything remains the same until (2'). It remains to justify that the upper bound in (38) can be decreased to $m := k(\ell + 1) - \lfloor \frac{\ell}{2} \rfloor + 1$. If

rank $A_\gamma \leq m$, then we are already done. From now on we assume that $r := \text{rank } A_\gamma > m$. Since γ admits a $(\mathbb{R} \setminus \{0\})$ -representing measure, which we denote by μ , it is nrg and $\text{rank } \gamma = \text{rank } A_\gamma = \text{rank } A_\gamma[r - 1]$. Hence, $A_{\gamma^{(2k-2(r-1), 2k)}}$ is pd and in particular also its submatrix $A_{\gamma^{(2k-2(m-1), 2k)}}$ is pd. We denote by $\gamma(z_1, \dots, z_\ell)$ a sequence obtained from the sequence γ by replacing the moments $\gamma_{-2k\ell}, \gamma_{-2k\ell+1}, \dots, \gamma_{-2k\ell+\ell-1}$ with variables z_1, \dots, z_ℓ . By [46, Theorem 3.1], the sequence $\gamma^{(2k-2(m-1), 2k)}$ has a m -atomic $(\mathbb{R} \setminus \{0\})$ -rm (to apply [46, Theorem 3.1] we used that $2k - 2(m - 1) = -2k\ell + 2\lfloor \frac{\ell}{2} \rfloor < 0$). We denote the measure obtained in this way by μ_1 and generate its moment sequence $\gamma(z_1, \dots, z_\ell)$, where z_1, \dots, z_ℓ are the moments of degrees $-2k\ell, -2k\ell + 1, \dots, -2k\ell + \ell - 1$, respectively. If none of the moments $\beta_{i,j}$ of the sequence $\beta^{(2k-1)}$ depends on $\gamma_{-2k\ell}, \gamma_{-2k\ell+1}, \dots, \gamma_{-2k\ell+\ell-1}$, then μ_1 will generate the K -rm for $\beta^{(2k-1)}$ as in the proof of Claim 1 of Theorem 4.1. But by definition (36), there is indeed no moment from $\beta^{(2k-1)}$ depending on $\gamma_{-2k\ell}, \gamma_{-2k\ell+1}, \dots, \gamma_{-2k\ell+\ell-1}$, which concludes the proof of Theorem 4.2. ■

A corollary to Theorem 4.1 is an improvement of the bounds on the degrees of sums of squares in the Positivstellensatz [11, Corollary 6.4] for the curves of the form $yx^\ell = 1, \ell \in \mathbb{N} \setminus \{1\}$.

Corollary 4.3: *Let $K := \{(x, y) \in \mathbb{R}^2 : yx^\ell = 1\}$, where $\ell \in \mathbb{N} \setminus \{1\}$ and $k \geq \ell + 1$. If $r(x, y) \in \mathbb{R}[x, y]_{2k}$ is strictly positive on K , then r admits a decomposition*

$$r(x, y) = \sum_{i=1}^{\ell_1} f_i(x, y)^2 + (yx^\ell - 1) \sum_{i=1}^{\ell_2} g_i(x, y)^2 - (yx^\ell - 1) \sum_{i=1}^{\ell_3} h_i(x, y)^2,$$

where $\ell_1, \ell_2, \ell_3 \in \mathbb{Z}_+, f_i, g_i, h_i \in \mathbb{R}[x, y]$ and

$$\deg f_i^2 \leq 2m, \quad \deg((yx^\ell - 1)g_i^2) \leq 2m, \quad \deg((yx^\ell - 1)h_i^2) \leq 2m$$

with $m = k + \ell + 1$. where $\ell_1, \ell_2, \ell_3 \in \mathbb{Z}_+$.

Proof: By the equivalence (1) \Leftrightarrow (3) of Theorem 4.1, the set K has the property $(R_{k,\ell})$ in the notation of [52, p. 2713]. Now the result follows by [52, Theorem 1.5]. ■

Remark 4.2: The bound on m in Theorem 4.3 in [11, Corollary 6.4] is quadratic in k and ℓ , namely $(2k + 2)(2 + \ell) - (1 + \ell)$.

4.2. A solution to the TMP based on the feasibility of a linear matrix inequality

In this subsection we give another alternative solution to the TMP on curves yx^ℓ , where $\ell \in \mathbb{N} \setminus \{1\}$, which is based on the feasibility of a linear matrix inequality associated to the univariate sequence γ , obtained from the original sequence β as in the proof of the results in the previous subsection. The feasibility question appears as a result of the fact that γ is not fully determined by β , but β admits a K -representing measure if and only if γ can be completed to a sequence admitting a $(\mathbb{R} \setminus \{0\})$ -representing measure.

For $n_1, n_2 \in \mathbb{Z}, n_1 \leq n_2$, we denote by $[n_1 : n_2]$ the set of all integers between n_1 and n_2 . Let the sets N_1, N_2 form a partition of $[n_1 : n_2]$, i.e. $N_1, N_2 \subseteq [n_1 : n_2], N_1 \cup N_2 = [n_1 : n_2]$

and $N_1 \cap N_2 = \emptyset$. Let $\Gamma_1 := (\gamma_t)_{t \in N_1}$ be a sequence of real numbers indexed by integers from N_1 and $\Gamma_2 := (\mathbf{y}_t)_{t \in N_2}$ a tuple of variables indexed by integers from N_2 . Let

$$F_{\Gamma_1}(\Gamma_2) : \mathbb{R}^{|N_2|} \rightarrow \mathbb{R}^{|N_1|+|N_2|} \tag{39}$$

be a function with the output a sequence $(\tilde{\gamma}_t)_{t \in [n_1:n_2]}$ where $\tilde{\gamma}_t = \begin{cases} \gamma_t, & \text{if } t \in N_1, \\ \mathbf{y}_t, & \text{if } t \in N_2. \end{cases}$

In Theorem 4.4 the set N_1 will be the set of indices, for which the corresponding univariate sequence γ is determined by β , while the indices of the non-determined part will belong to N_2 . Since we can either get a sequence with the lowest and highest degree terms both of odd degree or both of even degree or only the highest term of odd degree, for which the solutions to the STHMP are slightly different, we separate three cases for $N_1 \cup N_2 = [n_1 : n_2]$ (see (41)).

Theorem 4.4: Let $K := \{(x, y) \in \mathbb{R}^2 : yx^\ell = 1\}$, $\ell \in \mathbb{N} \setminus \{1\}$, and

$$\beta := \beta^{(d)} = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq d},$$

where $\lceil \frac{d}{2} \rceil \geq \ell + 1$. Define

$$N_1 := \{t \in \mathbb{Z}_- : t = -i\ell + j \text{ for some } 0 \leq j < \ell, i \in \mathbb{Z}_+ \text{ and } i + j \leq d\},$$

$$\gamma_t = \begin{cases} \beta_{t,0}, & \text{if } t \in (\mathbb{N} \cup \{0\}) \cap N_1, \\ \beta_{t+\ell \lceil \frac{|t|}{\ell} \rceil, \lceil \frac{|t|}{\ell} \rceil}, & \text{if } t \in N_1 \setminus (\mathbb{N} \cup \{0\}), \end{cases} \tag{40}$$

and $\Gamma_1 := (\gamma_t)_{t \in N_1}$. Let

$$[n_1 : n_2] := \begin{cases} [-d\ell - 2 : d + 2], & \text{if } d \text{ is even,} \\ [-d\ell - 2 : d + 1], & \text{if only } \ell \text{ is even,} \\ [-d\ell - 1 : d + 1], & \text{if } d, \ell \text{ are odd,} \end{cases} \tag{41}$$

$\Gamma_2 := (\mathbf{y}_t)_{t \in N_2}$ be a tuple of variables with $N_2 = [n_1 : n_2] \setminus N_1$ and $F_{\Gamma_1}(\Gamma_2)$ be defined as in (39). Then the following statements are equivalent:

- (1) β has a K -representing measure.
- (2) $\beta_{i+\ell,j+1} = \beta_{i,j}$ for every $i, j \in \mathbb{Z}_+$ such that $i + j \leq d - \ell - 1$ and there exists a tuple $\Gamma_2 = (\mathbf{y}_t)_{t \in N_2} \in \mathbb{R}^{N_2}$ such that $A_{F_{\Gamma_1}(\Gamma_2)} \geq 0$.

Proof: Assume that d is even. Observing the proof of Theorem 4.1 one can notice that $F_{\Gamma_1}(\Gamma_2)$ corresponds to the sequence $\tilde{\gamma}$. The original sequence β determines only γ_t for $t \in N_1$ by (40), while for $t \in N_2$, \mathbf{y}_t are variables. By the proof of Theorem 4.1, β will have a K -rm iff it satisfies the rg relations coming from the column relation $YX^\ell = 1$ and there exists $\tilde{\gamma}$ such that $A_{\tilde{\gamma}} \geq 0$. This proves Theorem 4.4 for even d .

Observing the proof of Theorem 4.2 in case d is odd one can notice that only

$$\gamma^{(-d\ell, d)} = (\gamma_{-d\ell}, \gamma_{-d\ell+1}, \dots, \gamma_{d-1}, \gamma_d)$$

needs to have a $(\mathbb{R} \setminus \{0\})$ -rm to obtain a K -rm for β . In case $d\ell$ is even, this is equivalent to $A_{\gamma^{(-d\ell-2, d+1)}} \geq 0$, where

$$\gamma^{(-d\ell-2, d+1)} = (\gamma_{-d\ell-2}, \gamma_{-d\ell-1}, \dots, \gamma_d, \gamma_{d+1})$$

for some $\gamma_{-d\ell-2}$, $\gamma_{-d\ell-1}$ and γ_{d+1} . Since $\gamma^{(-d\ell-2, d+1)}$ corresponds to the sequence $F_{\Gamma_1}(\Gamma_2)$ for even ℓ and odd d , this proves Theorem 4.2 in this case. If $d\ell$ is odd, then it suffices that there are $\gamma_{-d\ell-1}$ and γ_{d+1} such that $A_{\gamma^{(-d\ell-1, d+1)}} \geq 0$, where

$$\gamma^{(-d\ell-1, d+1)} = (\gamma_{-d\ell-1}, \gamma_{-d\ell}, \dots, \gamma_d, \gamma_{d+1}).$$

Since $\gamma^{(-d\ell-1, d+1)}$ corresponds to the sequence $F_{\Gamma_1}(\Gamma_2)$ for odd $d\ell$, this proves Theorem 4.2 in this case. ■

We will present the statement of Theorem 4.4 on a few examples. The following example is for $\ell = 2$ and a sequence β of even degree.

Example 4.5: Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$ be a bivariate sequence of degree $2k$, $k \geq 3$, and $K := \{(x, y) \in \mathbb{R}^2 : yx^2 = 1\}$. For the existence of a K -rm β must satisfy the relations $\beta_{i,j} = \beta_{i+2, j+1}$ for every $i, j \in \mathbb{Z}_+$ such that $i + j \leq 2k - 3$. In the notation of Theorem 4.4,

$$N_1 := \{t \in \mathbb{Z} : -4k \leq t \leq 2k, t \neq -4k + 1\},$$

$$[n_1 : n_2] := [-4k - 2 : 2k + 2], \quad N_2 = \{-4k - 2, -4k - 1, -4k + 1, 2k + 1, 2k + 2\},$$

the formula (36) is equal to

$$\gamma_t = \begin{cases} \beta_{t,0}, & \text{if } t \in \mathbb{Z}_+ \cap N_1, \\ \beta_{t+2 \lceil \frac{|t|}{2} \rceil, \lceil \frac{|t|}{2} \rceil}, & \text{if } t \in N_1 \setminus (\mathbb{Z}_+ \cap N_1), \end{cases}$$

the function $F_{\Gamma_1} : \mathbb{R}^5 \rightarrow \mathbb{R}^{6k+4}$ is defined by

$$\begin{aligned} F_{\Gamma_1}(\Gamma_2) &= F_{\Gamma_1}(\gamma_{-4k-2}, \gamma_{-4k-1}, \gamma_{-4k+1}, \gamma_{2k+1}, \gamma_{2k+2}) \\ &:= (\gamma_{-4k-2}, \gamma_{-4k-1}, \gamma_{-4k}, \gamma_{-4k+1}, \gamma_{-4k+2}, \dots, \gamma_{2k}, \gamma_{2k+1}, \gamma_{2k+2}), \end{aligned}$$

and the matrix $A_{F_{\Gamma_1}(\Gamma_2)}$ is equal to

$$\begin{pmatrix} \gamma_{-4k-2} & \gamma_{-4k-1} & \gamma_{-4k} & \gamma_{-4k+1} & \gamma_{-4k+2} & \cdots & \gamma_k & \gamma_{k+1} \\ \gamma_{-4k-1} & \gamma_{-4k} & \gamma_{-4k+1} & \gamma_{-4k+2} & \ddots & & \gamma_{k+1} & \gamma_{k+2} \\ \gamma_{-4k} & \gamma_{-4k+1} & \gamma_{-4k+2} & \ddots & & & \gamma_{k+2} & \vdots \\ \gamma_{-4k+1} & \beta_{0,2k-1} & \ddots & & & & \vdots & \vdots \\ \gamma_{-4k+2} & \ddots & & & & & \vdots & \gamma_{2k} \\ \vdots & & & & & & \gamma_{2k} & \gamma_{2k+1} \\ \gamma_{k+1} & \cdots & \cdots & \cdots & \gamma_{2k-1} & \gamma_{2k} & \gamma_{2k+1} & \gamma_{2k+2} \end{pmatrix}$$

The question of feasibility of $A_{F_{\Gamma_1}}(\Gamma_2)$ can be answered analytically, since the structure of the missing entries is simple enough. Actually it is even easier to work with $A_{(\gamma_{-4k}, \gamma_{-4k+1}, \gamma_{-4k+2}, \dots, \gamma_{2k})}$ and answer the feasibility question together with the conditions from the solution of [46, Theorem 2.1] (see [46, Theorem 4.1]).

The following example demonstrates the statement of Theorem 4.4 for the case $\gamma x^3 = 1$ and a sequence β of even degree.

Example 4.6: Let $\beta = (\beta_{i,j})_{i,j \in \mathbb{Z}_+, i+j \leq 2k}$ be a bivariate sequence of degree $2k$, $k \geq 4$, and $K := \{(x, y) \in \mathbb{R}^2 : \gamma x^3 = 1\}$. For the existence of a K -rm β must satisfy the relations $\beta_{i,j} = \beta_{i+3,j+1}$ for every $i, j \in \mathbb{Z}_+$ such that $i + j \leq 2k - 4$. In the notation of Theorem 4.4,

$$\begin{aligned}
 N_1 &:= \{t \in \mathbb{Z} : -6k \leq t \leq 2k, t \notin \{-6k + 1, -6k + 2, -6k + 5\}\}, \\
 [n_1 : n_2] &:= [-6k - 2 : 2k + 2], \\
 N_2 &:= \{-6k - 2, -6k - 1, -6k + 1, -6k + 2, -6k + 5, 2k + 1, 2k + 2\},
 \end{aligned}$$

the formula (36) is equal to

$$\gamma_t = \begin{cases} \beta_{t,0}, & \text{if } t \in \mathbb{Z}_+ \cap N_1, \\ \beta_{t+3 \lceil \frac{|t|}{3} \rceil, \lceil \frac{|t|}{3} \rceil}, & \text{if } t \in N_1 \setminus (\mathbb{Z}_+ \cap N_1), \end{cases}$$

the function $F_{\Gamma_1} : \mathbb{R}^7 \rightarrow \mathbb{R}^{8k+4}$ is defined by

$$\begin{aligned}
 F_{\Gamma_1}(\Gamma_2) &= F_{\Gamma_1}(\gamma_{-6k-2}, \gamma_{-6k-1}, \gamma_{-6k+1}, \gamma_{-6k+2}, \gamma_{-6k+5}, \gamma_{2k+1}, \gamma_{2k+2}) \\
 &:= (\gamma_{-6k-2}, \gamma_{-6k-1}, \gamma_{-6k}, \gamma_{-6k+1}, \gamma_{-6k+2}, \gamma_{-6k+3}, \gamma_{-6k+4}, \\
 &\quad \gamma_{-6k+5}, \gamma_{-6k+6}, \dots, \gamma_{2k}, \gamma_{2k+1}, \gamma_{2k+2}),
 \end{aligned}$$

and the matrix $A_{F_{\Gamma_1}(\gamma_{-6k-2}, \gamma_{-6k-1}, \gamma_{-6k+1}, \gamma_{-6k+2}, \gamma_{-6k+5}, \gamma_{2k+1}, \gamma_{2k+2})}$ is equal to

γ_{-6k-2}	γ_{-6k-1}	γ_{-6k}	γ_{-6k+1}	γ_{-6k+2}	γ_{-6k+3}	γ_{-6k+4}	γ_{-6k+5}	\dots	γ_k	γ_{k+1}
γ_{-6k-1}	γ_{-6k}	γ_{-6k+1}	γ_{-6k+2}	γ_{-6k+3}	γ_{-6k+4}	γ_{-6k+5}	\dots	\dots	γ_{k+1}	γ_{k+2}
γ_{-6k}	γ_{-6k+1}	γ_{-6k+2}	γ_{-6k+3}	γ_{-6k+4}	γ_{-6k+5}	\dots			γ_{k+2}	\vdots
γ_{-6k+1}	γ_{-6k+2}	γ_{-6k+3}	γ_{-6k+4}	γ_{-6k+5}	\dots				\vdots	\vdots
γ_{-6k+2}	γ_{-6k+3}	γ_{-6k+4}	γ_{-6k+5}	\dots					\vdots	\vdots
\vdots	γ_{-6k+4}	γ_{-6k+5}	\dots						\vdots	\vdots
\vdots	\dots	\dots							\vdots	\vdots
\vdots	\dots								\vdots	γ_{2k}
\vdots									γ_{2k}	γ_{2k+1}
γ_{k+1}	\dots	\dots	\dots	\dots	\dots	\dots	γ_{2k-1}	γ_{2k}	γ_{2k+1}	γ_{2k+2}

In contrast to the situation $\gamma x^2 = 1$ from Example 4.5, the structure of the missing entries here is too complicated for the analytic approach and we believe the feasibility question can only be answered numerically using linear matrix inequality solvers.

Remark 4.3: It would be interesting to know, to what extent does the result [50, Corollary 7.6] (see Remark 3.1.(3.1)) extend to even degree sequences on all plane curves. As

explained in Remark 3.1. (3), one needs one more atom in the upper bound for curves of the form $y = q(x)$ with $\deg q = 2$ and the same is true if $\deg q \leq 1$ by Remark 3.1.(4). On the other hand, the results of the present paper suggest that for curves $\mathcal{Z}(p)$, $\deg p \geq 3$, the upper bound could be $k \deg p$. Also from the concrete solution to the TMP on the curve $\mathcal{Z}(y^2 - x^3)$ [48, Corollary 4.3] it follows that the same bound works. However, the forthcoming result of Bhardwaj [59] shows that also for degree 3 curves the upper bound has to be loosened, by constructing a truncated moment sequence of degree $2k = 6$ on a curve $\mathcal{Z}(p)$, where $p(x, y) = y^2 - x^3 + ax - 1$, $a = \frac{524287}{262144}$, with a minimal measure consisting of 10 atoms, which is $k \deg p + 1$.

Acknowledgments

I would like to thank the anonymous referees for very useful comments and suggestions for improving the manuscript. Numerical examples in this paper were obtained using the software tool *Mathematica* [59].

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This work was supported by the Slovenian Research Agency grants J1-2453, J1-3004, P1-0288.

ORCID

A. Zalar  <http://orcid.org/0000-0001-9630-4014>

References

- [1] Curto R, Fialkow L. Solution of the truncated complex moment problem for flat data. *Mem Amer Math Soc.* 1996;119, no. 568.
- [2] Curto R, Fialkow L. Flat extensions of positive moment matrices: recursively generated relations. *Mem Amer Math Soc.* 1998;136:648.
- [3] Curto R, Fialkow L. Truncated K -moment problems in several variables. *J Oper Theory.* 2005;54:189–226.
- [4] Laurent M. Revisiting two theorems of Curto and Fialkow on moment matrices. *Proc Amer Math Soc.* 2005;133:2965–2976.
- [5] Curto R, Fialkow L. Solution of the singular quartic moment problem. *J Oper Theory.* 2002;48:315–354.
- [6] Curto R, Fialkow L. Solution of the truncated parabolic moment problem. *Integral Equ Oper Theory.* 2004;50:169–196. doi: 10.1007/s00020-003-1275-3.
- [7] Curto R, Fialkow L. Solution of the truncated hyperbolic moment problem. *Integral Equ Oper Theory.* 2005;52:181–218. doi: 10.1007/s00020-004-1340-6.
- [8] Fialkow L. The truncated moment problem on parallel lines. *Varied Landsc Oper Theory.* 2014;20:99–116.
- [9] Curto R, Yoo S. Concrete solution to the nonsingular quartic binary moment problem. *Proc Amer Math Soc.* 2016;144:249–258. doi: 10.1090/proc/12698.
- [10] Fialkow L, Nie J. Positivity of Riesz functionals and solutions of quadratic and quartic moment problems. *J Funct An.* 2010;258:328–356.
- [11] Fialkow L. Solution of the truncated moment problem with variety $y = x^3$. *Trans Amer Math Soc.* 2011;363:3133–3165.

- [12] Curto R, Fialkow L. Recursively determined representing measures for bivariate truncated moment sequences. *J Oper Theory*. 2013;70:401–436.
- [13] Curto R, Fialkow L, Möller HM. The extremal truncated moment problem. *Integral Equ Oper Theory*. 2008;60:177–200. doi: 10.1007/s00020-008-1557-x.
- [14] Curto R, Yoo S. Non-extremal sextic moment problems. *J Funct Anal*. 2015;269:758–780. doi: 10.1016/j.jfa.2015.04.014.
- [15] Yoo S. Sextic moment problems on 3 parallel lines. *Bull Korean Math Soc*. 2017;54:299–318.
- [16] Yoo S. Sextic moment problems with a reducible cubic column relation. *Integral Equ Oper Theory*. 2017;88:45–63.
- [17] Blekherman G. Positive Gorenstein ideals. *Proc Amer Math Soc*. 2015;143:69–86. doi: 10.1090/S0002-9939-2014-12253-2.
- [18] Blekherman G, Fialkow L. The core variety and representing measures in the truncated moment problem. *J Oper Theory*. 2020;84:185–209.
- [19] di Dio PJ, Schmüdgen K. The multidimensional truncated moment problem: atoms, determinacy, and core variety. *J Funct Anal*. 2018;274:3124–3148. doi: 10.1016/j.jfa.2017.11.013.
- [20] Fialkow L. The core variety of a multisequence in the truncated moment problem. *J Math Anal Appl*. 2017;456:946–969. doi: 10.1016/j.jmaa.2017.07.041.
- [21] Kimsey DP. The cubic complex moment problem. *Integral Equ Oper Theory*. 2014;80:353–378. doi: 10.1007/s00020-014-2183-4.
- [22] Fialkow L. Positivity, extensions and the truncated complex moment problem. *Contemp Math*. 1995;185:133–150.
- [23] Kimsey DP. On a minimal solution for the indefinite multidimensional truncated moment problem. *J Math Anal Appl*. 2021;500(1):125091. doi:10.1016/j.jmaa.2021.125091.
- [24] Kimsey DP, Putinar M. The moment problem on curves with bumps. *Math Z*. 2021;298(3-4):935–942. doi:10.1007/s00209-020-02633-2.
- [25] Wolfram Research, Inc. *Mathematica*, Version 12.0. Champaign, IL: Wolfram Research, Inc; 2020.
- [26] Akhiezer NI. *The classical moment problem and some related questions in analysis*. New York: Hafner Publishing Co; 1965.
- [27] Akhiezer NI, Krein M. *Some questions in the theory of moments*. Providence: American Math. Soc; 1962. (Transl. Math. Monographs; Vol. 2).
- [28] Krein MG, Nudelman AA. *The Markov moment problem and extremal problems*. Translations of Mathematical Monographs, Amer. Math. Soc. 1977.
- [29] Schmüdgen K. *The moment problem*. Cham: Springer; 2017. (Graduate Texts in Mathematics; Vol. 277).
- [30] Alpay D, Jorgensen PET, Kimsey DP. Moment problems in an infinite number of variables. *Infin Dimens Anal Quantum Probab Relat Top*. 2015;18.1550024
- [31] Ghasemi M, Kuhlmann S, Marshall M. Moment problem in infinitely many variables. *Israel J Math*. 2016;212:989–1012.
- [32] Infusino M, Kuna T, Lebowitz JL, et al. The truncated moment problem on \mathbb{N}_0 . *J Math Anal Appl*. 2017;452:443–468. doi: 10.1016/j.jmaa.2017.02.060.
- [33] Nie J. The \mathcal{A} -truncated K -moment problem. *Found Comput Math*. 2014;14:1243–1276.
- [34] Curto R, Ghasemi M, Infusino M, et al. The truncated moment problems for unital commutative \mathbb{R} -algebras. arxiv preprint. Available at: <https://arxiv.org/pdf/2009.05115.pdf>.
- [35] Ando T. Truncated moment problems for operators. *Acta Sci Math Szeged*. 1970;31:319–334.
- [36] Burgdorf S, Klep I. The truncated tracial moment problem. *J Oper Theory*. 2012;68:141–163.
- [37] Bhardwaj A, Zalar A. The singular bivariate quartic tracial moment problem. *Complex Anal Oper Theory*. 2018;12:1057–1142. doi: 10.1007/s11785-017-0756-3.
- [38] Bhardwaj A, Zalar A. The tracial moment problem on quadratic varieties. *J Math Anal Appl*. 2021;498:124936doi: 10.1016/j.jmaa.2021.124936.
- [39] Bakonyi M, Woerdeman HJ. *Matrix completions, moments, and sums of Hermitian squares*. Princeton: Princeton University Press, Princeton; 2011.
- [40] Dyukarev YM, Fritzsche B, Kirstein B, et al. On distinguished solutions of truncated matricial Hamburger moment problems. *Complex Anal Oper Theory*. 2009;3:759–834.

- [41] Dette H, Studden WJ. Matrix measures, moment spaces and Favard's theorem for the interval $[0, 1]$ and $[0, \infty)$. *Linear Algebra Appl.* **2002**;345:169–193.
- [42] Kimsey DP, Trachana M. On a solution of the multidimensional truncated matrix-valued moment problem. *Milan J Math.* **2022**;90:17–101. doi: [10.1007/s00032-021-00346-7](https://doi.org/10.1007/s00032-021-00346-7).
- [43] Kimsey DP, Woerdeman H. The truncated matrix-valued K -moment problem on $\mathbb{R}^d, \mathbb{C}^d, \mathbb{T}^d$. *Trans Amer Math Soc.* **2013**;365:5393–5430. doi: [10.1090/S0002-9947-2013-05812-6](https://doi.org/10.1090/S0002-9947-2013-05812-6).
- [44] Curto R, Fialkow L. Recursiveness, positivity, and truncated moment problems. *Houston J Math.* **1991**;17:603–635.
- [45] Curto R, Yoo S. Cubic column relations in the truncated moment problems. *J Funct Anal.* **2014**;266:1611–1626.
- [46] Zalar A. The strong truncated Hamburger moment problem with and without gaps. *J Math Anal Appl.* **2022**;516:126563. doi: [10.1016/j.jmaa.2022.126563](https://doi.org/10.1016/j.jmaa.2022.126563).
- [47] Simonov KK. Strong truncated matrix moment problem of Hamburger. *Sarajevo J Math.* **2016**;215:181–204.
- [48] Zalar A. The truncated Hamburger moment problem with gaps in the index set. *Integral Equ Oper Theory.* **2021**;93:36. doi: [10.1007/s00020-021-02628-6](https://doi.org/10.1007/s00020-021-02628-6).
- [49] Zalar A. The truncated moment problem on the union of parallel lines. *Linear Algebra Appl.* **2022**;649:186–239. doi: [10.1016/j.laa.2022.05.008](https://doi.org/10.1016/j.laa.2022.05.008).
- [50] Riener C, Schweighofer M. Optimization approaches to quadrature: a new characterization of Gaussian quadrature on the line and quadrature with few nodes on plane algebraic curves, on the plane and in higher dimensions. *J Complex.* **2018**;45:22–54.
- [51] Lasserre JB. *Moments, positive polynomials and their applications*. London: Imperial College Press; **2009**.
- [52] Curto R, Fialkow L. An analogue of the Riesz–Haviland theorem for the truncated moment problem. *J Funct Anal.* **2008**;255:2709–2731.
- [53] Stochel J. Solving the truncated moment problem solves the moment problem. *Glasgow J Math.* **2001**;43:335–341.
- [54] Ramana MV. An exact duality theory for semidefinite programming and its complexity implications. *Math Program.* **1997**;77:129–162.
- [55] Klep I, Schweighofer M. An exact duality theory for semidefinite programming based on sums of squares. *Math Oper Res.* **2012**;38:569–590. doi: [10.1287/moor.1120.0584](https://doi.org/10.1287/moor.1120.0584).
- [56] Porkolab L, Khachiyan L. On the complexity of semidefinite programs. *J Glob Optim.* **1997**;10:351–365.
- [57] Richter H. Parameterfreie Abschätzung und Realisierung von Erwartungswerten. *Bl der Deutsch Ges Versicherungsmath.* **1957**;3:147–161.
- [58] Albert A. Conditions for positive and nonnegative definiteness in terms of pseudoinverses. *SIAM J Appl Math.* **1969**;17:434–440.
- [59] Bhardwaj A. Flat extensions for the truncated moment problem on curves $y^2 = qx$ with $\deg q = 3$. in preparation.