

research article

CD56-positive diffuse large B-cell lymphoma: comprehensive analysis of clinical, pathological, and molecular characteristics with literature review

Gorana Gasljevic, Lucka Boltezar, Srdjan Novakovic, Vita Setrajcic-Dragos, Barbara Jezersek-Novakovic, Veronika Kloboves-Prevodnik

Institute of Oncology Ljubljana, Ljubljana, Slovenia

Radiol Oncol 2023; 57(2): 249-256.

Received 9 January 2023
Accepted 19 February 2023

Correspondence to: Assoc. prof. Veronika Kloboves-Prevodnik, M.D., Ph.D., Institute of Oncology Ljubljana, Zaloška 2, SI-1000 Ljubljana, Slovenia. E-mail: vkloboves@onko-i.si

Disclosure: No potential conflicts of interest were disclosed.

This is an open access article distributed under the terms of the CC-BY license (<https://creativecommons.org/licenses/by/4.0/>).

Background. Diffuse large B-cell lymphoma (DLBCL) is the most common non-Hodgkin lymphoma. The expression of CD56 in DLBCL is highly unusual. Little is known about its incidence and clinical importance. So far, no genetic profiling was performed in CD56 positive DLBCL.

Patients and methods. Tissue microarrays have been constructed, sectioned, and stained by H&E and immunohistochemistry for 229 patients with DLBCL diagnosed 2008–2017. For CD56 positive cases, clinical data was collected including age at diagnosis, stage of the disease, International Prognostic Index (IPI) score, treatment scheme and number of chemotherapy cycles, radiation therapy, treatment outcome, and possible relapse of the disease. Overall survival (OS) and progression-free survival (PFS) were calculated. For four patients, RNA was extracted and targeted RNA (cDNA) sequencing of 125 genes was performed with the Archer FusionPlex Lymphoma kit.

Results. CD56 expression was found in 7 cases (3%). The intensity of expression varied from weak to moderate focal, to very intensive and diffuse. All patients had *de novo* DLBCL. The median age at the time of diagnosis was 54.5 years. Five of them were women and 2 males. According to the Hans algorithm, 6 patients had the germinal centre B cells (GBC) type and one non-GBC (activated B-cell [ABC]) type, double expressor. Genetic profiling of four patients according to Schmitz's classification showed that 1 case was of the BN2 subtype, 1 of EZB subtype, 2 were unclassified. The six treated patients reached a complete response and did not experience progression of the disease during the median follow-up period of 80.5 months.

Conclusions. We report on one of the largest series of CD56+DLBCL with detailed clinicopathological data and for the first time described genetical findings in a limited number of patients. Our results show that CD56 expression is rare, but seems to be present in prognostic favourable subtypes of DLBCL not otherwise specified (NOS) as tested by immunohistochemical or genetic profiling.

Key words: diffuse large B-cell lymphoma not otherwise specified; CD56; immunohistochemistry; genetic profiling; prognosis

Introduction

CD56, also known as the neural cell adhesion molecule (NCAM), is a member of the immunoglobu-

lin superfamily that plays important functional roles during nervous system development, differentiation, and immune surveillance. In addition to neurons and glial cells, CD56 is normally also

expressed in neuroendocrine tissues and some cells of the hematopoietic system like NK cells and activated T lymphocytes.¹ In the hematopathology service, it is mainly used as a marker of NK cells and their neoplastic counterparts. Its aberrant expression is useful as a proof of clonal plasma cell proliferation, while it can also be used as prognostic marker in plasmacytoma, as well as in acute myeloid leukemia (AML) and acute lymphoblastic leukaemia (ALL).²⁻⁵

Diffuse large B-cell lymphoma (DLBCL) is the most common lymphoma, representing approximately one third of all non-Hodgkin lymphomas.² Cases of DLBCL that do not fit the distinctive clinical presentation, tissue morphology, neoplastic cell phenotype, and/or pathogen-associated criteria of other subtypes of DLBCL are termed "DLBCL not otherwise specified (DLBCL NOS)" and represent 80–85% of all DLBCL cases.² The WHO 2016 classification of hematopoietic neoplasms² requires that the neoplastic cells in DLBCL NOS be further defined based on whether they are derived from germinal centre B cells or activated B-cells as identified by gene expression profiling (GEP) or are germinal centre B cells (GBC) or non-GBC as identified by immunohistochemical (IHC) analyses. In general, DLBCL NOS is an aggressive disease with an overall long-term survival rate in patients treated with standard chemotherapy regimens of ~60%.^{7,8} Patients with activated B-cell (ABC) DLBCL and non-GBC variants have significantly worse prognoses than patients with the GBC variant.⁶ Expression of markers in DLBCL NOS neoplastic cells that have clinical significance as prognostic or predictive factors include CD5, MYC, BCL2, BCL6, CD20, CD19, CD22, CD30, PD-L1, and PD-L2.^{2,6} For example, 5–10% of DLBCL NOS cases express CD5 and have a very poor prognosis that is not improved by even aggressive treatment regimens, while the expression of CD30 represents a favourable prognostic indicator.²

Very little is known about the incidence and clinical importance of CD56 expression in DLBCL. In the last 30 years, the literature has only a few case reports or small series of CD56+ DLBCL with conflicting results on its importance.¹⁰⁻¹⁸ It could have a prognostic value; however, since new target drugs are becoming available and among them is also anti-CD56 antibody, CD56 could serve as a potential target for the treatment of patients who do not respond to standard therapeutic schemes.

The purpose of this study was to evaluate CD56 expression in DLBCL in our series, to estimate its

relationship to epidemiological factors, to roughly estimate its value as a prognostic marker, and to describe, for the first time the molecular findings in a subset of cases.

Patients and methods

Specimens

Data bases of the Department of Pathology Institute of Oncology Ljubljana (IOL) have been searched for all cases of DLBCL diagnosed between 2008 and 2017. Only the cases in which appropriate amount of material was present that could allow the construction of tissue microarrays (229) have been chosen for the study. Tissue microarrays have been constructed, sectioned, and stained by H&E and immunohistochemistry for the Hans algorithm as previously described.¹⁹ Also, for the cases that were CD56 positive, flow cytometric and/or immunocytochemical staining results and data were retrieved and re-analysed from the database of the Department of Cytopathology.

Patients

For selected patients, clinical data was collected including age at diagnosis, stage of the disease, IPI score, treatment scheme and number of cycles, potential radiation therapy, outcome and possible relapse of the disease were also noted. Overall survival (OS) and progression-free survival (PFS) were calculated. Subjects were censored at their last visit to the IOL and for those who finished follow-up at IOL, a vital status from the Cancer Registry of the Republic of Slovenia. All procedures followed in this evaluation were in accordance with the ethical standards of the responsible committee on human experimentation (Ethical Committee of Institute of Oncology Ljubljana, approval number: ERID-KESOPKR-23 and the Ethical Committee of the Republic of Slovenia, approval number: 58/02/15) and the Helsinki Declaration of 1975, as revised in 2000.

Immunohistochemistry

3–4 µm thick, formalin-fixed paraffin-embedded sections of constructed TMAs were used for immunohistochemical staining with the monoclonal antibody CD56. Staining was performed on the Ventana Benchmark platform using the MRQ 42 clone (Cell Marque) in dilution 1:200.

Flow cytometric analysis and immunocytochemistry

The preparation of FNAB (fine needle aspiration biopsy) lymph node sample, cell counting, sample preparations for flow cytometric immunophenotyping, acquisition of cells with flow cytometer and measurement result analysis were performed as previously described.²⁰ Monoclonal antibodies against CD45, CD19, CD20, CD3, CD10, CD5, CD23, FMC7, κ and λ LCs (BD Biosciences, New Jersey, U.S.) were used. The samples were acquired using a four-colour flow cytometer FACSCalibur (BD Biosciences, New Jersey, U.S.), a six-colour flow cytometer FACSCanto II (BD Biosciences, New Jersey, U.S.) or a ten-colour FACSCanto X (BD Biosciences, New Jersey, U.S.). The measurement results were analysed using CellQuest (BD Biosciences, New Jersey, U.S.) or BD FACSDiva software (BD Biosciences, New Jersey, U.S.). For immunocytochemical staining, methanol and Delaunay-fixed cytopines were prepared. Stainings were carried out on the Ventana Benchmark Ultra platform using antibodies against CD56, CK AE1/AE3, CK18 (DAKO), CD20 (Cell Marque, Rocklin, California, U.S.), synaptophysin (Termo Scientific, Waltham, Massachusetts, U.S.), CD3 and TTF-1 (Leica Biosystems, Nussloch, Germany).

Molecular analysis - NGS sequencing

RNA was extracted from 4 paraffin-embedded tissue samples was extracted using the MagMAX™ FFPE DNA/RNA Ultra Kit (ThermoFisher, Waltham, MA, USA). Samples were treated with DNase, during the extraction process. Targeted RNA (cDNA) sequencing of 125 genes was performed with the Archer FusionPlex Lymphoma kit (Invitae-ArcherDX, San Francisco, CA, USA). The final NGS library was quantified using the KAPA Library Quantification Kit (KAPA Biosystems, Merck, Ljubljana, Slovenia) and pair-end sequenced on a MiSeqDx system (Illumina, San Diego, CA, USA). The trimmed FASTQ file was uploaded to Archer Analysis software Version 6.0.3.2, which performed variant and fusion calls along with the determination of cell of origin (ABC or GCB). Variants were considered true positive if the frequency of the variant allele was above 10%, with minimum coverage of 100x.²⁰ All variants reported in GnomAD were excluded. Fusions were considered true positive if the fusion event was covered with a minimum of 5 unique reads and the percentage of reads supporting the event was above 10%.^{21,22}




FIGURE 1. (A) Morphology of diffuse large B-cell lymphoma (DLBCL), CD56+; H&E, 20x; (B) Strong expression of CD56 in DLBCL not otherwise specified (NOS) (tissue microarray), 4x; (C) weak to moderate CD56 expression in DLBCL NOS (tissue microarray), 4x.

Statistical analysis

For numeric and demographic variables descriptive statistics were used (median, range, standard deviation, percentage). Overall survival and progression-free survival were calculated using the Kaplan-Meier method. Statistical analyses were performed using IBM SPSS Statistics, version 26.

Results

Among 229 DLBCL, NOS cases included in the study, CD56 expression was found in 7 cases (3%). The intensity of CD56 expression varied from moderate focal to very intensive and diffuse positive reaction (Figure 1). Reanalysis of the five cases in which fine needle aspiration biopsy (FNAB) of the lymph node was performed prior to surgical biopsy and histological examination showed that CD56 was not included in routine flow-cytometry

TABLE 1. Clinicopathological characteristics of patients with CD56 positive diffuse large B-cell lymphoma (DLBCL), review of the literature with our series

Publication	Coun	No of pat	Case No	Sex Age	GC type**	Non-GC type	LN	Extranodal disease and site	Clinical stage	IPI	LDH	Surg	CT and No. of cycles	RT	Response	FU
Kern 1993 ²³	USA	1	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Muroi 1998 ⁷	Jap	2	1	M,49	Yes		Yes	Liver, Spleen, Pericard, Ef. Liver	NA	NA	NA	No	CHOP, NAx	No	PR?	NA
			2	F,62		Yes	Yes		NA	NA	NA	No	CHOP, NAx	No	PR?	NA
Sekita 1999 ¹⁸	Jap	1	1	M,16	Yes		Yes		I	NA	NA	No	CHOP, 6x	No	CR	10 m
Hammer 1998 ⁵	USA	4	1	M,51	NA	NA	Yes	Stomach	NA	NA	NA		NA	NA	NA	NA
			2	M,69	NA	NA	No		NA	NA	NA		NA	NA	NA	NA
			3	M,76	NA	NA	Yes		NA	NA	NA		NA	NA	NA	NA
			4	M,54	NA	NA	Yes		NA	NA	NA		NA	NA	NA	NA
Otsuka 2004 ⁴	Jap	2	1	NA	Yes		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
			2	NA	Yes		NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Weisberger *2006 ¹¹	USA	10	1	M,41	Yes	Yes	No	Ileocecal valve	NA	NA	NA	NA	NA	NA	NA	NA
			2	M,52	Yes		Yes	SpineAbdomen Brain	NA	NA	NA	NA	NA	NA	NA	NA
			3	M,54	Yes		Yes		NA	NA	NA	NA	NA	NA	NA	NA
			4	M,83	Yes		No		NA	NA	NA	NA	NA	NA	NA	NA
			5	M,49	Yes		Yes		NA	NA	NA	NA	NA	NA	NA	NA
			6	F,57	Yes		No		NA	NA	NA	NA	NA	NA	NA	NA
			7	F,69	Yes		Yes		NA	NA	NA	NA	NA	NA	NA	NA
			8	M,77	Yes		Yes		NA	NA	NA	NA	NA	NA	NA	NA
			9	M,84	Yes		Yes		NA	NA	NA	NA	NA	NA	NA	NA
			10	M,77	Yes		No		NA	NA	NA	NA	NA	NA	NA	NA
Isobe 2007 ¹³	Jap	3	1	M,80	Yes		Yes	Ascites	NA	NA	NA	No	THP-COP, 3x	No	NR	DOD
			2	F,87	Yes		No	Ileum	NA	NA	NA	Yes	No	No	CR	22 m
			3	M,73	Yes		Yes	Ileum	NA	NA	NA	Yes	R-CHOP, 6x	No	CR	22 m
Chen 2010 ⁴	Ch	1	1	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Gomyo 2010 ⁶	Jap	7	1	M,29		Yes	Yes	Spleen	IIIB	HI	↑	No	R-CHOP, aPBSCT	No	CR	A, 24 m
			2	F,60		Yes	Yes	WR	IIA	L	N	No	R-CHOP 3x	Yes	CR	A, 50 m
			3	F,22	Yes		No	WR	IA	L	N	No	CHOP 3x	Yes	CR	A, 57 m
			4	M,64		Yes	Yes	Pl. Ef, Adr. gl, Submand. gl	IIIA	H	↑	No	CHOP 5x	No	CR	D, 4 m
			5	M,63	Yes		No	Nasal cavity	IA	L	N	No	RCHOP 3x	Yes	CR	(pneumonia)
			6	M,50	Yes		No	Intra-extradural mass	IA	L	N	No	Res+CHOP 4x	Ye	RCR	A, 43 m
			7	F,45	Yes		No	Subcutis	IWA	HI	↑	No	R-CHOP 8x	sNo		A, 70 mA, 5 m
Stacchini 2012 ²	It	5	1	M,72	Yes		Yes	Spleen, Stomach, Pancr.	NA	NA	NA	NA	NA	NA	NA	NA
			2	M,15	Yes		Yes	Stomach, Liver	NA	NA	NA	NA	NA	NA	NA	NA
			3	M,71	Yes		No	Nasopharynx	NA	NA	NA	NA	NA	NA	NA	NA
			4	M,60					NA	NA	NA	NA	NA	NA	NA	NA
			5	M,21	Yes	Yes	No		NA	NA	NA	NA	NA	NA	NA	AWD 12m
Gu 2013 ¹⁰	SK	1	1	F,5		Yes		WR	I		N	Yes	COPAD, 6x	No	CR	NA
Liu 2020 ⁸	Ch	1	1	M,14	Yes, DH		Yes	Nasopharynx	IV	NA	↑	No	CTX+CP R-Hyper-CVAD AB R-DA-EPOCH, 6x IT DM+CTB, 4x	No	CR	NA
Gasljevic 2022	Slo	7	1	F,56	Yes		Yes	Skeletal muscle	IA	0	N	Yes	R-CHOP, 3x	No	CR	A, 63 m
			2	F,51	Yes		Yes	Small bowel	IA	0	N	No	R-CHOP, 3x	No	CR	A, 73 m
			3	M,57	Yes				IIA	0	↑	No	R-CHOP, 6x	No	CR	A, 55 m
			4	M,56		Yes, DE		Spleen, Liver, Adrenal gland	IVB	0	↑	No	R-EPOCH, 6x +IT, 2x	No	CR	A, 40 m
			5	F,53	Yes		Yes		IIA	3	N	Yes	CHOP, 3x	Yes	CR	A, 62 m
			6	F,30		Yes	Yes		IVB	3	↑	No	R-CHOP, 8x	Yes	CR	A, 182 m
			7	F,79	NA	NA	Yes		IVB	5	↑	No	No	No	NA	DOD

A = alive; aPBSCT = autologous peripheral blood stem cell transplantation; AWD = alive with disease; Ch = China; CHOP = cyclophosphamide, doxorubicin hydrochloride, vincristine sulfat, prednisone; Coun = country; CP = prednisone; CR = complete response; CT = chemotherapy; CTX = cyclophosphamide; D = dead; DA-EPOCH = etoposide, doxorubicin, vindesine, dexamethasone, cyclophosphamide; DE = double expressor; DOD = dead of disease; F = female; FU = follow-up; Gl = gland; Hyper-CVAD AB = A: cyclophosphamide, vindesine, liposomal doxorubicin, dexamethason, B: methotrexate, cytarabine; IPI = International Prognostic Index; It = Italy; IT = intrathecal; Jap = Japan; LN = lymph nodes; M = male; m = months; N = normal; NA = not available; NR = no response; Pancr = pancreas; Pl. E = pleural effusion; PR = partial response; R = rituximab; res = resection; RT = radiotherapy; SK = South Korea; Slo = Slovenia; Submand = submandibular; THP-COP = pirarubicin, cyclophosphamide, vincristine sulfat, prednisone; WR = Waldeyers ring

* only histologically proven cases are considered

** on the basis of the CD10 positivity

work-out. There was only one case²³ (case 1 in Table 1) in which immunocytochemistry for CD56 was stained since tumour cells showed co-expression of cytokeratin and the diagnosis of metastatic neuroendocrine carcinoma has been made.

All patients had *de novo* DLBCL. The median age at the time of diagnosis was 54.5 years (range 30–57). Five of them were women and 2 males. Five patients were diagnosed with DLBCL, GC type, 2 with DLBCL non-GC (ABC) type, one being a dou-

TABLE 2. Genetic profile of CD56 positive diffuse large B-cell lymphoma (DLBCL) samples

Case number in Table 2	COO IHC	COO AFPL	fusion	variants			VAF (%)	variant classification	Schmitz et al., 2018 ³² classification
				gene	nucleotide change	amino acid change			
1	GCB	GCB	ND	RANBP1	NM_002882.3:c.23A>G	NP_002873.1:p.[His8Arg]	13,7	Uncertain significance	unclassified
2	GCB	GCB	ND	ND	ND	ND	ND	ND	unclassified
3	GCB	GCB	ND	CD79B	NM_000626.2:c.587A>T	NP_000617.1:p.[Tyr196Phe]	49,0	Pathogenic	E2B
				CD79B	NM_000626.2:c.568A>G	NP_000617.1:p.[Met190Val]	50,1	Uncertain significance	
				EZH2	NM_001203247.1:c.1922A>G	NP_001190176.1:p.[Tyr641Cys]	53,7	Pathogenic	
				MYD88	NM_001172567.1:c.656C>G	NP_001166038.1:p.[Ser219Cys]	37,7	Uncertain significance	
				SH3BP5	NM_004844.4:c.460G>A	NP_004835.2:p.[Ala154Thr]	19,3	Uncertain significance	
4	ABC	ABC	IGH-BCL6	CD79B	NM_000626.2:c.587A>C	NP_000617.1:p.[Tyr196Ser]	25,9	Pathogenic	BN2
				SH3BP5	NM_004844.4:c.460G>A	NP_004835.2:p.[Ala154Thr]	12,6	Uncertain significance	

AFPL = Archer SusionPlex lymphoma; COO= cell of origin; IHC = immunohistochemical analyses; ND = not detected; VAF = variant allele frequency;

ble expressor (DE). One patient refused staging and treatment and died shortly after being diagnosed and was therefore excluded from survival analysis.

Among the six patients who received treatment, three patients were in clinical stage 1, one in stage 2 while two were in clinical stage 4. Only patients in clinical stage 4 had constitutional symptoms. Four patients had disease localised in the lymph nodes while two of them also had extranodal infiltrates – one in the pectoral muscles and the other in the renal fascia and small bowel. Three patients had elevated LDH levels, in fact, both patients in clinical stage 4B and one in stage 2A. Those patients in stage 4 had the IPI score 3 and others had the IPI score 0.

Three patients underwent surgical procedure and were later treated with adjuvant 3 cycles of CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) and R-CHOP (rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone). Other 3 patients were treated with 6 or 8 cycles of R-CHOP. Two patients were also treated with adjuvant radiotherapy after completion of systemic treatment. The patient with non-GC type DE of DLBCL was treated with 6 cycles of R-EPOCH (rituximab, etoposide, cyclophosphamide, doxorubicin, vincristine, prednisone) together with 2 doses of intrathecally administered methotrexate and cytosine arabinoside for central nervous system prophylaxis.

The 6 treated patients reached a complete response and did not experience progression of the disease during the follow-up period, meaning that

5-year PFS and OS are 100%. Median follow-up was 80.5 months (range 42–197).

The clinicopathological characteristics of our cohort together with all cases reported in the literature are shown in Table 1. Genetic profiling of 4 patients was performed as described in *Patients and methods*, and the results are presented in Table 2.

Discussion

CD56 expression in DLBCL NOS is very rare. Its incidence is reported to be 0.5 to 7% of DLBCLs, but is actually unknown since CD56 is generally not included in the immunohistochemical or flow cytometric panel for the diagnosis of DLBCL.¹⁰⁻¹⁸ In our series of patients with DLBCL NOS expression of CD56 was present in 3% of patients and varied in intensity from weak to very strong and diffuse. In one of those cases, that phenomenon resulted in an incorrect diagnosis of lymph node metastasis of the neuroendocrine tumour. In fact, in the general pathology service the main use of CD56 is to prove neuroblastoma and neuroendocrine differentiation in tumours of different origin while in hematopathology service it is used as a marker of NK cells, as a proof of clonal plasma cell proliferation, and as a prognostic marker in plasmacytoma, acute myeloid leukemia (AML), and acute lymphoblastic leukemia (ALL).²⁻⁵ Since neuroendocrine carcinomas could be unevenly and weakly positive or even negative for cytokeratins²⁴, it is of the greatest importance for the pathologist to be aware that strong expression of CD56 could be present al-

so in some entities that are by definition not CD56 positive.

Throughout the papers published so far, there has been much speculation about this phenomenon with regard to its expression in special clinicopathological settings and its possible prognostic value. From an epidemiological point of view, some authors⁹ suggested that it could be related to racial and/or geographical factors since, at the time of the publication of the paper, almost 50% of all reported cases were reported from Japan. Thorough analysis of all the cases with available information shows that 18 out of 45 cases (40%) have arisen in the population of far east (Japan, Korea, China; Table 1), while 27 (60%) were reported in the western population, Caucasians mainly (USA, Italy, Slovenia; Table 1). These results suggest that CD56+DLBCL is not related to racial / ethnic factors opposite to some other CD56 positive lymphoproliferative diseases such as NK/T cell lymphoma, nasal type.² The age distribution is very wide with cases described in paediatric/ adolescent population as well as in the older patient most of the patients being in 6–7th decade of life. In our series, the vast majority of patients were middle aged, in the beginning of the sixth decade. The distribution of gender showed that among the far east patients, somewhat higher number of men are reported (6 female *vs.* 9 males; for 3 cases there is no information about gender) while in the western world there is a predominance of males (7 females *vs.* 19 males; 26% *vs.* 74%). However, our series shows contradictory results in which most patients (70%) are women, so it can be assumed that the higher incidence reported in males so far could be only a mere coincidence.

There are two main biologically distinct molecular subtypes of DLBCL: GCB and ABC. ABC DLBCL is associated with substantially worse outcomes when treated with standard chemoimmunotherapy. Based on gene expression studies, Hans *et al.*²⁵ developed an algorithm to discriminate GBC from non-GBC types in regard to immunohistochemical expression of CD10, bcl6 and MUM1 with cutoff of 30%. In addition to GCB and ABC subtypes, double-hit lymphomas and double-expressor lymphomas, which overexpress myc and bcl2 protein, are aggressive DLBCLs and are also associated with a poor prognosis. On the basis of immunohistochemical results, a few authors^{9,11-13} found a relation of CD56 expression to DLBCL of GBC origin. Of the 45 summarized cases, for 8 cases there was no information about immunophenotype. Twenty-eight out of 36 (76%) were of GBC

type and the remaining 24% were of non-GBC (ABC) type. One reported case⁸ was double hit lymphoma with translocations of *MYC* and *bcl-6*, while in our series one DLBCL of non-GC (ABC) type DLBCL showed so-called double expressor profile with expression of bcl2 and myc protein expression being > 30%. Somewhat lower percentage of GBC types are reported in Eastern patients compared to the Western (10/15 and 17/21 or 75% *vs.* 81%). This finding could be related to the previously recognized and reported lower frequency of the DLBCL GBC subtype in Asian countries.²⁶

In addition, it has been suggested that CD56 expression in DLBCL could be related to a more frequent extranodal presentation associated to the adhesive properties of CD56.^{9,11} In neural cells, it mediates cell-to-cell adhesion by CD56 molecules of adjacent cells binding together.²⁷ It may be involved in homophilic adhesion for NK and T cells due to the C2-set Ig regions and fibronectin regions within its extracellular domain.²⁸ However, its function with respect to B-cell ontogeny is unclear. The expression of CD56 has been detected in a human pluripotent stem cell.²⁸ A subset of very early precursor B cells has the innate capacity for CD56 expression that is down-regulated and extinguished later in differentiation. It has been shown that lymphomagenesis is a stepwise process progression of which is enabled by accumulation of genetic events.⁸ In follicular or mantle cell lymphoma, for example³⁰, first events such as t(14,18) and t(11,14) namely, do occur in progenitor B cells. Drawing parallels to this, we could assume that CD56+ DLBCL could arise from the precursor B-cell that, for whatever reason, did not down-regulate CD56 expression and then collected additional mutations that resulted in lymphoma development. Some authors^{9,11-13} underlined frequent extranodal infiltrates in CD56+DLBCL with spleen, stomach, ileum, and nasal cavity being most frequently involved. Of 40 cases with available information, 16 (40%) presented with isolated lymphadenopathy while 24 (60%) had extranodal infiltrates with or without lymphadenopathy (14 *vs.* 10). Four of our patients presented with isolated lymphadenopathy while two had extranodal disease, which is concordant with majority of our patients having limited stage disease and were therefore treated adjuvantly after surgery.

The expression of CD 56 can be used as a prognostic marker in certain hematopathological entities; it can predict the occurrence of brain infiltration in ALL⁵, the aggressiveness of multiple myeloma³, and relapsed AML.⁴ So far, its prognostic

importance in DLBCL has not been confirmed. All of our patients achieved complete remission, and remained in remission which can be at least partially attributed to low IPI scores and low clinical stages; however, two patients with clinical stage 4 also achieved and maintained complete remission. None of our patients had a high IPI score of 4 or 5 which are known to have the lowest survival.³¹ In most of them, DLBCL was of GCB subtype, which also carry a better prognosis.²⁴

Schmitz *et al.*³² classified DLBCL cases according to genetic findings into 4 categories, namely MCD (based on the cooccurrence of *MYD88*^{L265P} and *CD79B* mutations), BN2 (based on *bcl6* fusions and *NOTCH2* mutations), N1 (based on *NOTCH1* mutations) and the EZB group (based on *EZH2* mutations and *bcl2* translocations). These subtypes differed phenotypically and in response to immunochemotherapy, with favourable survival in the BN2 and EZB groups. Genetic profiling of four patients from our series according to Schmitz classification³², showed that 1 case was of BN2 subtype, one belongs to the EZB group, while two were unclassified. Although data are limited and demand testing in larger cohorts of patients, so far it can be concluded that CD56 expression is more often present in cases of DLBCL NOS with prognostically favourable genetic findings.

CD56 is expressed in some aggressive tumour types such as small lung cell carcinoma and neuroblastoma. To date, it has been used as a target molecule for antibody-based immunotherapy in phase I and II clinical trials for small cell lung carcinoma³³; a favourable safety profile has been demonstrated. That led to the development of CAR-T therapy directed against CD56 in neuroblastoma. In the xenograft neuroblastoma model, anti-CD56 therapy led to the tumour burden control but had only modest effect on survival.³⁴ More studies are needed in regard to neuroblastoma therapy and other CD56 positive tumours but CD56 could eventually serve as a potential target for the treatment of CD56+ DLBCL patients who do not respond to the standard therapeutic schemes.

In conclusion, here we report one of the largest series of CD56+DLBCL with detailed clinicopathological data and for the first time described genetic findings in a limited number of patients. Our results show that CD56 expression is rare but seems to be present in prognostic favourable subtypes of DLBCL NOS as tested by immunohistochemical or genetic profiling.

Acknowledgement

The publication of this article was financed by the Slovenian Research Agency (ARRS), grant No. P3-0289. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Authors' contributions: GG and VKP conceived the idea, designed the study, collected the data, and wrote the initial draught of the article. LB and BJN collected and analysed the data and revised the manuscript. SN and VŠD performed the molecular testing, interpreted the data, and revised the manuscript.

References

1. Van Acker H, Capsomidis A, Smits EL, Van Tendeloo VF. CD56 in the immune system: more than a marker for cytotoxicity? *Front Immunol* 2017; **8**: 892-90. doi: 10.3389/fimmu.2017.00892
2. Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, et al. *WHO classification of tumours of hematopoietic and lymphoid tissues*. 4th edition. Lyon: IARC; 2017.
3. Chang H, Samiee S, Yi QL. Prognostic relevance of CD56 expression in multiple myeloma: a study including 107 cases treated with high-dose melphalan-based chemotherapy and autologous stem cell transplant. *Leuk Lymphoma* 2006; **47**: 43-7. doi: 10.1080/10428190500272549
4. Chaudhri NA, Almhareb F, Walter CU, Nounou R, Khalil S, Bakshi N, et al. Expression of CD56 in acute myeloid leukemia (AML) is associated with poor outcome when patients treated with stem cell transplant in second remission but not in the first remission. *Blood* 2011; **118**: 4880-86. doi: 10.1182/blood.V118.21.4880.4880
5. Ravandi F, Cortes J, Estrov Z, Thomas D. CD56 expression predicts occurrence of CNS disease in acute lymphoblastic leukaemia. *Leuk Res* 2002; **26**: 643-49. doi: 10.1016/s0145-2126(01)00188-6
6. Grimma KE, O'Malley DP. Aggressive B cell lymphomas in the 2017 revised WHO classification of tumors of hematopoietic and lymphoid tissues. *Ann Diagn Pathol* 2019; **38**: 6-10. doi: 10.1016/j.anndiagpath.2018.09.014
7. Feugier P, Van Hoof A, Sebban C, Solal-Celigny P, Bouabdallah R, Fermé C, et al. Long-term results of the R-CHOP study in the treatment of elderly patients with diffuse large B-cell lymphoma: a study by the group d'Etude des Lymphomes de l'Adulte. *J Clin Oncol* 2005; **23**: 4117-26. doi: 10.1200/JCO.2005.09.131
8. Liu Y, Barta SK. Diffuse large B-cell lymphoma: 2019 update on diagnosis, risk stratification and treatment. *Am J Hematol* 2019; **94**: 604-16. doi: 10.1002/ajh.25460
9. Gomyo H, Kajimoto K, Miyata Y, Maeda A, Mizuno I, Yamamoto K, et al. CD56-positive diffuse large B-cell lymphoma: possible association with extranodal involvement and bcl-6 expression. *Hematology* 2010; **15**: 157-61. doi: 10.1179/102453309X12583347113573
10. Gu MJ, Ha JO. CD56 positive diffuse large B-cell lymphoma: a case report and literature review. *Int J Clin Exp Pathol* 2013; **6**: 3023-25.
11. Weisberger J, Gorczyca W, Kinney MC. CD56-Positive Large B-cell Lymphoma. *App Imm Mol Morphol* 2007; **14**: 369-74. doi: 10.1097/01.pai.0000208279.66189.43
12. Stacchini A, Barreca A, Demurtas A, Aliberti S, di Celle PF, Novero D. Flow cytometric detection and quantification of CD56 (neural cell adhesion molecule, NCAM) expression in diffuse large B cell lymphomas and review of the literature. *Histopathology* 2012; **60**: 452-59. doi: 10.1111/j.1365-2559.2011.04098.x
13. Isobe Y, Sugimoto K, Takeuchi K, Ando J, Masuda A, Mori T, et al. Neural cell adhesion molecule (CD56)-positive B-cell lymphoma. *Eur J Haematol* 2007; **79**: 166-9. doi: 10.1111/j.1600-0609.2007.00893.x

14. Otsuka M, Yakushijin Y, Hamada M, Hato T, Yasukawa M, Fujita S. Role of CD21 antigen in diffuse large-B-cell lymphoma and its clinical significance. *Br J Hematol* 2004; **127**: 416-24. doi: 10.1111/j.1365-2141.2004.05226.x
15. Hammer RD, Vnencak-Jones CL, Manning B, Glick AD, Kinney MC. Microvillus lymphomas are B-cell neoplasms that frequently express CD56. *Mod Pathol* 1998; **11**: 239-46. PMID: 9521469
16. Chen B, Sun WY, Zhang G. [CD56 positive diffuse large B-cell lymphoma: report of a case]. [Chinese]. *Zhonghua Bing Li Xue Za Zhi* 2010; **39**: 343-4. PMID: 20654160
17. Muroi K, Omine K, Kuribara R, Uchida M, Izumi T, Hatake K, et al. CD56 expression in B-cell lymphoma. *Leuk Res* 1998; **22**: 201-2. doi: 10.1016/s0145-2126(97)00153-7
18. Sekita T, Tamaru J, Isobe K, Harigaya K, Masuoka S, Katayama T, et al. Diffuse large B-cell lymphoma expressing the natural killer cell marker CD56. *Pathol Int* 1999; **49**: 752-8. doi: 10.1046/j.1440-1827.1999.00929.x
19. Boltezar L, Kloboves-Prevodnik V, Pohar-Perme M, Gasljevic G, Jezersek-Novakovic B. Comparison of the algorithms classifying the ABC and GCB subtypes in diffuse large B-cell lymphoma. *Oncol Lett* 2018; **15**: 6903-12. doi: 10.3892/ol.2018.8243
20. Brozic A, Pohar Marinsek Z, Novakovic S, Kloboves-Prevodnik V. Inconclusive flow cytometric surface light chain results; can cytoplasmic light chains, Bcl-2 expression and PCR clonality analysis improve accuracy of cytological diagnoses in B-cell lymphomas? *Diagn Pathol* 2015; **10**: 191-6. doi: 10.1186/s13000-015-0427-5
21. Crotty R, Hu K, Stevenson K, Pontius MY, Sohani AR, Ryan RJH, et al. Simultaneous identification of cell of origin, translocations, and hotspot mutations in diffuse large B-cell lymphoma using a single RNA-sequencing assay. *Am J Clin Pathol* 2021; **155**: 748-54. doi: 10.1093/ajcp/aqaa185
22. Mokánszki A, Bicskó R, Gergely L, Méhes G. Cell-free total nucleic acid-based genotyping of aggressive lymphoma: Comprehensive analysis of gene fusions and nucleotide variants by next-generation sequencing. *Cancers* 2021; **13**: 3032. doi: 10.3390/cancers13123032
23. Kern WF, Spier CM, Miller TP, Grogan TM. NCAM (CD56)-positive malignant lymphoma. *Leuk Lymphoma* 1993; **12**: 1-10. doi: 10.3109/10428199309059565
24. Kirpatrick D, Swalling A, Kasmeridis H, Farshid G. Metastatic neuroendocrine carcinoma negative for cytokeratin immunohistochemistry. *Pathology* 2017; **51**: S140-S141. doi: 10.1016/j.pathol.2016.09.057
25. Hans CP, Weisenburger DD, Greiner TC, Gascoyne RD, Delabie J, Ott G, et al. Confirmation of the molecular classification of diffuse large B-cell lymphoma by immunohistochemistry using a tissue microarray. *Blood* 2004; **103**: 275-82. doi: 10.1182/blood-2003-05-1545
26. Shiozawa E, Yamochi-Onizuka T, Takimoto M, Ota H. The GCB subtype of diffuse large B-cell lymphoma is less frequent in Asian countries. *Leuk Res* 2007; **31**: 1579-83. doi: 10.1016/j.leukres.2007.03.017
27. Rutishauser U, Acheson A, Hall AK, Mann DM, Sunshine J. The neural adhesion molecule (NCAM) as a regulator of cell-cell interactions. *Science* 1988; **240**: 53-7. doi: 10.1126/science.3281256
28. Lanier LL, Chang C, Azuma M, Ruitenberg JJ, Hemperly JJ, Phillips JH. Molecular and functional analysis of human natural killer cell-associated neural cell adhesion molecule (N-CMCD56). *J Immunol* 1991; **146**: 4421-26.
29. Young HE, Steele TA, Bray RA, Detmer K, Blake LW, Lucas PW, et al. Human pluripotent and progenitor cells display cell surface cluster differentiation markers CD10, CD13, CD56, and MHC class-I. *Proc Soc Exp Biol Med* 1999; **221**: 63-71. doi: 10.1046/j.1525-1373.1999.d01-55.x
30. Navarrete M, Opezzo P. The pathogenesis of follicular lymphoma beyond apoptosis resistance. *Trans Canc Res* 2017; **6**: S529-S532. doi: 10.21037/tcr.201
31. Ziepert M, Hasenclever D, Kuhnt E, Glass B, Schmitz N, Pfreundschuh M, et al. Standard International prognostic index remains a valid predictor of outcome for patients with aggressive CD20+ B-cell lymphoma in the rituximab era. *J Clin Oncol* 2010; **28**: 2373-80. doi: 10.1200/JCO.2009.26.2493
32. Schmitz R, Wright GW, Huang DW, Johnson CA, Phelan JD, Wang JQ, et al. Genetics and pathogenesis of diffuse large B-cell lymphoma. *NEJM* 2018; **378**: 1396-407. doi: 10.1056/NEJMoa1801445.
33. Shah MH, Lorigan P, O'Brien MER, Fossella FV, Moore KN, Bhatia S, et al. Phase I study of IMG901, a CD56-targeting antibody- drug conjugate, in patients with CD56-positive solid tumours. *Invest New Drugs* 2016; **34**: 290-99. doi: 10.1007/s10637-016-0336-9.
34. Crossland DL, Denning WL, Ang S, Olivares S, Mi T, Switzer K, et al. Antitumor activity of CD56-chimeric antigen receptor T cells in neuroblastoma and SLCL models. *Oncogene* 2018; **37**: 3686-97. doi:10.1038/s41388-018-0187-2