

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 32, 2022, 1





ANNALES

**Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies**

Series Historia Naturalis, 32, 2022, 1

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 32, leto 2022, številka 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnsič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačič (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / *Società storica del Litorale - Capodistria*®Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / *Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente*®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / *Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran*
SI-6330 Piran / *Pirano*, Fornace/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 30. 06. 2022.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS) in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:** 300 izvodov/copie/copiesRevija *Annales, Series Historia Naturalis* je vključena v naslednje podatkovne baze / *La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in:* BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).To delo je objavljeno pod licenco / *Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY-NC 4.0.*Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website https://zdjp.si/en/p/annalesshn/
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito https://zdjp.si/en/p/annalesshn/

VSEBINA / INDICE GENERALE / CONTENTS 2022(1)

SREDOZEMSKI MORSKI PSI
SQUALI MEDITERRANEI
MEDITERRANEAN SHARKS

Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ

Observations on Thresher Shark, *Alopias vulpinus* (Chondrichthyes: Alopiidae) from the Coast of Algeria (Southwestern Mediterranean Sea) 1
Opazovanja morskih lisic, Alopias vulpinus (Chondrichthyes: Alopiidae) ob alžirski obali (jugozahodno Sredozemsko morje)

Elif ÖZGÜR ÖZBEK & Hakan KABASAKAL

Notes on Smoothback Angel Shark, *Squatina oculata* (Squatiniiformes: Squatinidae) caught in the Gulf of Antalya 9
Zapis o pegastih sklatih, Squatina oculata (Squatiniiformes: Squatinidae), ujetih v Antalijskem zalivu

Alessandro PAGANO & Alessandro DE MADDALENA

Underwater Observations of the Rare Angular Roughshark *Oxynotus centrina* (Chondrichthyes: Squalidae) in the Waters of Santa Tecla (Sicily, Italy) 17
Podvodna opazovanja redkega morskega prašiča, Oxynotus centrina (Chondrichthyes: Squalidae) v vodah Sante Tecele (Sicilija, Italija)

Deniz ERGÜDEN, Deniz AYAS & Hakan KABASAKAL

Morphometric Measurements of Three Young Carcharhinid Species from Northeastern Levant (Mediterranean Sea) 25
Morfometrične meritve mladičev treh vrst morskih psov iz družine Carcharhinidae iz severnovzhodnega Levanta (Sredozemsko morje)

Hakan KABASAKAL

Projections on the Future of Deep-Sea Sharks in the Sea of Marmara, Where Deep Zones Are Threatened by Deoxygenation: a Review 35
Napovedi o prihodnosti globomorskih morskih psov v Marmarskem morju, ogroženem zaradi pomanjkanja kisika: pregled

BIOINVAZIJA
BIOINVASIONE
BIOINVASION

Alan DEIDUN, Bruno ZAVA & Maria CORSINI-FOKA

Distribution Extension of *Lutjanus argentimaculatus* (Lutjanidae) and *Psenes pellucidus* (Nomeidae) to the Waters of Malta, Central Mediterranean Sea 49
Širjenje areala vrst Lutjanus argentimaculatus (Lutjanidae) in Psenes pellucidus (Nomeidae) v malteške vode (osrednje Sredozemsko morje)

Sami M. IBRAHIM, Abdulrazziq A. ABDULRAZIQ, Abdulghani ABDULGHANI, Sara A.A. AL MABRUK, David SALVATORI, Bruno ZAVA, Maria CORSINI-FOKA & Alan DEIDUN

First Record of *Enchelycore anatina* (Muraenidae) from Libyan Waters and an Additional Record from Southern Italy (Western Ionian Sea) 59
Prvi zapis o pojavljanju kavljazobe murene Enchelycore anatina (Muraenidae) iz libijskih voda in dodatni zapis za južno Italijo (zahodno Jonsko morje)

Rasha Ali HENEISH & Samir Ibrahim RIZKALLA

Morphometric and Meristic Characteristics of a New Record of Bluespot Mullet *Crenimugil seheli* (Pisces: Mugilidae) in Egyptian Mediterranean waters 67
Novi zapis o pojavljanju vrste Crenimugil seheli (Pisces: Mugilidae) v egiptovskih sredozemskih vodah in njene morfometrične in meristične značilnosti

Yana SOLIMAN, Adib SAAD, Vienna HAMMOUD & Christian CAPAPÉ

Heavy Metal Concentrations in Tissues of *Siganus rivulatus* (Siganidae) from the Syrian Coast (Eastern Mediterranean Sea) 75
Vsebnost težkih kovin v tkivih marmoriranega morskega kunca Siganus rivulatus (Siganidae) iz sirske obale (vzhodno Sredozemsko morje)

IHTIOLOGIJA
 ITTIOLOGIA
 ICHTHYOLOGY

Jihade ALAHYENE, Brahim CHIAHOU, Hammou EL HABOUZ & Abdelbasset BEN-BANI

Length Based Growth Estimation of the Blue Shark *Prionace glauca* from the Moroccan Central Atlantic Coast 85
Dolžinsko-masni odnos in ocena rasti pri sinjem morskem psu (Prionace glauca) iz osrednje atlantske obale Maroka

Okan AKYOL, Altan LÖK & Funda ERDEM

Occurrence of *Cubiceps gracilis* (Nomeidae) in the Eastern Mediterranean Sea 101
Pojavljanje klateža, Cubiceps gracilis (Nomeidae), v vzhodnem Sredozemskem morju

Farid HEMIDA, Boualem BRAHMI, Christian REYNAUD & Christian CAPAPÉ

Occurrence of the Rare Driftfish *Cubiceps gracilis* (Nomeidae) from the Algerian Coast (Southwestern Mediterranean Sea) 107
Pojavljanje redkega klazeža Cubiceps gracilis (Nomeidae) z alžirske obale (jugozahodno Sredozemsko morje)

Deniz ERGÜDEN & Cemal TURAN

A Rare Occurrence of *Carapus acus* (Carapidae) in the Eastern Mediterranean, Turkey 113
Redko pojavljanje strmorinca Carapus acus (Carapidae) v vzhodnem Sredozemskem morju (Turčija)

Laith JAWAD, Murat ŞIRIN, Miloslav PETRTÝL, Ahmet ÖKTENER, Murat ÇELİK & Audai QASIM

Skeletal Abnormalities in Four Fish Species Collected from the Sea of Marmara, Turkey 119
Skeletne anomalije pri štirih vrstah rib iz Marmarskega morja (Turčija)

RAZMNOŽEVALNA EKOLOGIJA
 ECOLOGIA RIPRODUTTIVA
 REPRODUCTIVE ECOLOGY

Amaria Latefa BOUZIANI, Khaled RAHMANI, Samira AIT DARNA, Alae Eddine BELMAHI, Sihem ABID KACHOUR & Mohamed BOUDERBALA

Gonadal Histology in *Diplodus vulgaris* from the West Algerian Coast 137
Histologija gonad pri navadnem šparu (Diplodus vulgaris) iz zahodne alžirske obale

Cheikhna Yero GANDEGA, Nassima EL OMRANI, Rezan O. RASHEED, Mohammed RAMDANI & Roger FLOWER

The Growth and Reproduction of Two Sparidae, *Pagrus caeruleostictus* and *Pagellus bellottii* in Northern Mauritanian Waters (Eastern Tropical Atlantic) 143
Rast in razmnoževanje dveh vrst pagrov, Pagrus caeruleostictus in Pagellus bellottii v severnih mavretanskih vodah (vzhodni tropski Atlantik)

Nassima EL OMRANI, Hammou EL HABOUZ, Abdellah BOUHAIMI, Jaouad ABOU OUALID, Abdellatif MOUKRIM, Jamila GOUZOULI, Mohammed RAMDANI, Roger FLOWER & Abdelbasset BEN-BANI

The Reproductive Biology of the Pouting *Trisopterus luscus* from the Atlantic Coast of Morocco 155
Reproduktivna biologija francoskega moliča (Trisopterus luscus) iz atlantske obale Maroka

Mourad CHÉRIF, Rimel BENMESSAOUD & Christian CAPAPÉ

Growth Patterns and Age Structure of *Mullus surmuletus* (Mullidae) from the Northern Coast of Tunisia (Central Mediterranean Sea) 173
Rastni parametri in starostna struktura progastih bradačev Mullus surmuletus (Mullidae) iz severne tunizijske obale (osrednje Sredozemsko morje)

FLORA
 FLORA
 FLORA

Martina ORLANDO-BONACA, Erik LIPEJ, Romina BONACA & Leon Lojze ZAMUDA

Improvement of the Ecological Status of the *Cymodocea nodosa* Meadow near the Port of Koper 185
Izboljšanje ekološkega stanja morskega travnika kolenčaste cimodoceje (Cymodocea nodosa) v bližini kopskega pristanišča

FAVNA
 FAVNA
 FAVNA

Manja ROGELJA, Martin VODOPIVEC & Alenka MALEJ

Cestum veneris Lesueur, 1813 (Ctenophora) – a Rare Guest in the Northern Adriatic Sea 197
Cestum veneris Lesueur, 1813 (Ctenophora) – redek gost v severnem Jadranu

Adla KAHRIĆ, Dalila DELIĆ & Dejan KULIJER

Notospermus annulatus (Nemertea: Lineidae), a New Record for Bosnia and Herzegovina 205
Notospermus annulatus (Nemertea: Lineidae), prvi zapis o pojavljanju za Bosno in Hercegovino

Andrea LOMBARDO & Giuliana MARLETTA

Report of an Interesting *Trapania* (Gastropoda: Nudibranchia: Goniodorididae) Specimen from Central Eastern Sicily 211
Zapis o zanimivem primerku iz rodu Trapania (Gastropoda: Nudibranchia: Goniodorididae) iz osrednje vzhodne Sicilije

Abdelkarim DERBALI & Othman JARBOUI

Stock Assessment, Cartography and Sexuality of the Wedge Clam *Donax trunculus* in the Gulf of Gabes (Tunisia) 217
Ocena staleža, kartografija in spolnost klinaste školjke Donax trunculus v gabeškem zalivu (Tunizija)

Abdelkarim DERBALI, Aymen HADJ TAIEB & Othman JARBOUI

Length-Weight Relationships and Density of Bivalve Species in the Shellfish Production Area of Zarzis (Tunisia, Central Mediterranean Sea) 229
Dolžinsko-masni odnos in gostota školjk na gojišču školjk v predelu Zarsisa (Tunizija, osrednje Sredozemsko morje)

Toni KOREN

The Diversity of Moths (Lepidoptera: Heterocera) of Significant Landscape Donji Kamenjak and Medulin Archipelago, Istria, Croatia 237
Raznolikost nočnih metuljev (Lepidoptera: Heterocera) Pomembne pokrajine Donji Kamenjak in Medulinski arhipelag, Istra, Hrvaška

OCENE IN POROČILA

RECENSIONI E RELAZIONI

REVIEWS AND REPORTS

Ines Mandić Mulec & Nives Ogrinc

Recenzija knjige: Mikrobna biogeokemija vod 263

Kazalo k slikam na ovitku 265

Index to images on the cover 265

IMPROVEMENT OF THE ECOLOGICAL STATUS OF THE *CYMODOCEA NODOSA* MEADOW NEAR THE PORT OF KOPER

Martina ORLANDO-BONACA

Marine Biology Station Piran, National Institute of Biology, SI-6330 Piran, Fornače 41, Slovenia
e-mail: martina.orlando@nib.si

Erik LIPEJ

Ulica XXX divizije 10, SI-6320 Portorož, Slovenia

Romina BONACA

Ulica Vena Pilona 5, SI-6000 Koper, Slovenia

Leon Lojze ZAMUDA

Marine Biology Station Piran, National Institute of Biology, SI-6330 Piran, Fornače 41, Slovenia

ABSTRACT

Seagrass beds are more or less the marine counterpart of tropical rainforests, and their health is related to different anthropogenic stressors, including navigation routes and port activities. In the Mediterranean Sea, Cymodocea nodosa is considered an effective indicator of environmental change, due to its universal distribution, sensitivity to various natural and anthropogenic pressures, and the measurability of the species' responses to impacts. The aim of this study is to present the changes in the assessment of the ecological status of the C. nodosa meadow near the port of Koper, which was evaluated as Bad in 2018. The results show a significant improvement in the ecological status of the meadow, which can be attributed to a reduction in anthropogenic stressors.

Key words: *Cymodocea nodosa*, status evaluation, MediSkew index, Port of Koper, northern Adriatic Sea

MIGLIORAMENTO DELLO STATO ECOLOGICO DELLA PRATERIA DI *CYMODOCEA NODOSA* VICINO AL PORTO DI CAPODISTRIA

SINTESI

Le praterie di fanerogame marine vengono considerate la controparte marina delle foreste pluviali tropicali, e la loro salute è legata a diversi fattori di stress antropogenici, tra cui le rotte di navigazione e le attività portuali. Nel Mediterraneo, Cymodocea nodosa è considerata un indicatore efficace del cambiamento ambientale, a causa della sua distribuzione universale, della sensibilità a varie pressioni naturali e antropogeniche, e della misurabilità delle risposte della specie agli impatti. Lo scopo di questo studio è di presentare i cambiamenti nella valutazione dello stato ecologico della prateria di C. nodosa vicino al porto di Capodistria, che è stato valutato come Cattivo nel 2018. I risultati mostrano un miglioramento significativo dello stato ecologico della prateria, che può essere attribuito a una riduzione dei fattori di stress antropogenici.

Parole chiave: *Cymodocea nodosa*, valutazione dello stato, indice MediSkew, Porto di Capodistria, Adriatico settentrionale

INTRODUCTION

Seagrass meadows are among the most productive environments in the seas and oceans worldwide (Spalding et al., 2003; Brodersen et al., 2018). They provide habitat niches, food, and protection from predators for many different organisms in lagoons and marine ecosystems (Hemminga & Duarte, 2000; Como et al., 2008; Tuya et al., 2014; Espino et al., 2015). These environments are also important for human well-being (Nordlund et al., 2018; Unsworth et al., 2018), as they provide a range of ecosystem services, including moderating wave action and thus protecting the coastline from erosion (Ondiviela et al., 2014; Cabaço & Rui Santos, 2014), stabilising sediments (Terrados & Borum 2004; Widdows et al., 2008), regulating nutrient cycling and sequestering carbon (Duarte et al., 2010; Luisetti et al., 2013), purification of seawater (Richir et al., 2013), and providing a system for education and research (Effrosynidis et al., 2018). For these reasons, they have been included as priority habitats in a number of legal regulations, including the European Habitats Directive (HD, 92/43/EEC).

Seagrass beds are more or less the marine counterpart of tropical rainforests, and their health is associated with various types of anthropogenic stressors. These pressures include navigation routes and port activities, seabed dredging, commercial and recreational activities such as fishing and mooring, runoff from urban and agricultural areas, wastewater, and more recently, increasing climate change and ocean acidification (Short et al., 2011; Tuya et al., 2002; Marbà et al., 2014; Orlando-Bonaca et al., 2015, 2019; Repolho et al., 2017). Such pressures affect light and nutrient resources (Hemminga & Duarte, 2000), and cause physical damage to different sea bottom types (Montefalcone et al., 2008; Marbà et al., 2014). Rapid and widespread declines in seagrass meadows have been reported from many coastal areas over the past fifteen years (Orth et al., 2006; Tuya et al., 2013; Fabbri et al., 2015). Seagrasses have disappeared at a rate of 110 km² per year since 1980, a value similar to the rates of loss described for mangroves, coral reefs, and tropical rainforests (Waycott et al., 2009). In terms of cover, one third of the world’s seagrass meadows are reported to have already disappeared (Waycott et al., 2009).

Four native seagrass species are found in the Adriatic Sea: *Posidonia oceanica* (Linnaeus) Delile, *Cymodocea nodosa* (Ucria) Ascherson, *Zostera marina* Linnaeus and *Zostera noltei* Hornemann (Lipej et al., 2006). In the Mediterranean Sea, *C. nodosa* is considered an effective indicator of environmental change, due to its universal distribution,

sensitivity to various natural and anthropogenic pressures, and the measurability of the species’ responses to these impacts (Orfanidis et al., 2007, 2010; Oliva et al., 2012; Orlando-Bonaca et al., 2015; Papathanasiou et al., 2016). Although *C. nodosa* exhibits great phenotypic plasticity and can adapt to various natural and anthropogenic stressors through physiological and morphological adaptations, a sharp decline has been reported in coastal areas (Orth et al., 2006; Short et al., 2011; Tuya et al., 2013, 2014; Fabbri et al., 2015; Mačić & Zordan, 2018; Najdek et al., 2020) in recent decades.

In the northern Adriatic Sea, there is still a lack of long-time data series to support the conservation status of *C. nodosa* meadows, which is included in Annex II (List of Endangered or Threatened species) of the Convention for the Protection of the Mediterranean Sea Against Pollution (the Barcelona Convention). The ecological status of *C. nodosa* meadows in the Gulf of Trieste was assessed using the MediSkew index (Orlando-Bonaca et al., 2015; 2016), which was developed in accordance with the requirements of the EU Water Framework Directive (WFD, 2000/60/EC) and the Marine Strategy Framework Directive (MSFD, 2008/56/EC). The ecological status of the *C. nodosa* meadow growing near the Port of Koper was first evaluated in 2018 (Orlando-Bonaca et al., 2019), and subsequently monitored in 2020 and 2021. An annual monitoring programme is planned for the future, as shipping routes and port activities are considered one of the main pressures on the

Tab. 1: Boundaries among status classes for the MediSkew index (classes High and Good indicate a Good Environmental Status).

Tab. 1: Meje med posameznimi razredi stanja za MediSkew indeks (razreda Zelo dobro in Dobro označujeta Dobro okoljsko stanje).

Status classes	Absolute values of MediSkew
High	0 ≤ MediSkew < 0.2
Good	0.2 ≤ MediSkew < 0.4
Moderate	0.4 ≤ MediSkew < 0.6
Poor	0.6 ≤ MediSkew < 0.8
Bad	0.8 ≤ MediSkew ≤ 1

status of *C. nodosa* meadows (Orlando-Bonaca et al., 2015). The aim of this study is to present the changes in the assessment of the ecological status of the *C. nodosa* meadow near the port of Koper from 2018 using the MediSkew index.

MATERIAL AND METHODS

Study area, fieldwork and laboratory work

The Port of Koper is a Slovenian multi-purpose port on the northern Adriatic Sea, mainly connecting markets in Central and South-eastern Europe with the Mediterranean Sea and the Far East. The marine part of the cargo port consists of tree basins,

associated mooring piers and specialized loading terminals. The highest water turbidity values were measured during manoeuvres of the large ships (Žagar et al., 2014). Dredging of the sedimentary bottom was carried out in the Port of Koper along the access channels to Basin I (Luka Koper, 2015). Moreover, construction works, including dredging for the construction of a new RORO berth in the Basin III, were officially opened on May 27, 2019, and completed on March 31, 2020 (Franka Cepak, pers. comm.), resulting in a high sedimentation/resuspension rate.

The seagrass meadow located near the Port of Koper was sampled in July 2018, 2020 and 2021. Two sites (LuKp1 and LuKp2) were selected (Fig.




Fig. 1: Map of sampling sites for *Cymodocea nodosa* near the Port of Koper (LuKp1 and LuKp2) and the reference site in the Moon Bay (Cy2). The site LuKp1 is located in the part of the *C. nodosa* meadow with a higher density, while the site LuKp2 is located in a less dense part of the meadow. Other colours on the map: the largest, light green circle = *Zostera noltei*; grey circles = boulders with *Dictyota dichotoma*, *Padina pavonica* and turf; red circle = boulders with *Cystoseira compressa* and *P. pavonica*; dark brown area = boulders with *P. pavonica*, *Halopteris scoparia* and turf; light brown area = boulders with *Padina pavonica* and turf.

Sl. 1: Zemljevid mest vzorčenja kolenčaste cimodoceje blizu Luke Koper (LuKp1 in LuKp2) in referenčno mesto vzorčenja v Mesečevem zalivu (Cy2). LuKp1 se nahaja v zelo gostem delu travnika kolenčaste cimodoceje, medtem ko se LuKp2 nahaja v manj gostem delu. Druge barve na karti: največji, svetlozeleni krog = *Zostera noltei*; sivi krogi = skale prekrte z vrstama *Dictyota dichotoma* in *Padina pavonica* ter turfom; rdeči krog = skale prekrte z vrstama *Cystoseira compressa* in *P. pavonica*; temno rjavo območje = skale prekrte z vrstama *P. pavonica* in *Halopteris scoparia* ter turfom; svetlo rjavo območje = skale prekrte z vrsto *Padina pavonica* in turfom.

Tab. 2: Statistic parameters (minimum, maximum, mean, median) and absolute value of skewness ($|G|$) of \ln -transformed lengths of photosynthetically active parts of *Cymodocea nodosa* leaves from the sampling areas near the Port of Koper (LuKp1 and LuKp2) in 2018, 2020 and 2021, and in Moon Bay (Cy2, Strunjan Nature Reserve) in 2018. The reference median value in 2018 was 13.95 cm.

Tab. 2: Statistični parametri (minimum, maksimum, povprečje, mediana) in absolutna vrednost koeficienta asimetrije ($|G|$) \ln -transformiranih dolžin fotosintetsko aktivnega dela listov kolenčaste cimodoceje (*C. nodosa*) na točkah vzorčenja blizu Luke Koper (LuKp1 in LuKp2) v 2018, 2020 in 2021 ter v Mesečevem zalivu (Cy2, Naravni rezervat Strunjan) v 2018. Referenčna mediana v 2018 je bila 13,95 cm.

Area	Date	Min length (cm)	Max length (cm)	Mean (cm)	Median (cm)	$ G $
Str_3	12.7.2018	5.4	30.5	14.5	13.95	0.261
Str_4	12.7.2018	8.1	22.7	13.5	13.20	0.022
LuKp1_1	17.7.2018	5.9	66.2	37.8	41.25	1.423
LuKp1_2	17.7.2018	6.0	57.1	34.7	37.05	1.162
LuKp2_1	17.7.2018	3.7	58.8	30.7	30.45	1.533
LuKp2_2	17.7.2018	6.9	52.2	27.3	28.25	1.130
LuKp1_1	14.7.2020	5.4	62.5	32.0	31.90	1.044
LuKp1_2	14.7.2020	7.4	57.7	29.9	29.25	0.706
LuKp2_1	14.7.2020	5.1	61.3	29.2	28.90	0.979
LuKp2_2	14.7.2020	7.3	55.9	31.4	31.25	0.955
LuKp1_1	1.7.2021	8.7	55.8	27.33	25.90	0.355
LuKp1_2	1.7.2021	7.3	57.1	28.12	27.20	0.442
LuKp2_1	1.7.2021	11.5	47.7	24.72	22.95	0.142
LuKp2_2	1.7.2021	5.7	46.2	24.15	23.15	0.659

1) along the same isobath (3 m) and, within each site, two areas (LuKp1_1, LuKp1_2, and LuKp2_1, LuKp2_2) were chosen, approximately 100 m apart. In each area, five metallic frames (25 cm x 25 cm) were randomly placed on the bottom by SCUBA divers. These five squares were considered replicates of one sample. All shoots of *C. nodosa* located in each frame were carefully uprooted. The samples were labelled and individually placed in plastic bags.

In July 2018, samples of *C. nodosa* were also collected in the Strunjan Nature Reserve (sampling site Cy2, areas Str_3 and Str_4). Due to the low Pressure Index for Seagrass Meadows (PISM) value, the area Str_3 was selected as the reference area for *C. nodosa* in the Gulf of Trieste in 2009 (Orlando-Bonaca et al., 2015), and it has to be sampled and assessed every 5 years.

The samples of *C. nodosa* were stored in a freezer at -20°C in the laboratory of the Marine

Tab. 3: MediSkew index values for the sampling areas of *Cymodocea nodosa* in the Port of Koper and in the Moon Bay (Strunjan) and assessment of the Ecological Status (according to the WFD) and Environmental Status (according to the MSFD).

Tab. 3: Vrednosti indeksa MediSkew na točkah vzorčenja s kolenčasto cimodocejo in opredelitev ekološkega stanja (glede na OVS) in okoljskega stanja (glede na ODMS) za morski travnik ob Luki Koper in v Mesečevem zalivu (Strunjan).

Year	Area	Area's MediSkew	Site's MediSkew	Meadow's MediSkew	Ecolog. Status	Environ. Status	N of leaves	N of adult leaves			
2018	Str_3	0.065	0.04	-	High	Good / Achieved	300	213			
	Str_4	0.024					300	218			
	LuKp1_1	1.00	0.935	0.825	Bad	Not good / Not achieved	300	225			
	LuKp1_2	0.87					300	204			
	LuKp2_1	0.79	0.715				300	247			
	LuKp2_2	0.64					300	218			
2020	LuKp1_1	0.71	0.635				0.640	Poor	Not good / Not achieved	251	181
	LuKp1_2	0.56								300	223
	LuKp2_1	0.62	0.645	300	246						
	LuKp2_2	0.67		300	222						
2021	LuKp1_1	0.39	0.415	0.37	Good	Good / Achieved	300	238			
	LuKp1_2	0.44					300	207			
	LuKp2_1	0.26	0.325				300	231			
	LuKp2_2	0.39					300	212			

Biology Station Piran. The day before the analysis, they were slowly defrosted in a refrigerator. Seagrass shoots were then kept in plastic wash basins containing seawater. Twenty shoots from each quadrat were randomly selected (Orfanidis *et al.*, 2007). For each leaf (usually 5-6 leaves per shoot), the following parameters were measured to the nearest mm: length of the leaf sheath, length of the photosynthetic part and its width. The age of the leaf was designated as adult (when the leaf sheath

was well-developed), intermediate (when the leaf sheath was weakly developed at the leaf base), and juvenile (when the leaf sheath was absent). The above measurements were made on at least 60 undamaged, photosynthetically active leaves (adult and/or intermediate) from each frame. One sample consisted of five replicates of 60 leaves (300 leaves in total).

Additionally, in May 2020, the meadow and other vegetation types in the area were checked by

applying a field method based on visual observation of sea-bottom segments covered with vegetation in the infralittoral belt. The survey consisted of a cruise along the coastline in a small boat. Sublittoral communities were identified using a large Aquascope Underwater Viewer and directly annotated in a graphic display. This graphic support was prepared at an appropriate small scale and was suitable for use in the field. The final result is a division of the shoreline into several sectors, each identified by a community category (see Fig. 1). The information obtained on the distribution of communities was transcribed into a georeferenced graphic support in a Geographical Information System. All vegetation types between 1 and 4 m depth, were mapped.

Data analysis

To quantify changes in the photosynthetic part of the leaf length distribution for each *C. nodosa* sampling area near the Port of Koper, the MediSkew index was calculated (for details, see Orlando-Bonaca *et al.*, 2015). The boundaries among the status classes for the MediSkew index were set equidistantly (Tab. 1). Five status classes are sufficient for the assessment of the Ecological Status (ES) according to the WFD. In addition, High and Good classes indicate Good Environmental Status (EnS) according to the MSFD, while the classes Moderate, Poor, and Bad are considered Not Good EnS.

RESULTS AND DISCUSSION

The surveyed *C. nodosa* meadow near the Port of Koper can be considered as a part of the biocoenosis of superficial muddy sands in sheltered waters. The part of the meadow closest to the Port of Koper has a higher density of shoots than the part to the north (different green colours in Fig. 1). Within the meadow, rocky biotopes were also found, which include small communities dominated by *Padina pavonica* (Linnaeus) Thivy, *Dictyota dichotoma* (Hudson) J.V. Lamouroux, *Halopteris scoparia* (Linnaeus) Sauvageau and *Cystoseira compressa* (Esper) Gerloff & Nizamuddin. A monospecific patch of *Zostera noltei* was also found close to the Port (see Fig. 1).

The parameters of *C. nodosa* per sampling area are shown in Table 2. The leaves of *C. nodosa* were significantly shorter in the areas within the reference site in the Moon Bay (Cy2) than in the areas near the Port of Koper in all years, and consequently so were the median values (Tab. 2). The skewness |G| was the highest in the LuKp2_1 area in 2018 (Tab. 2). However, the

results show that mean and median leaf length values decreased in all sampled areas in the Port of Koper in 2020 and additionally in 2021. Leaf lengths were still much longer than those at the reference area, but there is a very clear trend of decreasing leaf lengths since 2018 near the Port of Koper (Tab. 2).

It should be emphasized that in 2020, many adult leaves of *C. nodosa* at LuKp1_1 were broken, without apical parts, and therefore we could not measure 300 undamaged leaves for this area (Tab. 3), as indicated in the methodology. All samples collected in 2021 had fewer damaged leaves and the number of adult leaves of *C. nodosa* exceeded 200 per sample (Tab. 3).

The ES (according to WFD) and the EnS (according to MSFD) of sampling areas and sites were assessed according to the boundaries in Table 1. The MediSkew index values for each sampled area in the Port of Koper are presented in Table 3. The two areas of the sampling site LuKp2, furthest from the Port Basin III, improved the ES from Poor in 2018 and 2020 to Good in 2021.

The improvement in the status of the LuKp1 sampling site is also impressive (Tab. 3). The area LuKp1_2 was assessed as Bad ES in 2018, while it remained Moderate in 2020 and 2021. The area LuKp1_1 improved from Bad ES in 2018 to Poor in 2020 and to Good in 2021. The ES of the entire meadow of *C. nodosa* near the Port of Koper was evaluated as Good in 2021, which is two orders of magnitude better than in 2020 (Tab. 3).

The results obtained from 2018 to 2021 show a significant improvement in the ES of the *C. nodosa* meadow. The Good ES achieved in 2021 may be related to the reduction of anthropogenic pressures, as the construction of the new RORO berth was completed in March 2020. This construction resulted in higher sediment resuspension in recent years, leading to increased turbidity and consequently less light. Seagrasses are generally light-limited (Touchette & Burkholder, 2000). Thus, when exposed to low light levels due to high water turbidity, they respond by increasing biomass distribution to the leaves. The increase in leaf size allows marine plants to capture more light and convert it into photosynthetic production (Greve & Binzer, 2004). That resuspension of sediments and water turbidity are critical to the health of *C. nodosa* meadows is confirmed by recent research (Orfanidis *et al.*, 2020). Additionally, the decrease in anthropogenic pressures near the port area in 2020 was also influenced by the Covid-19 pandemic, which led to a decrease in maritime traffic, especially cruise ship traffic in the Port of Koper,

as reported in many local media. March et al. (2021) have attempted to assess the impact of the pandemic on maritime traffic globally, which in turn has implications for the blue economy and ocean health.

The results of the present study are very encouraging, and as the Port of Koper has prepared a long-term monitoring programme in the harbour area and its surroundings, we hope to confirm the improved ES of the *C. nodosa* meadow near the Port in the long term.

ACKNOWLEDGEMENTS

The authors are grateful to the Port of Koper that financially supported this study. We would like to thank also Lovrenc Lipej, Milijan Šiško, Tihomir Makovec, Borut Mavrič, Domen Trkov, Aljoša Gračner, Matej Marinac and Tristan Bartole for their help during the fieldwork and laboratory work. Special thanks are due to Milijan Šiško for the preparation of Figure 1. We thank also the reviewers for their careful reading of the manuscript and their constructive remarks.

IZBOLJŠANJE EKOLOŠKEGA STANJA MORSKEGA TRAVNIKA KOLENČASTE CIMODOCEJE (*CYMODOCEA NODOSA*) V BLIŽINI KOPRSKEGA PRISTANIŠČA

Martina ORLANDO-BONACA

Morska biološka postaja Piran, Nacionalni inštitut za biologijo, SI-6330 Piran, Fornače 41, Slovenia
e-mail: martina.orlando@nib.si

Erik LIPEJ

Ulica XXX divizije 10, SI-6320 Portorož, Slovenia

Romina BONACA

Ulica Vena Piloni 5, SI-6000 Koper, Slovenia

Leon Lojze ZAMUDA

Morska biološka postaja Piran, Nacionalni inštitut za biologijo, SI-6330 Piran, Fornače 41, Slovenia

POVZETEK

*Morski travniki so bolj ali manj morski ekvivalent tropskega deževnega gozda, njihovo zdravje pa je povezano z različnimi antropogenimi dejavniki, vključno s plovnimi potmi in pristaniškimi dejavnostmi. V Sredozemskem morju je kolenčasta cimodoceja (*Cymodocea nodosa*) zaradi svoje univerzalne razširjenosti, občutljivosti na različne naravne in antropogene pritiske ter merljivosti odzivov vrste na vplive, učinkovit kazalnik okoljskih sprememb. Namen študije je predstaviti spremembe v oceni ekološkega stanja morskega travnika kolenčaste cimodoceje v bližini koprškega pristanišča, ki je bilo 2018 opredeljeno kot Zelo slabo. Rezultati kažejo bistveno izboljšanje ekološkega stanja travnika, kar lahko pripišemo zmanjšanju antropogenih pritiskov.*

Ključne besede: *Cymodocea nodosa*, ocena stanja, MediSkew indeks, Luka Koper, severni Jadran

REFERENCES

- Brodersen, M.M., M. Pantazi, A. Kokkali, P. Panayotidis, V. Gerakaris, I. Maina, S. Kavadas, H. Kaberi & V. Vassilopoulou (2018):** Cumulative Impacts from Multiple Human Activities on Seagrass Meadows in Eastern Mediterranean Waters: The Case of Saronikos Gulf (Aegean Sea, Greece). *Environ. Sci. Pollut. Res.*, 25, 26809-26822. doi:10.1007/s11356-017-0848-7.
- Cabaço, S. & A. Rui Santos (2014):** Human-induced changes of the seagrass *Cymodocea nodosa* in Ria Formosa lagoon (Southern Portugal) after a decade. *Cah. Biol. Mar.*, 55, 101-108.
- Como, S., P. Magni, M. Baroli, D. Casu, G. De Falco & A. Floris (2008):** Comparative analysis of macrofaunal species richness and composition in *Posidonia oceanica*, *Cymodocea nodosa* and leaf litter beds. *Mar. Biol.*, 153(6), 1087-1101.
- Duarte, C.M., N. Marbà, E. Gacia, J.W. Fourqurean, J. Beggins, C. Barrón & E.T. Apostolaki (2010):** Seagrass community metabolism: Assessing the carbon sink capacity of seagrass meadows. *Global Biogeochem. Cy.*, 24, GB4032.
- Effrosynidis, D., A. Arampatzis & G. Sylaios (2018):** Seagrass detection in the Mediterranean: A supervised learning approach. *Ecol. Inform.*, 48, 158-170.
- Espino, F., A. Brito, R. Haroun & F. Tuya (2015):** Macroecological analysis of the fish fauna inhabiting *Cymodocea nodosa* seagrass meadows. *J. Fish Biol.*, 87(4), 1000-1018.
- Fabbri F., F. Espino, R. Herrera, L. Moro, R. Haroun, R. Riera, N. González-Henriquez, O. Bergasa, O. Monterroso, M. Ruiz de la Rosa & F. Tuya (2015):** Trends of the seagrass *Cymodocea nodosa* (Magnoliophyta) in the Canary Islands: population changes in the last two decades. *Scientia Marina*, 79(1), 7-13.
- Greve, T.M. & T. Binzer (2004):** Which factors regulate seagrass growth and distribution? In: J. Borum, C.M. Duarte, D. Krause-Jensen, & T.M. Greve (Eds). *European seagrasses: an introduction to monitoring and management. The M&MS project*: 19-23.
- Habitat Directive (1992):** Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora.
- Hemminga, M.A. & C.M.D. Duarte (2000):** Seagrass ecology. Cambridge University Press, 298 pp.
- Lipej, L., R. Turk & T. Makovec (2006):** Ogrožene vrste in habitatni tipi v slovenskem morju (Endangered species and habitat types in the Slovenian sea). Zavod RS za varstvo narave, Ljubljana, 264 pp.
- Luisetti, T., E.L. Jackson & R.K. Turner (2013):** Valuing the European 'coastal blue carbon' storage benefit. *Mar. Pollut. Bull.*, 71, 101-106.
- Luka Koper (2015):** More about port's history. Available at: <https://luka-kp.si/eng/more-about-port-s-history>.
- Mačić, V. & C. Zordan (2018):** Mapping of the *Cymodocea nodosa* (Ucria) Asch. meadows in the Kotor Bay and data comparison over the last four decades. *Studia Marina*, 31(1), 5-15.
- Marbà, N., E. Diaz-Almela & C.M. Duarte (2014):** Mediterranean seagrass (*Posidonia oceanica*) loss between 1842 and 2009. *Biol. Conserv.*, 176, 183-190.
- March, D., K. Metcalfe, J. Tintoré & B.J. Godley (2021):** Tracking the global reduction of marine traffic during the COVID-19 pandemic. *Nature Communications*, 12, 2415. <https://doi.org/10.1038/s41467-021-22423-6>.
- Marine Strategy Framework Directive (2008):** Directive 2008/56/EC of the European Parliament and of the Council of 17 June 2008 establishing a framework for community action in the field of marine environmental policy.
- Montefalcone, M., M. Chiantore, A. Lanzone, C. Morri & G. Albertelli (2008):** BACI design reveals the decline of the seagrass *Posidonia oceanica* induced by anchoring. *Mar. Pollut. Bull.*, 56(9), 1637-1645.
- Najdek, M., M. Korlević, P. Paliaga, M. Markovski, I. Ivančić, L. Iveša, I. Felja & G.J. Herndl (2020):** Dynamics of environmental conditions during the decline of a *Cymodocea nodosa* meadow. *Biogeosciences*, 17, 3299-3315. <https://doi.org/10.5194/bg-17-3299-2020>.
- Nikolić, V., A. Žuljević, L. Mangialajo, B. Antolić, G. Kušpilić & E. Ballesteros (2013):** Cartography of Littoral Rocky-Shore Communities (CARLIT) as a Tool for Ecological Quality Assessment of Coastal Waters in the Eastern Adriatic Sea. *Ecol. Indic.*, 34, 87-93. doi:10.1016/j.ecolind.2013.04.021.
- Nordlund, L.M., R.K.F. Unsworth, M. Gullstrom & L.C. Cullen-Unsworth (2018):** Global significance of seagrass fishery activity. *Fish and Fisheries*, 19(3), 399-412. <https://doi.org/10.1111/faf.12259>.
- Oliva, S., O. Mascaró, I. Llagostera, M. Pérez, & J. Romero (2012):** Selection of Metrics Based on the Seagrass *Cymodocea nodosa* and Development of a Biotic Index (Cymox) for Assessing Ecological Status of Coastal and Transitional Waters. *Estuar. Coast. Shelf S.*, 114, 7-17.
- Ondiviela, B., I.J. Losada, J.L. Lara, M. Maza, C. Galván, T.J. Bouma & J. van Belzen (2014):** The role of seagrasses in coastal protection in a changing climate. *Coast. Engineering*, 87, 158-168.
- Orfanidis, S., V. Papathanasiou & S. Gounaris (2007):** Body size descriptor of *Cymodocea nodosa* indicates anthropogenic stress in coastal ecosystem. *Transitional Waters Bulletin*, 2, 1-7.
- Orfanidis, S., V. Papathanasiou, S. Gounaris, & T. Theodosiou (2010):** Size distribution approaches for monitoring and conservation of coastal *Cymodocea* habitats. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 20, 177-188.

- Orfanidis, S., V. Papathanasiou, N. Mittas, T. Theodosiou, A. Ramfos, S. Tsioli, M. Kosmidou, A. Kafas, A. Mystikou & A. Papadimitriou (2020):** Further improvement, validation, and application of CymoSkew biotic index for the ecological status assessment of the Greek coastal and transitional waters. *Ecol. Indic.*, 118, 106727.
- Orlando-Bonaca, M., J. Francé, B. Mavrič, M. Grego, L. Lipej, V. Flander Putrle, M. Šiško & A. Falace (2015):** A new index (MediSkew) for the assessment of the *Cymodocea nodosa* (Ucria) Ascherson meadows's status. *Marine Environmental Research*, 110, 132-141.
- Orlando-Bonaca, M., L. Lipej & J. Francé (2016):** The most suitable time and depth to sample *Cymodocea nodosa* (Ucria) Ascherson meadows in the shallow coastal area. Experiences from the northern Adriatic Sea. *Acta Adriatica*, 57(2), 251-262.
- Orlando-Bonaca, M., J. Francé, B. Mavrič & L. Lipej (2019):** Impact of the Port of Koper on *Cymodocea nodosa* meadow. *Annales, Series Historia Naturalis*, 29(2), 187-194.
- Orth, R.J., T.J.B. Carruthers, W.C. Dennison, C.M. Duarte, J.W. Fourqurean, K.L. Heck, A.R. Hughes, G.A. Kendrick, W.J. Kenworthy, S. Olyarnik, F.T. Short, M. Waycott & S.L. Williams (2006):** A global crisis for seagrass ecosystems. *BioSci.*, 56, 987-996.
- Papathanasiou, V., S. Orfanidis & M.T. Brown (2016):** *Cymodocea nodosa* metrics as bioindicators of anthropogenic stress in N. Aegean, Greek coastal waters. *Ecol. Indic.*, 63, 61-70.
- Repolho, T., B. Duarte, G. Dionísio, J.R. Paula, A.R. Lopes, I.C. Rosa, T.F. Grilo, I. Caçador, R. Calado & R. Rosa (2017):** Seagrass ecophysiological performance under ocean warming and acidification. *Sci. Rep.*, 7, 41443. doi: 10.1038/srep41443.
- Richir, J., N. Luy, G. Lepoint, E. Rozet, A. Alvera Azcarate & S. Gobert (2013):** Experimental in situ exposure of the seagrass *Posidonia australis* (L.) Delile to 15 trace elements. *Aquat. Toxicol.*, 140-141, 157-173.
- Short, F.T., B. Polidoro, S.R. Livingstone, K.E. Carpenter, S. Bandeira, J.S. Bujang, H.P. Calumpang, T.J.B. Carruthers, R.G. Coles, W.C. Dennison, P.L.A. Erftemeijer, M.D. Fortes, A.S. Freeman, T.G. Jagtap, A.M. Kamal, G.A. Kendrick, W.J. Kenworthy, Y.A. La Nafie, I.M. Nasution, R.J. Orth, A. Prathep, J.C. Sanciangco, B. van Tussenbroek & S.G. Vergara (2011):** Extinction risk assessment of the world's seagrass species. *Biol. Conserv.*, 144, 1961-1971.
- Spalding, M., M. Taylor, C. Ravilious, F.T. Short & E. Green (2003):** The distribution and status of seagrasses. In: Green, E.P. et al. (Eds.). *World atlas of seagrasses*. pp. 5-26.
- Terrados, J. & J. Borum (2004):** Why are seagrasses important? - Goods and services provided by seagrass meadows. In: Borum, J., C.M. Duarte, D. Krause-Jensen, & T.M. Greve (Eds). *European seagrasses: an introduction to monitoring and management*. The M&MS project, pp. 8-10.
- Touchette B.W. & J.M. Burkholder (2000):** Overview of the physiological ecology of carbon metabolism in seagrasses. *Journal of Experimental Marine Biology and Ecology*, 250, 169-205.
- Tuya, F., J.A. Martín & A. Luque (2002):** Impact of a marina construction on seagrass bed at Lanzarote (Canary Islands). *J. Coast. Conserv.*, 8, 157-162.
- Tuya, F., H. Hernandez-Zerpa, F. Espino & R. Haroun (2013):** Drastic decadal decline of the seagrass *Cymodocea nodosa* at Gran Canaria (eastern Atlantic): Interactions with the green algae *Caulerpa prolifera*. *Aquat. Bot.*, 105, 1-6.
- Tuya, F., L. Ribeiro-Leite, N. Arto-Cuesta, J. Coca, R. Haroun & F. Espino (2014):** Decadal changes in the structure of *Cymodocea nodosa* seagrass meadows: Natural vs. human influences. *Estuar. Coast. Shelf S.*, 137, 41-49.
- Unsworth, R.K.F., L.M. Nordlund & L.C. Cullen-Unsworth (2018):** Seagrass meadows support global fisheries production. *Conservation Letters*, 12(1), e12566. <https://doi.org/10.1111/conl.12566>.
- Water Framework Directive (2000):** Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy.
- Waycott, M., C.M. Duarte, T.J. Carruthers, R.J. Orth, W.C. Dennison, S. Olyarnik, A. Calladine, J.W. Fourqurean, K.L. Heck JR., A.R. Hughes, G.A. Kendrick, W.J. Kenworthy, F.T. Short & S.L. Williams (2009):** Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *P. Natl. Acad. Sci. USA*, 106, 12377-12381.
- Widdows, J., N.D. Pope, M.D. Brinsley, H. Asmus & R.M. Asmus (2008):** Effects of seagrass beds (*Zostera noltii* and *Z. marina*) on near-bed hydrodynamics and sediment resuspension. *Mar. Ecol. Prog. Ser.*, 358, 125-136.
- Žagar, D., V. Ramšak, M. Jeromel, M. Perkovič, M. Ličer & V. Malačič (2014):** Modelling sediment resuspension caused by navigation, waves and currents (Gulf of Trieste, northern Adriatic). In: de Almeida, A.B. et al. (Eds.). *3rd IAHR Europe Congress: Water engineering and research*. Book of abstracts. Faculty of Engineering of the University of Porto (FEUP), Portugal, pp. 86-87.