




Detecting geodesic circles in hyperbolic surfaces with persistent homology

Blaž Jelenc¹ · Žiga Virk² 

Received: 13 August 2024 / Accepted: 29 December 2024
© The Author(s) 2025

Abstract

In this paper we provide conditions under which a geodesic circle on a hyperbolic surface admits arbitrarily small geodesically convex neighborhoods. This implies that persistent homology using selective Rips complexes detects the length and the position of such a loop via persistent homology in dimensions one, two, or three. In particular, if a surface has a unique systole, then the systole can always be detected with persistent homology. The existential results of the paper are complemented by the corresponding quantitative treatments which explain the choice of parameters of selective Rips complexes as well as conditions, under which the detection occurs via the standard Rips complexes. In particular, if a surface has a unique systole, then the parameters depend on the first spectral gap in the length spectrum.

Keywords Simple closed geodesic · Rips complex · Persistent homology · Hyperbolic surface · Systole

Mathematics Subject Classification 55N31 · 57R19 · 55U05 · 57N65 · 52A55

1 Introduction

Persistent homology is a well established tool in theoretical and applied topology. It encodes topological and geometric information when combined with the Rips complexes on sufficiently tame metric spaces. While the encoding of the homology of an underlying space is well understood as it happens at small scales, the precise nature of the geometric information carried by persistent homology is generally not well understood. The few cases in which certain interpretative results on persistent homology are known include the one-dimensional persistent homology of geodesic spaces (it encodes the shortest homology base by [18, 19]),

The authors would like to thank the anonymous reviewers for their helpful comments. The second named author was supported by the Slovenian Research Agency grants No. J1-4001 and P1-0292.

✉ Žiga Virk
ziga.virk@fri.uni-lj.si
Blaž Jelenc
blaz.jelenc@fmf.uni-lj.si

¹ University of Ljubljana, Faculty of Mathematics and Physics, Ljubljana, Slovenia

² University of Ljubljana, Faculty of Computer and Information Science, Institute IMFM], Ljubljana, Slovenia




Fig. 1 A scheme of the two-dimensional footprint detection as described by Theorem 6.1




Fig. 2 A scheme of the three-dimensional footprint detection as described by Theorem 6.3

parts of persistent homology of ellipses [2] and regular polygons [3], persistent homology of contracting subsets [22], and the complete homotopy type of the Rips filtration of a circle [1].

Especially [1] is of great interest: it demonstrates that the Rips complexes of a circle attain the homotopy types of odd-dimensional spheres and thus a one-dimensional geometric object (a loop) generates higher-dimensional algebraic objects in persistent homology. This idea has led to [21] in which a theory for the detection of parts of the persistent homology of a subset within the persistent homology of the ambient space (we refer to such parts as footprints) is presented. In particular, it turns out that under specific conditions, a portion of the persistent homology of a loop in a space can be observed in the persistent homology of the space itself. While this part consists of odd-dimensional homology elements when the underlying loop is a geodesic circle γ (i.e., a circle equipped with a geodesic metric, see Preliminaries below for more details), it turns out that an additional two-dimensional homology element may also be generated from the geometric position of γ when γ is not a member of a shortest homology base. This two-dimensional element as detected in [21] requires somehow generous conditions on the neighborhood of γ . To rectify this shortcoming, selective Rips complexes were introduced in [23] (see also [13, 16]) as a modification of Rips complexes that facilitates the mentioned two-dimensional footprint under fairly general conditions. The main result of [23] states that each geodesic circle, which is a locally shortest loop and admits an arbitrarily small geodesically convex neighborhood, can be detected either with one- or two-dimensional persistent homology using appropriate selective Rips complexes.

In this paper we recast the results of [21] and [23] into the setting of complete orientable hyperbolic surfaces: **we provide simple conditions under which geodesic circles induce a two-dimensional (Theorem 6.1) or a three-dimensional (Theorem 6.3) footprint in persistent homology.** It turns out that the somewhat technical conditions of the two mentioned paper can be deduced from the existence of sufficiently large geodesic charts. For a scheme of our results see Figs. 1 and 2. The technical results leading to such a connection include the existence of geodesically convex neighborhoods and an introduction of \widetilde{DC} isolated loops. The parameters of the results are also quantified (after Theorem 6.1), leading to specific bounds on parameters of required selective Rips complexes and settings in which the detection takes place with classical Rips complexes. In particular, when the systole of a surface is unique, we can deduce that the parameters of the selective Rips complexes depend on the first spectral gap in the length spectrum.

The paper consists of two distinct parts. The first part is a treatment of geometrical properties in the context of differential geometry. It contains Preliminaries in Sect. 2, Retractable neighborhoods in Sect. 3 and the existence of geodesically convex neighborhoods in Sect. 4. The second part contains the applications to persistent homology. Section 5 contains preliminaries and adaptations of results on persistent homology to our setting. In Sect. 6 the results of the previous sections are combined to draw the main conclusions of the paper.

2 Preliminaries on differential geometry

Our objects of interest will be orientable smooth hyperbolic (i.e., having potentially non-constant negative Gaussian curvature: $K < 0$) surfaces equipped with a Riemannian metric. We will assume that the surfaces are complete. In particular, the geodesics are defined for all times, and as a consequence, every pair of points x, y on a surface with $d(x, y) = \ell$ is connected by a **shortest geodesic** of length ℓ . We should point out that the term “geodesics” in this paper refers to locally shortest paths as is common in the context of differential geometry [6, 9], which is prevalent in this paper. In contrast, term “geodesics” or “geodesic segment” in some related works of more general metric context [5, 7, 12, 18, 19, 21, 23] refers to isometric embeddings of intervals. Such paths (and their traces) will be referred to as **shortest geodesic** in this paper. A subset A of a surface \mathcal{S} is **geodesically convex** if for each pair of points $x, y \in A$ each shortest geodesic between them lies in A .

Given $c > 0$ let S_c^1 denote a circle equipped with a geodesic metric (meaning the distance between any two points on S_c^1 is the length of a shortest segment between the points) of circumference c . A **geodesic circle** on a surface \mathcal{S} is an isometric embedding of S_c^1 into \mathcal{S} for some $c > 0$. We will frequently identify a geodesic circle with its trace. Loop $\alpha \subset \mathcal{S}$ is a **bottleneck loop** if there exists a neighborhood of α in which α is the shortest representative of its free homotopy class. The equator on a sphere is a geodesic circle which is not a bottleneck loop. It is not hard to construct a bottleneck loop, which is not a geodesic circle.

2.1 Variations of Arc Length

(see [9, p.238 and p.339] for some background)

It is well known that small perturbations of closed geodesics increase the length of the perturbed path in our setting (i.e., in smooth orientable hyperbolic surfaces), which implies that geodesic circles are automatically bottleneck loops. While the formal statement of this fact for our purposes could be deduced from Theorem 3.1, we recall the argument for the illustrative purposes.

Let $\gamma(s) : [0, L] \rightarrow \mathcal{S}$ be a naturally parametrized smooth loop. A variation is a smooth map

$$h : [0, L] \times (-\varepsilon, \varepsilon) \rightarrow \mathcal{S}$$

$$(s, t) \mapsto h(s, t)$$

such that $h(s, 0) = \gamma(s)$ and $h(0, t) = h(L, t)$ (and the coincidence also holds for all derivatives). Denote by

$$V(s) = \frac{\partial h}{\partial t}(s, 0)$$

the *variational vector field*. One can associate h to V by $h(s, t) = \exp_{\gamma(s)}(tV)$. Let

$$\mathcal{L}(t) = \int_0^L \left| \frac{\partial h}{\partial s}(s, t) \right| ds$$

be the length of the variation. Then

$$\mathcal{L}'(0) = - \int_0^L A(s) \cdot V(s) ds$$

where $A(s) = \frac{D}{ds} \frac{\partial h}{\partial s}$ (this is zero for a geodesic) and

$$\mathcal{L}''(0) = \int_0^L \left(\left| \frac{D}{ds} V(s) \right|^2 - K(s) |V(s)|^2 \right) ds.$$

Here the variational vector field $V(s)$ is orthogonal to $\gamma'(s)$. As each loop (and its length) in S can be approximated by a smooth loop, we obtain the following result.

Theorem 2.1 *Let $\gamma(s)$ be a simple closed geodesic on S . If the Gaussian curvature satisfies $K(s) < 0$ along the curve then γ is a bottleneck loop.*

Proof By the discussion above $\mathcal{L}'(0) = 0$ along γ and there exists $\kappa > 0$ such that $\mathcal{L}''(0) > \kappa$ on some tubular neighborhood of γ . Hence every nontrivial variation strictly increases length. □

Remark The converse does not hold in general, that is, if every variation increases length it does not necessarily mean that $K(s)$ is negative, see Fig. 3. Suppose that $K(s) = \cos s - \frac{1}{2}$ and let $V(s) = f(s)(\gamma'(s) \times \vec{n})$. Then the above integral for the second derivative is

$$\int_0^{2\pi} \left(f'(s)^2 - \left(\cos s - \frac{1}{2} \right) f(s)^2 \right) ds.$$

Write $f(s) = a_0 + \sum_{n=1}^{\infty} (a_n \cos(ns) + b_n \sin(ns))$ using the Fourier series and the integral equals

$$\begin{aligned} & 2\pi a_0 \left(\frac{1}{2} a_0 - a_1 \right) + \sum_{n=1}^{\infty} \pi \left(n^2 + \frac{1}{2} \right) (a_n^2 + b_n^2) \\ & = \pi(a_0 - a_1)^2 + \frac{\pi}{2} a_1^2 + \sum_{n=1}^{\infty} \pi \left(n^2 + \frac{1}{2} \right) b_n^2 + \sum_{n=2}^{\infty} \pi \left(n^2 + \frac{1}{2} \right) a_n^2. \end{aligned}$$

The minimal value (zero) is clearly attained when all $a_i, b_j = 0$, so for every nontrivial variation the integral is strictly larger than zero even though $K(s)$ is not strictly negative.

2.2 Geodesic coordinates

(see [17, p.242] for more details on the topic)

Let S be an orientable surface and let $\gamma(v)$ be a naturally parametrized simple closed geodesic on S . Let $\vec{A}(v)$ be a vector field along γ , perpendicular to γ and $|\vec{A}(v)| = 1$. Let $\sigma(u, v) = \Gamma_v(u)$ be the resulting parametrization of the surface, see Theorem 2.2.

Fig. 3 The indicated geodesics is still a bottleneck loop, despite the Gaussian curvature along it not being strictly negative





Fig. 4 Geodesic coordinates



Theorem 2.2 (See [17, Proposition 9.5.1] for a proof) *The parametrization $\sigma(u, v) = \Gamma_v(u)$ is a chart for \mathcal{S} in a neighbourhood of any point $(0, v)$ with the first fundamental form given by*

$$\begin{bmatrix} 1 & 0 \\ 0 & G(u, v) \end{bmatrix},$$

where G is a smooth function such that $G(0, v) = 1$ and $G_u(0, v) = 0$ (Fig. 4).

To summarize, the parametrization $\Gamma_v(u)$ is such that $u = \text{const.}$ and $v = \text{const.}$ form an orthogonal system of curves, with $\Gamma_v(0)$ is a geodesic curve and $\Gamma_{v_0}(u)$ is a geodesic curve and the first fundamental form is given by

$$\begin{bmatrix} 1 & 0 \\ 0 & G(u, v) \end{bmatrix} \text{ with } G(0, v) = 1 \quad G_u(0, v) = 0.$$

Theorem 2.2 describes geodesic coordinates along γ at any point of a geodesic γ . When γ is a closed geodesic of length ℓ on an orientable surface, we can combine these into a consistent **geodesic coordinates** along γ . In particular, there is a width $\varepsilon > 0$ and a smooth embedding

$$\begin{aligned} H &: (-\varepsilon, \varepsilon) \times [0, \ell] /_{0 \simeq \ell} \rightarrow \mathcal{S} \\ H &: (u, v) \mapsto H(u, v), \end{aligned}$$

such that $H(u, v) = \Gamma_v(u)$. Map H will represent such geodesic coordinates throughout the paper and will also be referred to as **geodesic chart**.

3 Retractable neighborhood

The main goal of this section is to formalize the following phenomenon: given a path α in a geodesic chart around a closed geodesic γ , sliding α “perpendicularly” towards γ results in ever shorter paths, see Fig. 5. This is done by Theorem 3.1:

- (1) First for paths α “parallel” (in terms of geodesic coordinates) to γ .
- (2) Then for general paths α .
- (3) Finally for sliding α towards a parallel to γ .

Theorem 3.1 *Let \mathcal{S} be an orientable surface, let $\gamma(v)$ be a naturally parameterized simple closed geodesic of length ℓ on \mathcal{S} , and assume the Gaussian curvature K of \mathcal{S} is negative on γ . Choose $\varepsilon > 0$ such that the Gaussian curvature is negative on the ε -geodesic chart of γ*

$$H : (-\varepsilon, \varepsilon) \times [0, \ell] /_{0 \simeq \ell} \rightarrow \mathcal{S}$$

$$H : (u, v) \mapsto H(u, v)$$

Then the following conclusions hold:

- (1) For any $a < b \in [0, \ell]$ the length of the curve $H(u_1, v)$, $v \in [a, b]$ is strictly smaller then the length of the curve $H(u_2, v)$, $v \in [a, b]$ for $|u_1| < |u_2|$.
- (2) Denote by $\kappa_t(u, v) = H((1 - t)u, v)$ the deformation retraction of the (image of the) geodesic chart onto γ . Let $H(u(\tau), v(\tau))$, $\tau \in [a, b]$ be a curve on \mathcal{S} . Then $\frac{d}{dt} \kappa_t(u(\tau), v(\tau)) < 0$, i.e. the length of the curve $\kappa_t(u(\tau), v(\tau))$ is decreasing as t increases from 0 to 1.
- (3) Choose $0 < \delta < \varepsilon$. Let $H(u(\tau), v(\tau))$, $\tau \in [a, b]$ be a curve on \mathcal{S} with $u(\tau) \geq \delta$. Define $v_t(u, v) = H((1 - t)u(\tau) + t\delta, v(\tau))$. Then $\frac{d}{dt} v_t(u(\tau), v(\tau)) < 0$, i.e. the length of the curve $v_t(u(\tau), v(\tau))$ is decreasing as t increases from 0 to 1.

Proof (1) In geodesic coordinates (see Theorem 2.2 for the properties that will be used throughout this argument) the Gaussian curvature is expressed as

$$K = -\frac{1}{2\sqrt{G}} \left(\frac{G_u}{\sqrt{G}} \right)_u = -\frac{1}{2\sqrt{G}} \left(\frac{G_{uu}\sqrt{G} - \frac{G_u^2}{2\sqrt{G}}}{|G|} \right),$$

by the Brioschi formula. This implies that $G_{uu}(0, v) > 0$. Since $G_u(0, v) = 0$ by the construction of geodesic coordinates, we get that $G_u(u, v) > 0$ for $u > 0$ and $G_u(u, v) < 0$ for $u < 0$. In particular $G(u, v) > 1$ for $|u| > 0$. The curve $H(u_0, v)$, $v \in [a, b]$ has length equal to

$$L(u_0) = \int_a^b \sqrt{G(u_0, v)} dv.$$

Since $L'(u_0) = \int_a^b \frac{G_u(u_0, v)}{2\sqrt{G(u_0, v)}} dv$, $L(u)$ is decreasing for negative u and increasing for positive u and this means that the length of the curve $H(u_1, v)$, $v \in [a, b]$ is strictly smaller then the length of the curve $H(u_2, v)$, $v \in [a, b]$ for $0 < u_1 < u_2$ (and similarly for $u_2 > u_1 > 0$).

(2) The length of the curve is

$$L(t) = \int_a^b \sqrt{(1 - t)^2 u'(\tau)^2 + G((1 - t)u(\tau), v(\tau)) v'(\tau)^2} d\tau.$$




Fig. 5 Sliding a path α towards γ results in ever shorter paths

The derivative

$$\frac{dL(t)}{dt} = \int_a^b \frac{-2(1-t)u'(\tau)^2 + G_u((1-t)u(\tau), v(\tau))v'(\tau)^2(-u(\tau))}{2\sqrt{(1-t)^2u'(\tau)^2 + G((1-t)u(\tau), v(\tau))}} d\tau$$

is always negative, since $-2(1-t)u'(\tau)^2$ is clearly negative and G_u is positive for positive u and negative for negative u . So as t flows from $0 \rightarrow 1$, $L(t)$ decreases. The proposed deformation retraction $\kappa_t(u, v) = H((1-t)u, v)$ indeed shortens all paths (Fig. 5).

(3) The proof is mostly the same as that of (2). □

Remark The theorem above deals with the case where K is negative. However, a version of the theorem for non-positive K can be proved in the same manner: we could replace *negative curvature* by *non-positive curvature* and change all strict inequalities (except for $\varepsilon > 0$, of course) to non-strict inequalities.

4 Geodesic convexity of neighborhoods

In this section we consider the existence of small geodesically convex neighborhoods of geodesic circles. We will first discuss why some of the assumptions are necessary for such a statement.

If there is no condition on the Gaussian curvature then the geodesic chart H (in particular, its image) is not geodesically convex in general. For example, think about the equator on a sphere, which has no small geodesically convex neighborhood. Even if $K \leq 0$ everywhere, a geodesic chart is not necessarily geodesically convex at all widths ε . We demonstrate this fact on the surface in Fig. 6, which has $K \leq 0$ everywhere. Choose a geodesic circle γ circumventing any of the visible holes on the said surface. Clearly there are geodesic charts of γ , that are not geodesically convex.

Furthermore, not every simple closed geodesic is geodesically convex as Fig. 7 shows, and the same goes for its small neighborhoods.

Now that the required conditions are established, we proceed by proving the existence of geodesically convex neighborhoods provided a wide enough geodesic chart exists.

Fig. 6 Scherk's singly periodic surface, also called the Scherk-tower





Fig. 7 Closed geodesic which is not geodesically convex and hence not a geodesic circle



Theorem 4.1 *Let \mathcal{S} be a complete orientable surface with $K < 0$ and let γ be a naturally parametrized simple closed geodesic of length ℓ . Suppose that*

$$H((-D, D) \times [0, \ell] /_{0 \simeq \ell}) \subset \mathcal{S}$$

is a geodesic chart around γ , where $D > \ell/4$. Then for each $\delta \leq \frac{D-\ell}{2}$, neighborhood $H((-\delta, \delta) \times [0, \ell])$ is geodesically convex. Furthermore, for each $\delta' < \frac{D-\ell}{2}$, neighborhood $H([-\delta', \delta'] \times [0, \ell])$ is geodesically convex.

Proof The situation is depicted in Fig. 8. Choose $p, q \in H((-\delta, \delta) \times [0, \ell])$ and let α be a shortest geodesic between them. Assume α does not lie entirely in $H((-\delta, \delta) \times [0, \ell])$. We analyse the situation by considering two separate cases:

- (1) Assume α lies in $H((-D, D) \times [0, \ell])$. Using case (3) of Theorem 3.1 we can slide the part of α lying outside $H((-\delta, \delta) \times [0, \ell])$ inside $H((-\delta, \delta) \times [0, \ell])$ while keeping endpoints p, q intact and decreasing the length of the path. The result is a path between p and q which is shorter than α , a contradiction.
- (2) Assume α does not lie in $H((-D, D) \times [0, \ell])$, see Fig. 9. On one hand this means α starts at p , has to reach the complement of $H((-D, D) \times [0, \ell])$ (which is at distance at least $\ell/4 + \delta$ from p) and then has to reach q again (meaning it has to traverse a distance at least $\ell/4 + \delta$ from p again), resulting in a lower bound $\ell/2 + 2\delta$ for the length of α . On the other hand we can construct a different path from p to q using γ . Start at p and follow a geodesic towards γ (of length less than δ) to reach $P_0 \in \gamma$. In a similar fashion initiate a new path segment by starting at q and follow a geodesic towards γ (of length less than δ) to reach $Q_0 \in \gamma$. Connect P_0 and Q_0 by a path along γ of length at most $\ell/2$. Concatenating these three paths we obtain a path from p to q of length less than $\ell/2 + 2\delta$. This contradicts the lower bound of the previous paragraph.

We conclude that α lies entirely in $H((-\delta, \delta) \times [0, \ell])$.

The statement for closed neighborhoods can be proved in the same way. □

Remark 4.2 Theorem 4.1 is stated for the case $K < 0$ as this is one of the assumptions of our eventual applications to persistent homology, and the proof of case (1) follows fairly easily. However, Theorem 4.1 as stated also holds when $K \leq 0$. In order to prove it we only need to modify case (1) of the corresponding proof as follows.

Assume $\alpha = H(u(t), v(t)): [0, 1] \rightarrow \mathcal{S}$ lies in $H((-D, D) \times [0, \ell])$. Without the loss of generality we may assume that $p = H(\delta, v_p) = H(u(0), v(0)), q = H(\delta, v_q), H(u(1), v(1))$ for some $\delta < D$, and $u(t) > \delta, \forall t \in (0, 1)$. The length of α equals

$$L_1 = \int_{t=0}^1 \sqrt{u'^2 + G(u(t), v(t)) \cdot v'^2} dt.$$

By the same reasoning the length of the projection of α onto $H([-\delta, \delta] \times [0, \ell])$ equals

$$L_2 = \int_{t=0}^1 \sqrt{G(\delta, v(t)) \cdot v'^2} dt.$$

By the assumptions:

- (1) u' is non-zero at some interval as $u(t) > \delta, \forall t \in (0, 1)$, and
- (2) $G(u(t), v(t)) \geq G(\delta, v(t))$ as $G(0, v) = 1$ and $G_{uu}(u, v) > 0$ for $u > 0$ (see part (1) of the proof of Theorem 3.1).

BLAŽ JELENC AND ŽIGA VIRK




Fig. 8 Setup of Theorem 4.1




Fig. 9 A sketch of a part of the proof of Theorem 4.1. If a shortest geodesic between p and q reaches the complement of $H((-D, D) \times [0, \ell])$ as depicted on the left, then there exists a shorter path between the two points (on right) along γ

As a consequence, $L_1 > L_2$.

The conditions of Theorem 4.1 contain the size of a geodesic chart. Throughout the rest of this section we explain how this condition may be dropped in the case of unique shortest closed geodesics on a surface.

We first recall the Cartan-Hadamard Theorem in Riemannian geometry.

Theorem 4.3 *Let \mathcal{S} be a complete surface with $K \leq 0$ and $p \in K$. Then the map*

$$\exp_p : T_p\mathcal{S} \rightarrow \mathcal{S} \quad p \in \mathcal{S}$$

is a universal covering projection.

Theorem 4.4 *Let \mathcal{S} be a complete orientable surface with $K \leq 0$ and let γ be a simple contractible loop. Then γ bounds a simple region (diffeomorphic to disc) in \mathcal{S} .*

Proof Let γ be a simple contractible loop. Then the lift to the universal covering space is also a simple loop $\tilde{\gamma}$. By the Jordan-Schönflies Theorem $\tilde{\gamma}$ bounds a simple region \mathcal{D} .

We now intend to prove that the covering projection restricted to \mathcal{D} is injective. For each non-trivial deck transformation g on the universal cover we have $\tilde{\gamma} \cap g(\tilde{\gamma}) = \emptyset$ as γ is simple. Furthermore, if $g(\mathcal{D}) \subseteq \mathcal{D}$, then g would have a fixed point (any point of the $\bigcap_{n \in \mathbb{N}} g^n(\mathcal{D}) \neq \emptyset$ is a fixed point by a standard argument), a contradiction. Hence $\mathcal{D} \cap g(\mathcal{D}) = \emptyset$ and thus the

covering projection is injective on \mathcal{D} . Consequently, the image of \mathcal{D} by the covering projection is a homeomorphic image of \mathcal{D} , whose boundary (in S) is γ . □

The (unmarked) **length spectrum** of a manifold is the collection of lengths of all closed geodesics. Each compact complete hyperbolic manifold has a discrete length spectrum. For a proof see, for example, [4, Lemma 3.1 and Remark 3.2]. Each shortest closed geodesic in a manifold is called **systole**. Each systole is a geodesic circle. The length of a systole is hence the first value of the length spectrum.

Proposition 4.5 combined with Remark 4.6 states that if the systole of a hyperbolic surface is unique, a choice of D as in Theorem 4.1 can be made.

Proposition 4.5 *Let S be a compact complete orientable surface with $K \leq 0$. Assume a closed loop γ of length ℓ is the unique systole of S and $L > \ell$ is the second value of the length spectrum, i.e., each closed loop of length less than L is either contractible or homotopic to γ .*

Then parameter D of Theorem 4.1 can be chosen to be $D = L/2 - \ell/4 > L/4$, i.e.,

$$H((-D, D) \times [0, \ell] /_{0 \simeq \ell}) \subset S$$

is a geodesic chart around γ .

Remark 4.6 Before we embark on the proof of Proposition 4.5 we briefly explain why a choice of L as assumed in Proposition 4.5 always exists if γ does. As S is compact then, as was mentioned before Proposition 4.5, the length spectrum of S is discrete and hence L can be chosen to be the second value of the length spectrum (and use the fact that each non-contractible loop has a representative as a closed geodesic). In particular, the pair ℓ, L represents the first spectral gap of the length spectrum.

Proof of Proposition 4.5 We need to prove that H is injective on $(-D, D) \times [0, \ell]$. Assume on the contrary, that there exists $D' < D$ such that there exist two geodesics which:

- start at points $p, q \in \gamma$;
- are perpendicular to γ at these points;
- have their first point of intersection at z with $d(z, \gamma) = D'$.

See Fig. 10 for a sketch of such a situation. The corresponding geodesic segments (from z to p and q , and a shortest segment from p to q along γ) form a geodesic triangle T with angles being $\pi/2, \pi/2$ and a non-trivial angle. Triangle T as a loop is of length at most $D + D + \ell/2 < L$ hence T is contractible. By Theorem 4.4 T bounds a disc. By the Gauss-Bonnet Theorem we get that $\int\int_D K dS > 0$, a contradiction. □

5 Preliminaries on persistent homology and footprint detection

Persistent homology is a type of parameterized version of homology. Ever since its introduction two decades ago the corresponding theory and applications witnessed intense development that expanded onto other fields of mathematics and science. For a general exposition on the topic see [10]. In this paper we will focus on a specific setting of hyperbolic surfaces. On the other hand, our treatment will be slightly more general than the standard approaches in persistent homology as we will not restrict our choice of coefficients to fields. We proceed by briefly presenting our setting. For a similar setting see [18, 19, 21, 23].




Fig. 10 A sketch of a part of the proof of Proposition 4.5: two geodesics perpendicular to γ having z as the first point of their intersection. Note that geodesics could also emerge from γ in different directions

Let X be a metric space and fix a scale $r > 0$. The **Rips complex** $\text{Rips}(X; r)$ is an abstract simplicial complex with the vertex set X defined by the following rule: a finite $\sigma \subset X$ is a simplex iff $\text{Diam}(\sigma) < r$.

Definition 5.1 [23] Let Y be a metric space, $r_1 \leq r_2, n \in \mathbb{N}$. **Selective Rips complex** $\text{sRips}(Y; r_1, n, r_2)$ is an abstract simplicial complex defined by the following rule: a finite subset $\sigma \subset Y$ is a simplex iff the following two conditions hold:

- (1) $\text{Diam}(\sigma) < r_1$;
- (2) there exist subsets $U_0, U_1, \dots, U_n \subset U$ of diameter less than r_2 such that $\sigma \subset U_0 \cup U_1 \cup \dots \cup U_n$.

The collection of Rips complexes of X for all positive $r > 0$ can be assembled together into the Rips filtration $\{\text{Rips}(X; r)\}_{r>0}$ of X consisting additionally of bonding maps $i_{r_1, r_2}: \text{Rips}(X; r_1) \rightarrow \text{Rips}(X; r_2)$, which are natural inclusions (identities on vertices) for all $r_1 < r_2$. Obtaining a filtration of selective Rips complexes we are required to make a more specific choice of positive increasing functions $r_1(t) \leq r_2(t)$.

Persistent homology is obtained by applying the homology functor to any filtration. When coefficients of a homology form a field the resulting persistent homology may under appropriate conditions (for example, if X is finite) be presented by a collection of intervals, which give rise to two well-known visualizations of persistent homology: persistence diagram and barcodes (see [10] for details).

5.1 Geodesic circles and persistent homology

We next present the known results which explain how geodesic circles in Riemannian manifolds generate algebraic objects (**footprints**) in persistent homology in various dimensions. Let γ be a geodesic circle of length ℓ in a Riemannian surface X and fix coefficients G for all homology groups in this section. Theorem 5.2 states that if γ is a member of a shortest homology base, then it induces a one-dimensional footprint which dies at $|\gamma|/3$. In this case it is the topological nature of γ that induces the footprint and hence the name **topological footprint**.

Theorem 5.2 [18] (Generating 1-dimensional (topological) footprint) *Let X be a compact semi-locally simply connected geodesic space and let G be an Abelian group. If γ is a member of a shortest homology base, then for each $r > 0$ a discretization of γ (called r -sample of γ in [18]) at scale r represents a homology class $Q_r \in H_1(\text{Rips}(X; r); G)$ such that:*

- For each $r_1 < r_2$ the inclusion $\text{Rips}(X; r_1) \rightarrow \text{Rips}(X; r_2)$ induced map maps Q_{r_1} to Q_{r_2} , and
- $Q_r \neq 0$ iff $r < |\gamma|/3$.

However, the focus of this paper is to also detect geodesic circles which are not contained in any shortest homology base and to detect γ via higher-dimensional homology.

In [21] a generic method for detecting γ through higher dimensional persistent homology is described. The result is inspired by [1], in which authors demonstrate that the homotopy type of a circle via Rips complexes attains all odd dimensional spheres. It turns out that this result may be used to prove that $\gamma \subset X$ sometimes induces odd-dimensional homology elements in persistent homology of X via Rips filtration. Additionally, two-dimensional elements may also appear when γ is not a member of a shortest homology base. It turns out that these two-dimensional elements appear under very weak assumptions if selective Rips complexes are used [23].

We continue by providing technical prerequisites, details and adjustments of these two results as we will later combine them with the results of the previous sections.

5.2 Detection via selective Rips complexes

We start with detection results via selective Rips complexes as described in [23]. Despite being chronologically more recent than the approach via deformation contractions described in the following subsection, we describe this setting first due to its simplicity. Broadly speaking, it turns out that each γ which is a geodesic circle, a bottleneck loop and has arbitrarily small convex neighborhood can be detected either with 1-dimensional persistent homology (in case γ is a member of a shortest homology base by [18]) or by a 2-dimensional persistent homology via selective Rips complexes. In this case it is the geometric property of a neighborhood of γ that induces the footprint in the absence of a topological footprint and hence the name **geometric footprint**.

Theorem 5.3 [23] (Generating 2-dimensional (geometric) footprint) *Let X be a geodesic locally compact, semi-locally simply connected space and let G be an Abelian group. Assume α is a geodesic circle in X satisfying the following properties:*

- (1) α is a bottleneck loop;
- (2) α is homologous to a non-trivial G -combination of loops $\beta_1, \beta_2, \dots, \beta_k$ of length at most $|\alpha|$, none of which is homotopic to α or α^- ;
- (3) α has arbitrarily small geodesically convex neighborhood.

Then there exist bounds $B_1 > |\alpha|/3$ and $B_2 > 0$ such that for all increasing bijections $a \geq b: (0, \infty) \rightarrow (0, \infty)$, and for all $r > 0$ such that $B_1 > a(r) > |\alpha|/3$ and $B_2 > b(r)$, there exists a non-trivial

$$Q_r \in H_2(\text{sRips}(X; a(r), 2, b(r)); G)$$

satisfying the following properties:

- (1) $\forall r_1 < r_2$ with $a(r_i) > |\alpha|/3$ and $b(r_i) < B, \forall i$, we have $i_{r_1, r_2}^G(Q_{r_1}) = Q_{r_2}$, where $i_{r_1, r_2}^G : H_2(\text{sRips}(X; a(r_1), 2, b(r_1)); G) \rightarrow H_2(\text{sRips}(X; a(r_2), 2, b(r_2)); G)$ is the natural inclusion induced map.
- (2) $\forall q : a(q) \leq |\alpha|/3$ there exists no Q_q with $i_{q, r}(Q_q) = Q_r$.

5.3 Detection via Rips complexes

The second way of detection was described in [21] and works in a fairly general setting. For our purposes we will adapt the results of [21] to our setting. We start by defining deformation contractions, i.e., maps inducing homotopy equivalence on Rips complexes (observe relation with retractable neighborhoods in Sect. 3, it will be employed later).

Definition 5.4 [14, 21] Let X be a metric space and $A \subset X$. A continuous map $F : X \times [0, 1] \rightarrow X$ is called a **deformation contraction** (we will abbreviate it as DC and write $X \xrightarrow{DC} A$) if:

- (1) $F(x, 0) = x, F(x, 1) \in A, F(a, t) = a, \forall x \in X, a \in A, t \in [0, 1]$, and
- (2) $d(F(x, t'), F(y, t')) \leq d(F(x, t), F(y, t)), \forall x, y \in X, t' > t$.

If additionally $d(F(x, t'), F(y, t')) < d(F(x, t), F(y, t))$ holds for all pairs $(x, y) \in (X \setminus A) \times X$ and for all $t' > t$, then F is called a strict deformation contraction (SDC or $X \xrightarrow{SDC} A$).

Proposition 5.5 [21] Suppose $X \xrightarrow{DC} A$ via a map F . Then the inclusions $\text{Rips}(A; r) \hookrightarrow \text{Rips}(X; r)$ are homotopy equivalences for each $r > 0$.

A local property used in [21] to deduce detection of a loop is that of DC-isolation (deformation contraction isolated). For our purposes a minor modification of this property \widetilde{DC} will be more useful.

Definition 5.6 Suppose $0 < D_1 \leq D_2$. A loop $\gamma \subset S$ in a complete surface with $K \leq 0$ is $\widetilde{DC}(D_1, D_2)$ -isolated if the following conditions hold for closed geodesic neighborhoods $N_1 = \overline{N}(\gamma, D_1/2)$ and $N_2 = \overline{N}(N_1, D_2)$ of γ :

- (1) N_2 is geodesically convex.
- (2) Sliding along geodesics perpendicular to γ towards γ in a geodesic chart N_2 (see Fig. 5 and (2) of Theorem 3.1) induces $N_2 \setminus \text{Int}(N_1) \xrightarrow{SDC} \partial N_1$ and $N_1 \xrightarrow{SDC} \gamma$, see Fig. 11.

Note that if γ is $\widetilde{DC}(D_1, D_2)$ -isolated then $\overline{N}(\gamma, r)$ is geodesically convex for each $r \leq D_2$ by Theorem 3.1.

Proposition 5.7 Let $0 < D_1 \leq D_2$ and suppose a loop $\gamma \subset S$ in a complete surface with $K \leq 0$ is $\widetilde{DC}(D_1, D_2)$ -isolated. In this case the boundary ∂N_1 consists of two simple closed loops denoted by γ', γ'' , see Fig. 11. For each $p \in \gamma$ let p' and p'' denote the points on γ' and γ'' respectively, which are closest to p , see Fig. 12. Then the following two statements hold:

- (1) There exists $D' > |\gamma|/3$ such that for each $p, q \in \gamma$ with $d(p, q) \geq |\gamma|/3$ we have $d(p', q') > |\gamma|/3 + D'$ and $d(p'', q'') > |\gamma|/3 + D'$.
- (2) For each $r \in (|\gamma|/3, D')$ the inclusion induced maps $\text{Rips}(\gamma'; r) \hookrightarrow \text{Rips}(N_1; r)$ and $\text{Rips}(\gamma''; r) \hookrightarrow \text{Rips}(N_1; r)$ are homotopically trivial.




Fig. 11 A \widetilde{DC} -isolated loop γ




Fig. 12 A sketch the setting of Proposition 5.7

Proof (1) Map $(p, q) \mapsto d(p', q')$ is continuous and (by Definition 5.6) does not attain value $|\gamma|/3$ on a compact domain $\{(p, q) \in \gamma; d(p, q) \geq |\gamma|/3\}$ hence it has a lower bound above $|\gamma|/3$. The same holds for γ'' and the minimum of these two lower bounds is D' .

(2) By Proposition 5.5 we have $\text{Rips}(N_1; r) \simeq \text{Rips}(\gamma; r)$ with the homotopy equivalence arising from our setting (see Definition 5.6) mapping $p' \mapsto p$. Thus $\text{Rips}(\gamma'; r) \hookrightarrow \text{Rips}(N_1; r)$ is homotopic to a map $\text{Rips}(\gamma'; r) \rightarrow \text{Rips}(\gamma; r)$ mapping $p' \mapsto p$. By (1) the image of this map is actually contained in $\text{Rips}(\gamma; |\gamma|/3)$, which is homotopy equivalent to the circle by [1]. On the other hand, $\text{Rips}(\gamma; r)$ is homotopy equivalent to a sphere of dimension at least three by [1], hence the map in question is nullhomotopic. \square

The property of being DC -isolated was used in [21] to deduce an appearance of odd-dimensional homology elements. In a similar fashion we can prove a similar result for \widetilde{DC} isolated loops in our setting. In this case the combinatorics of Rips complexes from [1] induces the footprint and hence the name **combinatorial footprint**.

Theorem 5.8 (Generating 3-dimensional (combinatorial) footprint via Rips complexes) *Suppose S is a complete orientable surface with $K \leq 0$, γ is a $\widetilde{DC}(D, D)$ isolated geodesic circle for some $D \in (|\gamma|/3, 2|\gamma|/5)$, and G is a group. Then there exists $D' \in (|\gamma|/3, 2|\gamma|/5)$*

such that the inclusion $\gamma \hookrightarrow X$ induces an inclusion

$$\{G\}_{r \in (|\gamma|/3, D')} \cong \{H_3(\text{Rips}(\gamma; r); G)\}_{r \in (|\gamma|/3, D')} \hookrightarrow \{H_3(\text{Rips}(X; r); G)\}_{r \in (|\gamma|/3, D')}.$$

In particular, for each $r \in (|\gamma|/3, D')$ there exists a non-trivial

$$Q_r \in H_3(\text{Rips}(X; r); G)$$

such that:

- (1) $\forall r_1 < r_2$ from $(|\gamma|/3, D')$ we have $i_{r_1, r_2}^G(Q_{r_1}) = Q_{r_2}$, where

$$i_{r_1, r_2}^G : H_3(\text{Rips}(X; r_1); G) \rightarrow H_3(\text{Rips}(X; r_2); G)$$

is the natural inclusion induced map.

- (2) $\forall q : a(q) \leq |\gamma|/3$ there exists no Q_q with $i_{q, r}(Q_q) = Q_r$.

Proof Note that $\{G\}_{r \in (|\gamma|/3, D)} \cong \{H_3(\text{Rips}(\gamma; r); G)\}_{r \in (|\gamma|/3, D)}$ follows from [1] as $\text{Rips}(\gamma; r) \simeq S^3, \forall r \in (|\gamma|/3, D)$. Choose D' as in Proposition 5.7.

We set a Mayer-Vietoris long exact sequence using the notation of Definition 5.6. For a fixed $r \in (|\gamma|/3, D)$ define

$$A = \text{Rips}(N_2; r) \simeq \text{Rips}(\gamma; r),$$

$$B = \text{Rips}(\mathcal{S} \setminus \text{Int}(N_1); r).$$

Note that

$$A \cap B = \text{Rips}(N_2 \setminus \text{Int}(N_1); r) \simeq \text{Rips}(\gamma'; r) \sqcup \text{Rips}(\gamma''; r)$$

as γ' and γ'' are at distance more than r . Furthermore, observe that each point of $\mathcal{S} \setminus N_2$ is at distance at least $D > r$ from each point of N_1 due to $N_2 = \overline{N}(N_1, D)$. As a result, each subset of \mathcal{S} of diameter less than r is contained either in $N_2 = \overline{N}(N_1, D)$ or $\mathcal{S} \setminus \text{Int}(N_1)$, implying

$$A \cup B = \text{Rips}(\mathcal{S}; r).$$

From the Mayer-Vietoris sequence we extract the following exact subsequence:

$$H_3(A \cap B; G) \rightarrow H_3(A; G) \oplus H_3(B; G) \rightarrow H_3(\text{Rips}(X, r); G).$$

By (2) of Proposition 5.6 the first map is trivial, which implies that the second map is an inclusion on $H_3(A; G) = H_3(\text{Rips}(\gamma; r); G)$. The formal conclusion follows from the functoriality of the Mayer-Vietoris sequence. \square

6 Final results

In this section we combine the geometric results of initial sections with the footprint detection results of Sect. 5 to describe detection (footprints) of geodesic circles on hyperbolic surfaces.




Fig. 13 A scheme of a geometric footprint detection




Fig. 14 A scheme of combinatorial footprint detection

6.1 Results with selective Rips complexes [two-dimensional footprint]

A scheme of the following result is provided in Fig. 13.

Theorem 6.1 *Let S be a compact complete orientable surface with $K < 0$ and let $\gamma \subset S$ be a geodesic circle. Assume either of the following holds:*

- (1) $H((-D, D) \times [0, \ell]) \subset S$ is a geodesic chart around γ , where $D > |\gamma|/4$.
- (2) Loop γ is the unique systole of S .

Then at $r = |\gamma|/3$ loop γ induces

- (i): A one-dimensional footprint in the sense of Theorem 5.2, or
- (ii): A two-dimensional footprint in the following sense: there exists a filtration of selective Rips complexes through which γ induces a two-dimensional footprint in the sense of Theorem 5.3.

Remark 6.2 Loops γ may induce both a one- and a two-dimensional footprint in the sense of Theorem 6.1 if it appears in a shortest homology base of $H_1(S; G)$ and is homologous to a non-trivial G -combination of loops $\beta_1, \beta_2, \dots, \beta_k$ of length at most $|\gamma|$, none of which is homotopic to γ or γ^- . In this case γ is not a member of each shortest homology base of $H_1(S; G)$.

Proof If γ is a member of a shortest homology base, conclusion (i) follows by Theorem 5.2. If that is not the case, (ii) and the conclusion of Theorem 5.3 follow if γ is a bottleneck loop (which holds always by Theorem 2.1) and has arbitrarily small geodesically convex neighborhood. The existence of the later follows either from Theorem 4.1 for assumption (1), or from Proposition 4.5 for assumption (2). □

Quantification of Theorem 6.1: Concerning conclusion (ii) we discuss the parameters of selective Rips complexes and conditions under which the induced two-dimensional footprint can be detected with the usual Rips complexes.

Starting with assumption (1) we have a geodesic chart of width D . Theorem 4.1 implies the existence of a geodesically convex geodesic chart of width $\tilde{T} = D/2 - |\gamma|/8$. The quantitative analysis of Theorem 5.3 in [23] implies that a selective Rips complex satisfies conclusion (ii) if the following holds: for $r > 0$ at which $a(r) = |\gamma|/3$ we need to have

$b(r) < \tilde{T} = D/2 - |\gamma|/8$ (recall notation of Theorem 5.3). In particular, the detection occurs with Rips complexes when $\tilde{T} > |\gamma|/3$, i.e., $D > 11|\gamma|/12$.

Continuing with assumption (2) we see that $D = L/2 - |\gamma|/4$, where $|g|$ and L are the first two values of the length spectrum of S and thus $L - |\gamma|$ is the first spectral gap. Consequently, a selective Rips complex satisfies conclusion (ii) if the following holds: for $r > 0$ at which $a(r) = |\gamma|/3$ we need to have $b(r) < (L - |\gamma|)/4$, i.e., the upper bound is the quarter of the first spectral gap. In particular, the detection occurs with Rips complexes when $(L - |\gamma|)/4 > |\gamma|/3$, i.e., $L > 7|\gamma|/3$. The ability of Rips complexes to detect γ with a two-dimensional footprint hence depends on the first spectral gap.

6.2 Results with Rips complexes [three-dimensional footprint]

A scheme of the following result is provided in Fig. 14.

Theorem 6.3 *Let S be a compact complete orientable surface with $K < 0$ and let $\gamma \subset S$ be a geodesic circle. Assume any of the following holds:*

- (1) $H([- \tilde{T}, \tilde{T}] \times [0, \ell]) \subset S$ is a geodesically convex geodesic chart around γ , where $\tilde{T} > 3|\gamma|/2$.
- (2) $H((-D, D) \times [0, \ell]) \subset S$ is a geodesic chart around γ , where $D > 13|\gamma|/4$.
- (3) Loop γ is the unique systole of S with $L > 7|\gamma|$ being the second smallest value of the length spectrum.

Then at $r = |\gamma|/3$ loop γ induces a three-dimensional footprint in the sense of Theorem 5.8.

Proof Assume (1) holds. Then by Theorem 3.1 there exists $\tilde{D} \in (|\gamma|/3, 2|\gamma|/5)$ such that γ is $\widetilde{DC}(\tilde{D}, \tilde{D})$ isolated, thus the conditions for Theorem 5.8 are satisfied.

Assumption (2) and Theorem 4.1 imply assumption (1). Assumption (3) and Proposition 4.5 imply assumption (1). \square

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Adamaszek, M., Adams, H.: The Vietoris-Rips complexes of a circle. *Pac. J. Math.* **290**, 1–40 (2017)
2. Adamaszek, M., Adams, H., Reddy, S.: On Vietoris-Rips complexes of ellipses. *J. Topol. Anal.* **11**, 661–690 (2019)
3. Adams, H., Chowdhury, S., Quinn Jaffe, A., Sibanda, B.: Vietoris-Rips Complexes of Regular Polygons, [arXiv:1807.10971](https://arxiv.org/abs/1807.10971)
4. Basmajian, A., Kim, Y.: Geometrically infinite surfaces with discrete length spectra. *Geom Dedicata* **137**, 219 (2008)
5. Bridson, M.R., Haefliger, A.: Metric spaces of non-positive curvature. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 319. Springer-Verlag, Berlin (1999)

6. Burago, D., Burago, Y., Ivanov, S.: *A Course in Metric Geometry*, Graduate Studies in Mathematics 33, AMS (2001)
7. Buyalo, S., Schroeder, V.: *Elements of asymptotic geometry*, EMS Monographs in Mathematics (2007)
8. Chazal, F., de Silva, V., Oudot, S.: Persistence stability for geometric complexes. *Geom. Dedicata* **173**, 193 (2014)
9. do Carmo, M.P.: *Differential Geometry of Curves and Surfaces*, Revised and Updated Second Edition. Dover Books on Mathematics. Dover Publications (2016)
10. Edelsbrunner, H., Harer, J.L.: *Computational Topology. An Introduction*. Amer. Math. Soc, Providence, Rhode Island (2010)
11. Gasparovic, E., Gommel, M., Purvine, E., Sazdanovic, R., Wang, B., Wang, Y., Ziegelmeier, L.: A Complete Characterization of the 1-Dimensional Intrinsic Čech Persistence Diagrams for Metric Graphs. In: Chambers E., Fasy B., Ziegelmeier L. (eds). *Research in Computational Topology*. Association for Women in Mathematics Series, vol 13. Springer, Cham, (2018)
12. Geoghegan, R.: *Topological methods in group theory*, Graduate Texts in Mathematics, 243. Springer, New York (2008)
13. Goričan, P.; Virk, Ž: Critical edges in Rips complexes and persistence. *Mediterr. J. Math.* **20**, 326 (2023)
14. Hausmann, J.-C.: On the Vietoris-Rips complexes and a cohomology theory for metric spaces. *Annal. Math. Studies* **138**, 175–188 (1995)
15. Katz, M.G.: *Systolic geometry and topology*, Mathematical Surveys and Monographs, vol. 137. American Mathematical Society, Providence, R.I. (2007)
16. Lemež, B., Virk, Ž: Reconstruction properties of selective Rips complexes. *Glasnik Matematički* **57**(2), 73–88 (2022)
17. Pressley, A.: *Elementary Differential Geometry*, 2nd edn. Springer, Berlin (2010)
18. Virk, Ž: 1-Dimensional Intrinsic Persistence of geodesic spaces. *J. Topol. Anal.* **12**, 169–207 (2020)
19. Virk, Ž: Approximations of 1-dimensional intrinsic persistence of geodesic spaces and their stability. *Revista Matemática Complutense* **32**, 195–213 (2019)
20. Virk, Ž.: Rips complexes as nerves and a Functorial Dowker-Nerve Diagram. *Mediterr. J. Math.* **18**, (2021)
21. Virk, Ž.: Footprints of geodesics in persistent homology, *Mediterr. J. Math.* **19**, (2022)
22. Virk, Ž: Contractions in persistence and metric graphs. *Bull. Malays. Math. Sci. Soc.* **45**, 2003–2016 (2022)
23. Virk, Ž: Persistent Homology with selective Rips complexes detects geodesic circles. *Mediterr. J. Math.* **21**, 170 (2024)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.