

A SYSTEMATIC REVIEW OF VITAMIN D STATUS AND DIETARY INTAKE IN VARIOUS SLOVENIAN POPULATIONS

SISTEMATIČNI PREGLED PREHRANSKEGA VNOSA IN PRESKRBLJENOSTI Z VITAMINOM D V SLOVENIJI

Maša HRIBAR^{1,2}, Evgen BENEDIK^{2,3}, Matej GREGORIČ⁴, Urška BLAZNIK⁴, Andreja KUKEČ⁵, Hristo HRISTOV¹, Katja ŽMITEK^{1,6}, Igor PRAVST^{1,2,6*}

¹Nutrition Institute, Tržaška cesta 40, 1000 Ljubljana, Slovenia

²University of Ljubljana, Biotechnical Faculty, Jamnikarjeva 101, 1000 Ljubljana, Slovenia

³University Medical Centre Ljubljana, Zaloška cesta 2, 1000 Ljubljana, Slovenia

⁴National Institute of Public Health, Trubarjeva 2, 1000 Ljubljana, Slovenia

⁵University of Ljubljana, Faculty of Medicine, Vrazov trg 2, 1000 Ljubljana, Slovenia

⁶VIST-Faculty of Applied Sciences, Gerbičeva cesta 53, 1000 Ljubljana, Slovenia

Received: Aug 26, 2021

Accepted: Nov 5, 2021

Review

ABSTRACT

Keywords:

vitamin D, dietary intake, serum 25-hydroxy-vitamin D, vitamin D deficiency, systematic reviews, Slovenia

Aim: Vitamin D (VitD) is involved in calcium and phosphate homeostasis, bone health, and normal functioning of the immune system. VitD status is monitored using serum 25-hydroxy-vitamin D (25(OH)D) as a biomarker. Serum 25(OH)D concentrations below 30 nmol/L indicate VitD deficiency and below 50 nmol/L indicate insufficiency. VitD can be synthesised endogenously in human skin when exposed to ultraviolet B (UVB) radiation. In the absence of sufficient UVB-light exposure, VitD intake becomes the main source of VitD, with a recommended daily intake of 20 µg. The aim of this study was to conduct a review and meta-analysis on the abovementioned topics, focusing on scientific studies in various Slovenian populations.

Methods: We conducted a systematic review and meta-analysis of published scientific papers, academic theses, or conference contributions reporting serum 25(OH)D status and VitD intake across various Slovenian populations. A search was carried out using Web of Science, Scopus, Medline, and the Slovenian library database.

Results: We identified 43 pertinent studies that addressed 25(OH)D status and 16 that addressed VitD intake. Serum 25(OH)D status was generally low across all populations, and notable seasonal variability was observed. VitD intakes were below 5 µg in all studies.

Conclusions: A general observation is that various population groups across Slovenia are at high risk of vitamin D insufficiency and deficiency, particularly during wintertime. Regarding vitamin D intake, all included studies reported daily intakes below the recommended level. We also identified key research gaps that need to be addressed to support further public health decision-making.

IZVLEČEK

Ključne besede:

vitamin D, prehranski vnos, serumski 25 hidroksi-vitamin D, preskrbljenost z vitaminom D, sistematični pregledi literature, Slovenija

Namen: Vitamin D (VitD) je vključen v metabolizem kalcija in fosforja, zdravje kosti in normalno delovanje imunskega sistema, hkrati pa je nezadostna preskrbljenost povezana z različnimi skeletnimi in neskeletnimi obolenji. Splošno uveljavljen biološki kazalnik preskrbljenosti z VitD je serumski koncentracija 25 hidroksi-vitamina D (25(OH)D). Koncentracije 25(OH)D pod 30 nmol/L so definirane kot hudo pomanjkanje, pod 50 nmol/L kot pomanjkanje, optimalna pa je koncentracija nad 75 nmol/L. VitD se endogeno sintetizira v koži ob izpostavitvi ultravijoličnim B (UVB) žarkom. Kadar je izpostavljenost UVB žarkom prenizka, postane vnos s hrano glavni vir VitD. V odsotnosti endogene sinteze je priporočeni vnos VitD 20 µg dnevno. Namen raziskave je bil sistematičen pregled znanstvene literature na področju preskrbljenosti s 25(OH)D in prehranskega vnosa VitD pri prebivalcih Slovenije.

Metode: Izvedli smo sistematičen pregled literature in metaanalizo znanstvenih člankov, zaključnih del in konferenčnih prispevkov, ki so poročali o preskrbljenosti s 25(OH)D in vnosu VitD pri različnih populacijskih skupinah prebivalcev Slovenije. Za iskanje literature smo uporabili podatkovne baze: Web of Science, Medline in Kooperativni online bibliografski sistemi in servisi (COBISS) brez omejevanja leta objave.

Rezultati: O preskrbljenosti s 25(OH)D je poročalo 43 znanstvenih prispevkov, medtem ko je o vnosu VitD poročalo 16 znanstvenih prispevkov. Preskrbljenost s 25(OH)D je bila relativno nizka v vseh zajetih populacijah, opaženo je bilo tudi znatno sezonsko nihanje. Vnos VitD je bil v vseh populacijah pod 5 µg.

Zaključki: Preskrbljenost s 25(OH)D je nizka, zato so mnogi prebivalci izpostavljeni povečanemu tveganju za (hudo) pomanjkanje, posebno v zimskem obdobju. Prav tako opažamo, da je prehranski vnos VitD precej nižji od priporočenega. Identificirane so bile tudi pomembne vrzeli, ki jih je treba nasloviti za sprejemanje na dokazih temelječih javnozdravstvenih ukrepov, ki bodo učinkoviteje podpirali zdravje prebivalcev Slovenije.

*Corresponding author: Tel. +386 590 68871; E-mail: igor.pravst@nutris.org

1 INTRODUCTION

Vitamin D (VitD) participates in numerous functions in the human body, including calcium and phosphate homeostasis, bone health, and the normal functioning of the immune system (1, 2). Deficient 25-hydroxy-vitamin D (25(OH)D) status causes rickets and osteomalacia (depending on age) (3, 4), contributes to the development of osteoporosis and myopathy, and has been linked to various non-skeletal disorders, such as acute and chronic respiratory infections, and risks of death due to cardiovascular disease, cancer, and other causes (5, 6). While numerous reports indicate a high prevalence of 25(OH)D deficiency, we should note that different thresholds are used in different studies. VitD status is determined considering the level of serum 25(OH)D, usually reported in both nanomoles per litre (nmol/L) and nanograms per millilitre (ng/mL; 1 ng/mL equals 2.5 nmol/L) (7). Thresholds for risk of 25(OH) deficiency and insufficiency are typically defined with serum concentrations below 30 and 50 nmol/L, respectively (3, 8, 9), while the Endocrine Society's guidelines highlight that optimal levels are above 75 nmol/L (3).

VitD can be synthesised endogenously in human skin when exposed to ultraviolet B (UVB) light. Considering the geographical location of Slovenia (latitudes 45-46° north), the intensity of UVB light is sufficient for VitD endogenous synthesis only from April to October, but not during wintertime (10). Therefore, seasonal variation in 25(OH)D status is expected in the population.

In the absence of UVB-induced endogenous synthesis, sufficient dietary intake of VitD is needed. Recommendations for VitD intake are not yet standardised. While the World Health Organization (WHO) recommends 10 µg VitD per day for adults below 65 years and 15 µg for the elderly population (11), the European Food Safety Authority (EFSA) and the nutrition societies of Germany, Austria, and Switzerland (D-A-CH) recommend a VitD intake of 15 and 20 µg per day, respectively for all above the age of 1 year (12). It should be noted that these are recommendations for dietary intake in the absence of endogenous UVB-induced VitD synthesis. However, the usual daily dietary VitD intake in adults is below 10 µg in most European countries (13). This is because only a few foods (i.e., fatty fish) are naturally rich in VitD (14), and those are not common in the European diet. Some countries, therefore, have introduced the fortification of particular foods with VitD (15) or prescribed VitD supplementation to certain subpopulation groups.

While several studies have investigated dietary intake or 25(OH)D status in various Slovenian populations, this topic has not yet been systematically reviewed. The aim of this study was thus to conduct a review and meta-analysis on the abovementioned topics, focusing on studies published either as scientific papers, academic thesis, or conference contributions.

2 METHODS

2.1 Search strategy and inclusion/exclusion criteria

We focused on two topics: A) serum 25(OH)D levels as a biomarker for VitD status, and B) VitD intake in various population groups in Slovenia. The main literature search was carried out in September 2020. The following four databases were searched (Box 1):

Box 1. Search strategy.

- Medline, Scopus, and Web of Science (WoS) - with the searching syntax: (TITLE-ABS-KEY (((“vitamin D”) OR (cholecalciferol)) AND sloven*)), and
- COBISS - Slovenian library database (Co-operative Online Bibliographic System & Services), with the searching syntax: kw=(“vitamin d” or cholecalciferol).

For this review, we included original scientific articles, academic thesis (diploma, masters, doctoral, and others), and conference contributions in the Slovene or English language. We included publications which provided data on either serum 25(OH)D levels or dietary VitD intake for the defined population. We included publications with original results, which were available as full texts or comprehensive abstracts, either in digital or hard copy. The literature search was conducted without any time limits, and therefore the identified results correspond to periods covered in the abovementioned databases.

In addition to the described literature search, we also asked members of the national expert working group on guidelines for sufficient vitamin D levels in the Slovenian population to provide information on the references with novel data on dietary VitD intake and 25(OH)D status in Slovenia that they are aware of. These members are experts and researchers in the field, and therefore well informed about nationally available data. The identified relevant publications, which were not already found in the literature search, were counted as additional records identified through other sources.

2.2 Data extraction and study selection process

The methods used in the present review follow the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement (16). After the literature search, we first removed all duplicates; secondly, we removed publications of which title/abstract did not fit the purpose of this review or were not original research. Finally, we removed publications where we could not obtain a full text. We combined publications describing the same data set with the same outcomes, referred to as a ‘research cluster’ (in Supplementary table 1, each presented in a separate row) from here onward. All publications were checked carefully, and the following data were extracted: description of the study period, study group(s), study results (distribution and

mean serum 25(OH)D levels, daily dietary VitD intake). To simplify comparisons, the results of identified studies were re-calculated to internationally accepted units (IU; conversion: 1 µg of vitamin D2 or D3=40 IU). Dietary VitD intake is presented in µg per day, while serum 25(OH)D levels are presented in nmol/L. The distribution of serum 25(OH)D levels is presented using the thresholds for VitD deficiency used in the original publication. However, with consideration of the most relevant thresholds (3, 8), we presented the prevalence of serum 25(OH)D levels below 30/50 nmol/L (VitD deficiency/insufficiency) and those above 50/75 nmol/L (sufficient/optimal level) separately.

2.3 Descriptive quantitative synthesis and meta-analysis

Statistical analyses were carried out using Stata Statistical Software, Release 15 (StataCorp LLC, College Station, TX, USA). The proportions of deficiency/optimal vitamin D status were analysed by applying the STATA Metaprop module for pooled meta-analyses of proportions. Analyses were carried out considering studies with healthy individuals, where seasonal proportions of deficiency/optimal 25(OH)D status were provided for a sample of at least 50 subjects.

3 RESULTS

The literature search resulted in 562 records. After the removal of duplicates (n=69) and the addition of records identified through other sources (n=9), a total of 502 records were screened for relevance. After the exclusion of 186 non-relevant records, we started collecting the full texts of 268 publications. Although we made a large effort to obtain full-text access to all of these records (we contacted the authors of publications, searched for records on various internet sources, and ordered the remaining missing publications through the Central Technical Library at the University of Ljubljana), we were unable to obtain full texts for 24 records. Therefore, the literature review was conducted using 58 relevant publications (Figure 1). Finally, we separated identified publications into two sections, depending on the reported study outcomes: a) serum 25(OH)D levels (25(OH)D status) and b) dietary intake of VitD. One publication (17) was used in both sections. In the section of 25(OH)D status, we included 43 texts and allocated them into 30 research clusters, and in the section of dietary VitD intake, we included 16 texts. The first reported research is dated as from 1993 (18) and the last from 2021 (19).

3.1 Serum 25(OH)D status

The literature review of 25(OH)D included 43 publications (Supplementary table 1), comprising 20 scientific journal articles, 11 conference papers, and 12 academic theses




Figure 1. PRISMA flow diagram (16) of the literature review process for vitamin D intake and status in various Slovenian populations. Notes: WOS: Web of Science; COBISS: Co-operative Online Bibliographic System & Services, 25(OH)D: 25-hydroxy-vitamin D; VitD: Vitamin D.

(details provided in Table 1). Altogether, we identified 30 research clusters, the majority referring to the adult population (n=21), one referred to older adults and adults, with an additional two for older adults and two for pregnant women. Two-thirds of the research clusters (n=21) reported data from patients (in- or out-patients), thirteen from the general population, and four reported separately for healthy controls and patient populations. Therefore, we considered these clusters twice, once in the healthy and once in the patient population. Four research clusters referred to children and one additional to newborn infants, giving five in total. In the meta-analyses, we included four studies that reported seasonal proportions of deficiency/optimal 25(OH)D status in a sample of at least 50 subjects.

3.1.1 Serum 25(OH)D concentrations in adults

Among the healthy adult population, three research clusters (20-22) were composed of healthy individuals, one cluster of presumably healthy individuals (23) (as the data collection was carried out retrospectively in a laboratory with self-pay service), and another three research clusters (24-27) of healthy individuals from control groups. A nationally representative study sample (adults 18-64 years)

was only used in one study (20), which was conducted in 2017-2018, and covered all calendar seasons. Serum 25(OH)D in adults ($n=125$; 18-64 years) was significantly affected by calendar season, sex, and physical activity level. Mean 25(OH)D level was 50.7 nmol/L (95% CI: 45.4-56.0 nmol/L), with 70.4 nmol/L (95% CI: 62.2-78.5) during the extended summer period (May-October), and 36.7 nmol/L (95% CI: 32.5-40.9 nmol/L) during the extended winter period (November-April). The distribution of mean serum 25(OH)D concentrations during the calendar year is presented in Figure 2. The next research cluster (22) included 240 individuals from periodic medical examinations. Blood samples were collected throughout the year, and seasonal variations in serum 25(OH)D levels are evident. Data are reported for every second month, and it is at its lowest in February (61.3 nmol/L) and highest in June (92.8 nmol/L). The third research cluster (21) included a healthy study population mostly from the central Slovenian region ($n=238$; 18-65 years). Blood samples were collected during winter months (January - March) and mean serum 25(OH)D level was 44.0 ± 17.0 nmol/L.




Figure 2. Box plots of the weighted mean serum 25(OH)D levels for different bi-monthly periods ($n=280$; monitoring time frame: 12 months in 2017-2018) with a presentation of outliers (●). Results of a nationally representative Nutrihealth study (ClinicalTrials.gov: NCT03284840), reproduced from (20) with approval of the authors.

3.1.1.1 Serum 25(OH)D concentrations in older adults

The population of healthy older adults in our review included three research clusters. The first cluster (28, 29) represented 80 adults aged between 51 and 93 years, living at home or institutionalised; the second cluster (20) included nationally representative data of 155 people aged 65 to 75 years, and the third cluster (30) represented 457 postmenopausal women aged from 55 to 90 years. Average serum concentrations of 25(OH)D were 33.0 ± 27.0 nmol/L,

47.7 nmol/L (95%CI: 43.9-51.5), and 48.2 ± 27.4 nmol/L, respectively. In the nationally representative cluster, seasonal 25(OH)D levels were reported. During the extended summer (May-October) and the extended winter (November-April), mean levels were 60.1 (95% CI: 54.0-66.2) and 39.0 nmol/L (95% CI: 35.0-43.0), respectively. In addition to the influence of seasonal variations, 25(OH)D levels were also significantly negatively affected by the body mass index (BMI). Seasonal variation of 25(OH)D was also observed in the third cluster, where decreasing levels of 25(OH)D concentration with increasing age were also reported (30).

3.1.1.2 Serum 25(OH)D concentrations in pregnant women

The review covered two research clusters of healthy pregnant women. One included data collected in the 'My Milk' research project ('The role of human milk in the development of breastfed child's intestinal microbiota') (17, 31, 32). We only included publications with a full study sample (17), as two references refer to a smaller subsample size (31, 32). The second cluster is described in six different publications (33-38). Both research clusters reported serum 25(OH)D levels throughout the year, with mean levels 43.4 ± 23.8 nmol/L and 74.7 ± 27.5 nmol/L for the first and second clusters, respectively.

3.1.1.3 Serum 25(OH)D concentrations in other populations

One research cluster included data on new-born infants (33-38); one was with healthy young athletes (16.9 ± 4.4 years) (39), and three clusters included paediatric patients (3.8-18 years) (40-44); one of these included a healthy control group (40-42). In infants, mean serum 25(OH)D concentrations were 55.2 ± 30.9 nmol/L and varied notably throughout the year. The lowest was observed in March (36.4 ± 22.6 nmol/L), and highest in September (72.9 ± 31.7 nmol/L) and the concentrations were approximately 10 nmol/L higher than in mothers (33-38). In young athletes (39), serum 25(OH)D was measured in samples collected in April (67.6 ± 36.2 nmol/L); there was no significant difference between a group of swimmers and artistic gymnasts.

Eighteen research clusters included data from adult patient populations: five research clusters included chronic kidney disease patients (24, 25, 45-50), three research clusters included type 2 diabetics (26, 51, 52), two clusters included patients with bone metabolism disorder or trauma (53, 54), two clusters included mixed data from University Medical Centre Ljubljana laboratories (55, 56), and single-cluster studies included patients after heart transplant (57), HIV patients (58), chronic intestinal failure patients on long-term home parenteral nutrition (59), and patients treated with antiepileptic drugs (27).

3.1.2 Distribution of serum 25(OH)D concentrations in the healthy population above 18 years

Six clusters reported data for 25(OH)D insufficiency (<50 nmol/L) in eight different (general), populations including pregnant women and newborns (17, 20, 21, 27, 33-39). The prevalence of 25(OH)D insufficiency in year-round samples varied from 14% in pregnant women (17) to 63% in a nationally representative sample of the elderly (20). However, due to notable seasonal differences, we focused on studies that reported seasonal results. Four clusters were included in meta-analyses, since they reported winter- and summertime insufficient serum 25(OH)D levels (<50 nmol/L) in at least 50 healthy subjects (Figure 3). The pooled summertime proportion of 25(OH)D insufficiency is 0.31 (95% CI 0.17, 0.46), where we should note heterogeneity between studies. The pooled wintertime proportion of 25(OH)D insufficiency is 0.77 (95% CI 0.68, 0.85), again with notable heterogeneity

between studies. This can be explained by the fact that these analyses included different population groups, for which very different 25(OH)D insufficiency proportions are also presented by studies that used the same design (i.e., adults and elderly in Hribar et al. (20)). Nevertheless, considerable seasonal variation in 25(OH)D insufficiency was observed, with much higher prevalence rates during wintertime ($p < 0.001$; $z = 5.62$).

3.1.3 Non-optimal 25(OH)D status

Seven research clusters (general populations) reported on the prevalence of non-optimal 25(OH)D concentrations using thresholds comparable with the Endocrine Society's guidelines (75 nmol/L) (3). Four clusters with at least 50 healthy subjects were used for meta-analyses. As seen in Figure 4, very few subjects were in the range of optimal 25(OH)D levels, even during summertime. Analyses showed a significantly different ($p < 0.001$; $z = 23.3$) pooled proportion of non-optimal 25(OH)D status between




Figure 3. Forest plot of studies with at least 50 healthy subjects evaluating the proportion of insufficient serum 25(OH)D level (<50 nmol/L). *Note: In one study (Soltirovska et al. 2015) sampling was carried out across the year, but the majority of the samples were collected during extended summer.




Figure 4. Forest plot of studies with at least 50 healthy subjects evaluating the proportion of non-optimal serum 25(OH)D level (<75 nmol/L). *Note: In one study (Dovnik et al. 2015), the non-optimal 25(OH)D level was set at 80 nmol/L.

summertime (0.69; 95% CI 0.51, 0.86), and wintertime (0.97; 95% CI 0.95, 0.99). While the summertime reports are quite heterogenous, this was not the case for wintertime ($z=118.4$), where the prevalence of non-optimal 25(OH)D status was above 95% in all reported studies.

3.2 Dietary intake of vitamin D

In total, sixteen publications reported dietary VitD intakes (Figure 1); of these, ten were journal articles and six were academic theses (details provided in Supplementary table 2). Most ($n=12$) publications referred to a healthy population (17, 19, 60-71), while two were on patient populations (72, 73). When considering the age and pregnancy status of the participants, one publication referred to children (<10 years) (60), six to adolescents (10-19 years) (17, 19, 61-65), eight to adults (>19 years) (19, 64-73), and one publication to pregnant women (17). One journal article covered both adolescents and adults (65), and one adolescents, adults, and the elderly (19). Two studies reported VitD intake separately - for food intake,

and intake with food and supplements (66, 71). Data on intake of VitD was collected using different methods: a) 24-hour dietary recalls (24h-recall) ($n=4$), b) different kinds of food frequency questionnaires (FFQ) ($n=4$), and c) dietary record (DR) ranging from 3 to 7 days ($n=6$). Two studies investigated VitD intake using both FFQ and 24h-recall (Supplementary table 2).

In the adult population, mean daily VitD intake ranged from 1.1 μg in older adults living in residential home (69, 70), to 14.2 μg in women after dietary intervention, with included food supplements and plant-based meal replacements ($n=109$; 3DDR) (66). In a teenage population mean VitD intake varied from 1.0 μg in teenage girls ($n=180$; 2 non-consecutive 24h-recalls) (64) to 4.0 μg in teenage boys ($n=1,010$; FFQ, portion size-adjusted with 3-day DR) (61). Preschool children had an average intake of 1.1 μg VitD per day ($n=129$; 3-day DR) (60). Data were available for two adult female patient populations. Celiac patients had an average intake of 2.6 μg VitD per day ($n=40$; 3-day DR) (72) and osteoporosis patients ($n=43$) had

an intake of 1.8 µg (FFQ) or 1.4 µg (24h-recall), depending on the dietary assessment method (73).

The Si.Menu project, conducted on a nationally representative sample, included 468 adolescents (10-17 years), 364 adults (18-64 years), and 416 elderly individuals (65-74 years). VitD intakes were estimated using a combination of two 24h-recalls adjusted with a food propensity questionnaire. Mean daily VitD intakes for adolescents, adults and the elderly were 2.7 µg, 2.9 µg, and 2.5 µg, respectively, and we should also note that the amount of VitD supplements was not recorded. Energy intake was a significant factor across all age groups, sex was determined as a significant factor, affecting VitD intakes in adolescents and adults, with lower VitD intakes observed in females. The most important dietary sources of VitD were eggs, fish, and fish products, and meat and meat products (19).

4 DISCUSSION

Increased interest in VitD research (74) is related to a growing body of evidence that connects sufficient 25(OH)D status with better health outcomes (75-78), while highlighting the high prevalence of VitD deficiency in different populations (13). This topic has recently gained even more attention due to the possible associations of VitD status with health outcomes of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infected patients (78-82).

In Slovenia, there is currently no official policy on food fortification and supplementation of diets with VitD in the general population. To support further public health decision-making, the Ministry of Health appointed a national expert working group on guidelines for sufficient vitamin D levels in the Slovenian population, under the umbrella of the National Institute of Public Health. To support this activity, this work aimed to review scientific publications, reporting dietary intake and/or 25(OH)D status in various Slovenian population subgroups. We reviewed all available publications with novel research data collected in Slovenia, including original scientific articles, conference papers, and academic theses.

4.1 Vitamin D status

We used reported data on serum 25(OH)D levels as a biomarker of 25(OH)D status across various populations in Slovenia, but due to methodological and analytical differences in different studies, data should be interpreted with caution. For example, in different studies, blood samples were collected in different seasons, or non-evenly distributed across the calendar year. We should

also mention that different assays were used for the quantification of 25(OH)D in different studies.

Only one study reported nationally representative data on 25(OH)D concentrations in adults (18-64 years) and the elderly (65-75 years) (20). The prevalence of insufficiency was 58.2% and 62.9% for adults and the elderly, respectively, and 83.3% and 84.4% of adults and the elderly, respectively, had 25(OH)D values below the optimal level. The levels of 25(OH)D were influenced by sex and physical activity, and most importantly by calendar season. In a much smaller cluster (n=22), 59.1% were insufficient; 90.1% were below the optimal status (27), and in a cluster referring to self-paying subjects from the laboratory information system (n=73), 58.9% were below optimal status (23). There were notable differences in the reported prevalence of 25(OH)D insufficiency (<50 nmol/L) and below optimal status (<75 nmol/L) in different studies, as these were conducted on different population groups and in different seasons. In 2016, Cashman et al. made a review of the available European data using the VitD Standardization Program (n=55,844) and reported that, overall, 40% of the European population had insufficient 25(OH)D status (74). In Southern Europe, more than one-third of studies had mean 25(OH)D concentrations below 50 nmol/L (83), and according to recent 2019 European review article, insufficiency occurs in 30-60% of the populations in Western, Southern and Eastern Europe (84). These results are comparable with those reported for various European countries (74, 83, 85). In the French adult population (18-89 years; population-based cohort), the prevalence of insufficiency was 34.6%, and 80.3% had non-optimal serum 25(OH)D concentrations (86). Similar results were observed in Spain (23-72 years; 43-49°N; nationally representative cohort), where the reported prevalence of insufficiency and non-optimal 25(OH)D status was 34.7% and 88.9%, respectively (87). In Germany (18-79 years) (74), the prevalence of 25(OH)D concentrations below 50 nmol/L was 54.5%, and below 75 nmol/L was 90.9%. Researchers from Portugal (>18 years; 32-42°N; nationally representative cohort) reported serum 25(OH)D insufficiency in 45.4% of the population and below optimal status in 96.4% (88). Even more worrying results came from Ukraine (20-95 years; 44-52°; nationally representative cohort), where the prevalence of insufficiency was 81.8%, with 95.4% below optimal status. Nationally representative data from Europe show that the Slovenian population is among those with a higher prevalence of insufficiency across the year.

We should also note seasonal variations, which were reported in a nationally representative Slovenian study (20). Higher serum 25(OH)D concentrations were observed in the extended summer period from May to October, with insufficient 25(OH)D status in 25.3% of adults and 40.2% of the elderly. A much more concerning situation was reported for an extended winter period from November

to April, when 81.6% of adults and 78.8% of the elderly had insufficient 25(OH)D status. Seasonal variation is also reported in other European countries. In Ukraine, the prevalence of insufficiency varied during the year. In the summer period (May–October) the prevalence for 25(OH)D insufficiency was 79.5%, and 85.8% during the winter period (November–April) (89). Seasonal variation was also noted in Portugal (>18 years; 32–42°N; nationally representative cohort), where in summer (July–September) the prevalence of insufficiency was 43.2%, rising to 76.0% in winter (January–March) (88). In Southern Italy (40°N; longitudinal) (90), seasonal variation in insufficiency was even more pronounced, and it varied from 4.4% in summer (August) to 70% in winter (February). In the present review, seasonal variation was also noted in pregnant women (17, 33–38) and across other populations (22, 23, 30, 43, 54).

4.2 Dietary vitamin D intake

The abovementioned seasonal variability in 25(OH)D status showed that in Slovenia (45–46°N), endogenous synthesis of VitD is not sufficient during wintertime, as was also observed in other regions (10). In such cases, dietary intake of VitD becomes very important. However, very few foods are rich in VitD (e.g., fatty fish), and these are not common in the Slovenian diet. Achieving adequate VitD intake is therefore challenging without the use of fortified foods and/or VitD dietary supplementation. In a recent Slovenian study (91) it was established that the 33% of population was taking VitD supplements (before the SARS-CoV-2 epidemic), and we should consider this when interpreting the data. In Slovenia, we follow the D-A-CH reference values for VitD intake, which recommend a daily intake of 20 µg VitD, for all above the age of 1 year in the absence of endogenous synthesis (12, 92, 93). Various dietary intake assessment tools have been used (FFQ, DR, and 24h-recall) to estimate dietary intake levels, and we should therefore compare the results with some caution. At the same time, the food database is often incomplete or lacking compositional data regarding VitD content in a certain food/dish, which may result in underestimation of dietary intake of VitD (94).

We reviewed sixteen publications reporting dietary intake of VitD in different Slovenian populations, one of which was nationally representative for adolescents, adults, and the elderly (19). All studies reported VitD intakes well below reference values across all the populations. In the adult and elderly population, daily mean VitD intakes ranged from 1.1 µg to 14.2 µg, the latter with supplements and plant-based meal replacements. A study by Lichthammer et al. (65) (FFQ) included populations from various Central-Eastern European countries (Austria, Poland, Hungary, and Slovenia). The mean daily VitD intake was very low in all countries, and was the lowest in Austria with 2.2 µg and then immediately followed by Slovenia with 2.6 µg.

A similar intake for the Austrian population was found in another study (2.1 µg per day; n=4,972; various dietary intake tools) (95). In a Slovenian nationally representative study, where the amount of supplementation was not included, mean daily VitD intakes were 2.7 µg, 2.9 µg, and 2.5 µg in adolescents, adults, and the elderly, respectively (19). A review of European studies on VitD intake showed a mean VitD intake of 3 to 5 µg per day (13, 96–99), with higher intakes in Northern Europe (up to 11 µg/day) and lower in Southern Europe (13, 84, 97, 99, 100). This can be explained by the high consumption of sea fish and VitD enriched foods in Northern Europe. Our review also covered six children or adolescent populations. In teenage populations, VitD intake varied from 1.0 µg in teenage girls (2x24h-recall) (64) to 4.0 µg in teenage boys (FFQ, portion size-adjusted with 3-day DR) (61). These results are comparable to an Austrian study (95) and other European studies (96, 98, 101, 102). A European nutrition and health report (100) also highlighted that VitD intake in children was generally very low, with the exception of some Nordic countries due to the general vitamin D food fortification policy.

4.3 Research gaps

This review has identified several studies reporting dietary intake and/or 25(OH)D status in various Slovenian populations, but the majority were conducted using the convenience sampling method and are therefore not nationally representative. Nevertheless, the reviewed results showed considerable seasonal variability in serum 25(OH)D levels in different populations, with a particularly high prevalence of 25(OH)D deficiency during the wintertime.

Only one study investigated 25(OH)D status in a nationally representative sample of adults and reported insufficient wintertime 25(OH)D levels in about 80% of adults and 25(OH)D deficiency in about 40%. The study sample also included those who reported vitamin D supplementation. The majority of other studies also showed that various population groups in Slovenia are mostly at risk in wintertime with regard to insufficient 25(OH)D status according to the WHO criteria (103). This literature review has also identified some important research gaps. The SARS-CoV-2 pandemic has significantly affected the dietary habits of Slovenians, and increased the consumption of food supplements and medicines with VitD. This might have affected 25(OH)D status in some populations studied in the last year. In addition, some population groups have not been sufficiently investigated. There is very limited data on the 25(OH)D status in healthy children and adolescents, and a lack of data on institutionalised populations, such as those living in elderly care institutions, who are rarely

exposed to sunlight and therefore at an even higher risk of 25(OH)D deficiency.

We also reviewed the data on dietary VitD intake in Slovenia. Several studies were identified that estimated VitD intakes in different populations using a variety of study designs and methods. One study reported nationally representative data. All studies reported very low VitD intakes. While the recommended daily intake of VitD in adults is 20 µg in the absence of endogenous synthesis, most studies reported dietary intakes below 5 µg. The majority of the studies did not estimate intakes of VitD with medicines and food supplements or fortified foods, though the latter practice has been negligible in Slovenia. VitD food supplementation practices should be further investigated in the future.

5 CONCLUSION

The general observation of our review is that various population groups across Slovenia are at high risk of 25(OH)D insufficiency and deficiency, particularly during the wintertime. Regarding VitD intake, the majority of included studies reported daily intakes below 5 µg. We also identified key research gaps that need to be addressed to support further public health decision-making.

ACKNOWLEDGMENTS

We acknowledge the support of the members of the national expert working group on guidelines for sufficient vitamin D levels in the Slovenian population, particularly Marija Pfeifer, Andraž Dovnik, and Aneta Soltirovska Šalamon. We also acknowledge the support of Joško Osredkar in data collection.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results. I.P. has led and participated in various other research projects in nutrition, public health, and food technology that were (co)funded by the Slovenian Research Agency, Ministry of Health of the Republic of Slovenia, the Ministry of Agriculture, Forestry, and Food of the Republic of Slovenia, and, for specific applied research projects, also by food businesses. I.P., K.Ž., and E.B. are members of a national expert working group on guidelines for sufficient

vitamin D levels in the Slovenian population.

FUNDING

The study was conducted within the national research program P3-0395 (Nutrition and Public Health) and project L7-1849 (Challenges in achieving adequate vitamin D status in the adult population) and supported by the Slovenian Research Agency and Ministry of Health of the Republic of Slovenia.

ETHICAL APPROVAL

This is a review article. No ethical approval is needed.

REFERENCES

- Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007;357(3):266-81. doi: 10.1056/NEJMra070553.
- Holick MF. Vitamin D: physiology, dietary sources and requirements. In: Caballero B, editor. *Encyclopedia of human nutrition*. 2nd ed. Oxford: Elsevier, Academic Press, 2005:368-77.
- Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2011;96(7):1911-30. doi: 10.1210/jc.2011-0385.
- Gallagher JC, Sai AJ. Vitamin D insufficiency, deficiency, and bone health. *J Clin Endocrinol Metab*. 2010;95(6):2630-3. doi: 10.1210/jc.2010-0918.
- Chowdhury R, Kunutsor S, Vitezova A, Oliver-Williams C, Chowdhury S, Kieft-de-Jong JC, et al. Vitamin D and risk of cause specific death: systematic review and meta-analysis of observational cohort and randomised intervention studies. *BMJ*. 2014;348:g1903. doi: 10.1136/bmj.g1903.
- Chang SW, Lee HC. Vitamin D and health - the missing vitamin in humans. *Pediatr Neonatol*. 2019;60(3):237-44. doi: 10.1016/j.pedneo.2019.04.007.
- Seamans KM, Cashman KD. Existing and potentially novel functional markers of vitamin D status: a systematic review. *Am J Clin Nutr*. 2009;89(6):1997s-2008s. doi: 10.3945/ajcn.2009.27230D.
- Institute of Medicine (US) Committee to Review Dietary Reference Intakes for Calcium and Vitamin D. *The National Academies Collection: reports funded by National Institutes of Health*. In: Ross AC, Taylor CL, Yaktine AL, Del Valle HB, editors. *Dietary reference intakes for calcium and vitamin D*. Washington: National Academies Press, National Academy of Sciences, 2011.
- Dietary reference values for vitamin D. *EFSA J*. 2016;14(10):e04547. doi: 10.2903/j.efsa.2016.4547.
- O'Neill CM, Kazantzidis A, Ryan MJ, Barber N, Sempos CT, Durazo-Arvizu RA, et al. Seasonal changes in vitamin D - effective UVB availability in Europe and associations with population serum 25-hydroxyvitamin D. *Nutrients*. 2016;8(9):533. doi: 10.3390/nu8090533
- World Health Organization. *Vitamin and mineral requirements in human nutrition*. 2nd ed. Geneva: World Health Organization, 2005.
- German Nutrition S. New reference values for vitamin D. *Ann Nutr Metab*. 2012;60(4):241-6. doi: 10.1159/000337547.
- Roman Viñas B, Ribas Barba L, Ngo J, Gurinovic M, Novakovic R, Cavelaars A, et al. Projected prevalence of inadequate nutrient intakes in Europe. *Ann Nutr Metab*. 2011;59(2-4):84-95. doi: 10.1159/000332762.

14. Benedik E. Sources of vitamin D for humans. *Int J Vitam Nutr Res*. 2021;8. doi: 10.1024/0300-9831/a000733.
15. Calvo MS, Whiting SJ. Survey of current vitamin D food fortification practices in the United States and Canada. *J Steroid Biochem Molecular Biology*. 2013;136:211-3. doi: 10.1016/j.jsbmb.2012.09.034.
16. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097. doi: 10.1371/journal.pmed.1000097.
17. Soltirovska Salamon A, Benedik E, Bratanič B, Velkavrh M, Rogelj I, Fidler Mis N, et al. Vitamin D status and its determinants in healthy Slovenian pregnant women. *Ann Nutr Metab*. 2015;67(2):96-103. doi: 10.1159/000439093.
18. Marc J. Razvoj analitičnega postopka sočasnega določanja koncentracij vitamina D3 in njegovih presnovkov v krvnem serumu pacientov na peritonealni in hemodializi Ljubljana: Univerza v Ljubljani, Fakulteta za naravoslovje in tehnologijo, 1993.
19. Hribar M, Hristov H, Lavriša Ž, Koroušić Seljak B, Matej G, Blaznik U, et al. Vitamin D intake in Slovenian adolescents, adults, and the elderly population. *Nutrients*. 2021;13(10):3528. doi: 10.3390/nu13103528.
20. Hribar M, Hristov H, Gregorič M, Blaznik U, Zaletel K, Oblak A, et al. Nutrihealth study: seasonal variation in vitamin D status among the Slovenian adult and elderly population. *Nutrients*. 2020;12(6):1-17. doi: 10.3390/nu12061838.
21. Žmitek K, Hribar M, Hristov H, Pravst I. Efficiency of vitamin D supplementation in healthy adults is associated with body mass index and baseline serum 25-hydroxyvitamin D level. *Nutrients*. 2020;12(5). doi: 10.3390/nu12051268.
22. Osredkar J, Marc J. Vitamin D in presnovki: fiziologija, patofiziologija in referenčne vrednosti. *Med Razgl*. 1996;35(4):543-65.
23. Avberšek-Lužnik I. Vpliv starosti, spola in letnega časa na koncentracije vitamina D pri naključnih preiskovancih gorenjske regije. In: Medpoklicno in medsektorsko povezovanje za razvoj kakovostne in odzivne zdravstvene obravnave: 9. mednarodna znanstvena konferenca. Bled, 2016:317-23.
24. Pajek M, Čuk I, Mlinšek G, Osredkar J, Pajek J. Physical function and 25-hydroxyvitamin D in dialysis patients - lessons learned from the Slovenian DIAGIB study. *Clin Nephrol*. 2017;88(1):S48-S52.
25. Šturma A. Povezanost serumskih koncentracij 25-hidroksi holekalciferola z gibalno zmogljivostjo dializnih bolnikov. Ljubljana: Univerza v Ljubljani, Fakulteta za farmacijo, 2017.
26. Trdan B. Povezanost koncentracij vitamina D in glukoze pri bolnikih s sladkorno boleznijo. Ljubljana: Univerza v Ljubljani, Fakulteta za farmacijo, 2016.
27. Troškot M, Duhovnik T. Preskrbljenost bolnikov, zdravljenih s protiepileptičnimi zdravili (karbamazepin, okskarbazepin), z vitaminom D in učinki nadomeščanja vitamina D na kostne kazalce. Ljubljana: Univerza v Ljubljani, 2013.
28. Gradišnik L. Dejavniki tveganja v povezavi s pomanjkanjem vitamina D pri starejših. Maribor: Univerza v Mariboru, 2017.
29. Velnar T, Gradišnik L, Železnik D, Krajnc B, Maver U. Dejavniki tveganja za pomanjkanje vitamina D pri starejših. *Farm Vestn*. 2018;69(3):188-94.
30. Pajk J. Sprememba serumske koncentracije 25-hidroksi vitamina D v pomenopavznem obdobju. Ljubljana: Univerza v Ljubljani, Fakulteta za farmacijo, 2011.
31. Duh K, Budič P. Vpliv antropometričnih mer, vitamina D in adiponektina pri materah na rast, mineralno kostno gostoto ter pojavnost alergijskih bolezni pri otrocih. Ljubljana: Univerza v Ljubljani, 2017.
32. Velkavrh M, Paro Panjan D, Benedik E, Fidler Mis N, Godnov U, Soltirovska Salamon A. The influence of maternal levels of vitamin D and adiponectin on anthropometrical measures and bone health in offspring. *Prilozi / Makedonska akademija na naukite i umetnostite, Oddelenie za medicinski nauki*. 2019;40(3):91-8.
33. Dovnik A. Sezonska nihanja vitamina D pri nosečnicah in novorojenčkih v mariborski regiji. Maribor: Univerza v Mariboru, 2016.
34. Dovnik A, Mujezinović F, Treiber M, Pečovnik-Balon B, Arko D, Takač I. Vpliv maternalnega vitamina D na pojav zlomov kosti in displazije kolčkov pri novorojenčku. In: VI. Osteološki dnevi. Maribor, 2017:201-8.
35. Dovnik A, Mujezinović F, Treiber M, Pečovnik-Balon B, Gorenjak M, Maver U, et al. Seasonal variations of vitamin D concentrations in pregnant women and neonates in Slovenia. *Eur J Obstet Gynecol Reprod Biol*. 2014;181:6-9. doi: 10.1016/j.ejogrb.2014.07.019.
36. Dovnik A, Mujezinović F, Treiber M, Pečovnik-Balon B, Gorenjak M, Maver U, et al. Determinants of maternal vitamin D concentrations in Slovenia: a prospective observational study. *Wien Klin Wochenschr*. 2017;129(1-2):21-8. doi: 10.1007/s00508-016-1142-2.
37. Dovnik A, Mujezinović F, Treiber M, Pečovnik-Balon B, Takač I. Sezonska nihanja koncentracije vitamina D pri nosečnicah in novorojenčkih v mariborski regiji. In: Novakovi dnevi, Nova Gorica, 2015:98-105.
38. Dovnik A, Mujezinović F, Treiber M, Pečovnik-Balon B, Takač I. Vpliv maternalnega vitamina D na zdravje novorojenčka. In: Čokolič M, Pečovnik-Balon B, Komadina R, Zavratnik A, editors. Osteološki dnevi. Maribor: Univerzitetni klinični center Maribor, 2015:175-82.
39. Jakše B, Sekulič D, Jakše B, Čuk I, Šajber D. Bone health among indoor female athletes and associated factors: a cross-sectional study. *Res Sports Med*. 2020;28(3):314-23. doi: 10.1080/15438627.2019.1696344.
40. Ferant Ž, Kozar N, Dolinšek J, Šikič Pogačar M, Mičetič-Turk D. Bone mineral density and body composition in pediatric patients with celiac disease and inflammatory bowel disease. *Health Med*. 2014;8(3):319-28.
41. Ferant Ž, Kozar N. Mineralna kostna gostota pri otrocih in mladostnikih s celiakijo in kronično vnetno črevesno boleznijo. Maribor: Univerza v Mariboru, 2012.
42. Šikič Pogačar M, Dolinšek J, Ferant Ž, Kozar N, Mičetič-Turk D. Raven vitamina D pri otrocih in mladostnikih s celiakijo in kroničnimi vnetnimi črevesnimi boleznimi v severovzhodni Sloveniji. Kongres klinične prehrane in presnovne podpore. Portorož: Slovensko združenje za klinično prehrano, 2013:189-91.
43. Blazina Š, Bratanič N, Širca-Čampa A, Blagus R, Orel R. Bone mineral density and importance of strict gluten-free diet in children and adolescents with celiac disease. *Bone*. 2010;47(3):598-603.
44. Tesovnik T, Kovač J, Hovnik T, Kotnik P, Battelino T, Trebušak Podkrajšek K. Association of average telomere length with body-mass index and vitamin D status in juvenile population with type 1 diabetes. *Zdr Varst*. 2015;54(2):74-8. doi: 10.1515/sjph-2015-0011.
45. Ekart R, Vodošek Hojs N, Bevc S, Dvoršak B, Jakopin E, Hren M, et al. 25-hydroxyvitamin D and 24-hour ambulatory pulse pressure in patients with chronic kidney disease. 23rd European Meeting on Hypertension and Cardiovascular Protection, June 14-17. Milan, 2013.
46. Jakopin E. Pomanjkanje vitamina D pri bolnikih zdravljenih s kronično hemodializo in njegovo nadomeščanje z visokimi odmerki holekalciferola. Maribor: Univerza v Mariboru, 2013.
47. Jakopin E, Pečovnik-Balon B, Ekart R, Gorenjak M. High-dose cholecalciferol supplementation for vitamin D deficiency in haemodialysis patients. *J Int Med Res*. 2011;39(3):1099-106. doi: 10.1177/147323001103900345.
48. Pečovnik-Balon B, Jakopin E, Bevc S, Knehtl M, Gorenjak M. Vitamin D as a novel nontraditional risk factor for mortality in hemodialysis patients. *Ther Apher Dial*. 2009;13(4):268-72.
49. Humar T. Antiproteinurično delovanje holekalciferola pri IgA nefropatiji. Ljubljana: Univerza v Ljubljani, Fakulteta za farmacijo, 2015.
50. Osredkar J, Humar T, Kovač D. Uporaba vitamina D pri zdravljenju pacientov z IgA nefropatijo. In: Skitek M, Černe D, Marc J, Štiglic AF, Kuželički NK, editors. Jesenovčevi dnevi. Ljubljana, 2015:110-28.
51. Krajnc M, Čokolič M, Pečovnik-Balon B. Coronary artery calcification, femoral neck BMD and serum 25-hydroxy vitamin D in type 2 diabetics. In: European Congress on Osteoporosis and Osteoarthritis (ESCEO13-

- IOF), 17-20 April 2013. Rome: International Osteoporosis Foundation and National Osteoporosis Foundation, 2013:S200-S1.
52. Kšela U, Zavrtnik A. Pomanjkanje vitamina D pri bolnikih s sladkorno boleznijo tipa 2 - naše izkušnje. In: IV. Osteološki dnevi. Maribor: Univerzitetni klinični center Maribor, 2013: 203-7.
 53. Vindišar F, Goličnik B, Komadina R, Marc J. Znižana hitrost glomerulne filtracije kot dejavnik tveganja za osteoporozo? In: II. Osteološki dnevi. Maribor: Medicinska fakulteta Univerze v Mariboru, 2009:154-62.
 54. Krajnc M, Čokolič M, Homšak E. Seasonal vitamin D variation in the area of Maribor. In: Slovenski endokrinološki kongres z mednarodno udeležbo, Bled. Ljubljana: Združenje endokrinologov Slovenije pri Slovenskem zdravniškem društvu, Slovensko osteološko društvo, 2018:200-1.
 55. Gros K. Določitev orientacijskih referenčnih vrednosti vitamina D (25-OH) v serumu. Ljubljana: Univerza v Ljubljani, Fakulteta za farmacijo, 2008.
 56. Salkič M. Klinični pomen porušenega razmerja kalcija in fosforja v serumu, ter korigiranega kalcija v serumu v odvisnosti od vitamina D. Ljubljana: Univerza v Ljubljani, Fakulteta za farmacijo, 2016.
 57. Rakuša M, Vrtovec B, Pogljajen G, Janež A, Jensterle Sever M. Endocrine disorders after heart transplantation : national cohort study. *BMC Endocrine Disorders*. 2020;20(54):10.
 58. Tomažič J, Ul K, Volčanšek G, Gorenšek S, Pfeifer M, Karner P, et al. Prevalence and risk factors for osteopenia/osteoporosis in an HIV-infected male population. *Wien Klin Wochenschr*. 2007;119(21-22):639-46. doi: 10.1007/s00508-007-0844-x.
 59. Rondaij T, Kozjek NR, Popovic P, Jordan T. Vitamin D deficiency in patients with chronic intestinal failure on home parenteral nutrition. *Clinical Nutrition ESPEN*. 2021;42:258-61. doi: 10.1016/j.clnesp.2021.01.026.
 60. Poličnik R, Pokorn D, Kulnik D, Mičetič-Turk D, Hlastan-Ribič C. Energy and nutrient intake among pre-school children in Central Slovenia. *Acta Aliment*. 2013;42(3):291-300. doi: 10.1556/AAlim.2012.0002.
 61. Fidler Mis N, Kobe H, Štimatec M. Dietary intake of macro-and micronutrients in Slovenian adolescents: comparison with reference values. *Ann Nutr Metab*. 2012;61(4):305-13. doi: 10.1159/000342469.
 62. Gregorič M. Ocena prehranskega vnosa pri mladostnikih z vidika varovanja zdravja. Ljubljana: Univerza v Ljubljani, Biotehniška fakulteta, 2015.
 63. Koch V, Gregorič M. Prehranska kakovost zajtrka slovenskih srednješolcev. *Zdr Varst*. 2009;48:131-42.
 64. Zdešar Kotnik K. Smiselnost uporabe vitaminskih in mineralnih prehranskih dopolnil pri mladostnikih. Ljubljana: Univerza v Ljubljani, Biotehniška fakulteta, 2019.
 65. Lichthammer A, Nagy B, Orbán C, Tóth T, Csajbók R, Molnár S, et al. A comparative study of eating habits, calcium and Vitamin D intakes in the population of Central-Eastern European countries. *New Med*. 2015;19(2):66-70. doi: 10.5604/14270994.1169799.
 66. Jeretina B. Dejavniki vpliva na pojav prezgodnje menopavze. Ljubljana: Univerza v Ljubljani, Zdravstvena fakulteta, 2019.
 67. Urh N, Babnik K, Rebec D, Poklar Vrtovec T. Ocena prehranskega stanja starejših v socialnovarstvenem zavodu. *Obz Zdrav Nege*. 2017;51(3):207-18.
 68. Juvan S. Kakovost vegetarijanske prehrane med študenti. Ljubljana: Univerza v Ljubljani, Biotehniška fakulteta, 1997.
 69. Gregorič M, Kotnik KZ, Pigac I, Gabrijelčič Blenkuš M. A web-based 24-h dietary recall could be a valid tool for the indicative assessment of dietary intake in older adults living in Slovenia. *Nutrients*. 2019;11(9):2234. doi: 10.3390/nu11092234.
 70. Pigac I. Določanje vnosa hranil pri oskrbovancih v domovih za starejše. Ljubljana: Univerza v Ljubljani, Biotehniška fakulteta, 2016.
 71. Jakše B, Jakše B, Pinter S, Pajek J, Godnov U, Fidler Mis N. Nutrient and food intake of participants in a Whole-Food Plant-Based Lifestyle Program. *J Am Coll Nutr*. 2021;40(4):333-48. doi: 10.1080/07315724.2020.1778584.
 72. Kocuvan Mijatov MA, Mičetič-Turk D. Dietary intake in adult female coeliac disease patients in Slovenia. *Zdr Varst*. 2016;55(2):86-93. doi: 10.1515/sjph-2016-0013.
 73. Lopert B. Ocena prehranskega vnosa kalcija in vitamina D pri osteoporoznih bolnikih. Ljubljana: Univerza v Ljubljani, Biotehniška fakulteta, 2019.
 74. Cashman KD, Dowling KG, Škrabáková Z, Gonzalez-Gross M, Valtueña J, De Henauw S, et al. Vitamin D deficiency in Europe: pandemic? *Am J Clin Nutr*. 2016;103(4):1033-44. doi: 10.3945/ajcn.115.120873.
 75. Autier P, Boniol M, Pizot C, Mullie P. Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol*. 2014;2(1):76-89. doi: 10.1016/s2213-8587(13)70165-7.
 76. Zittermann A. Vitamin D in preventive medicine: are we ignoring the evidence? *Br J Nutr*. 2003;89(5):552-72. doi: 10.1079/BJN2003837.
 77. Zittermann A, Pilz S, Hoffmann H, Marz W. Vitamin D and airway infections: a European perspective. *Eur J Med Res*. 2016;21:1-10. doi: 10.1186/s40001-016-0208-y.
 78. Autier P, Mullie P, Macacu A, Dragomir M, Boniol M, Coppens K, et al. Effect of vitamin D supplementation on non-skeletal disorders: a systematic review of meta-analyses and randomised trials. *Lancet Diabetes Endocrinol*. 2017;5(12):986-1004. doi: 10.1016/s2213-8587(17)30357-1.
 79. Radujkovic A, Hippchen T, Tiwari-Heckler S, Dreher S, Boxberger M, Merle U. Vitamin D deficiency and outcome of COVID-19 patients. *Nutrients*. 2020;12(9):2757. doi: 10.3390/nu12092757.
 80. Merzon E, Tworowski D, Gorohovski A, Vinker S, Golan Cohen A, Green I, et al. Low plasma 25(OH) vitamin D level is associated with increased risk of COVID-19 infection: an Israeli population-based study. *FEBS J*. 2020;287(17):3693-702. doi: 10.1111/febs.15495.
 81. Carpagnano GE, Di Lecce V, Quaranta VN, Zito A, Buonomico E, Capozza E, et al. Vitamin D deficiency as a predictor of poor prognosis in patients with acute respiratory failure due to COVID-19. *J Endocrinol Invest*. 2020:1-7. doi: 10.1007/s40618-020-01370-x.
 82. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, et al. Evidence that vitamin D supplementation could reduce risk of Influenza and COVID-19 infections and deaths. *Nutrients*. 2020;12(4). doi: 10.3390/nu12040988.
 83. Manios Y, Moschonis G, Lambrinou C-P, Tsoutsouloupoulou K, Binou P, Karachaliou A, et al. A systematic review of vitamin D status in southern European countries. *Eur J Nutr*. 2018;57(6):2001-36. doi: 10.1007/s00394-017-1564-2.
 84. Lips P, Cashman KD, Lamberg-Allardt C, Bischoff-Ferrari HA, Obermayer-Pietsch B, Bianchi ML, et al. Current vitamin D status in European and Middle East countries and strategies to prevent vitamin D deficiency: a position statement of the European Calcified Tissue Society *Eur J Endocrinol*. 2019;180(4):P23-54. doi: 10.1530/eje-18-0736.
 85. Kiely M, Cashman KD. Summary outcomes of the ODIN project on food fortification for vitamin D deficiency prevention. *Int J Environ Res Public Health*. 2018;15(11). doi: 10.3390/ijerph15112342.
 86. Souberbielle JC, Massart C, Brailly-Tabard S, Cavalier E, Chanson P. Prevalence and determinants of vitamin D deficiency in healthy French adults: the VARIETE study. *Endocrine*. 2016;53(2):543-50. doi: 10.1007/s12020-016-0960-3.
 87. González-Molero I, Rojo-Martínez G, Morcillo S, Gutierrez C, Rubio E, Pérez-Valero V, et al. Hypovitaminosis D and incidence of obesity: a prospective study. *Eur J Clin Nutr*. 2013;67(6):680-2. doi: 10.1038/ejcn.2013.48.
 88. Duarte C, Carvalheiro H, Rodrigues AM, Dias SS, Marques A, Santiago T, et al. Prevalence of vitamin D deficiency and its predictors in the Portuguese population: a nationwide population-based study. *Arch Osteoporosis*. 2020;15(1):36. doi: 10.1007/s11657-020-0695-x.
 89. Povoroznyuk VV, Balatska NI, Muts VY, Klymovytsky FV, Sytenko OV. Vitamin D deficiency in Ukraine: a demographic and seasonal analysis. *Gerontologija*. 2012;13:191-8.
 90. Carnevale V, Modoni S, Pileri M, Di Giorgio A, Chiodini I, Minisola S, et al. Longitudinal evaluation of vitamin D status in healthy subjects

- from southern Italy: seasonal and gender differences. *Osteoporos Int.* 2001;12(12):1026-30. doi: 10.1007/s001980170012.
91. Žmitek K, Hribar M, Lavriša Ž, Hristov H, Kušar A, Pravst I. Socio-demographic and knowledge-related determinants of vitamin D supplementation in the context of the COVID-19 pandemic: assessment of an educational intervention. *Frontiers Nutr.* 2021;8(290). doi: 10.3389/fnut.2021.648450.
 92. Benedik E, Fidler Mis N. New recommendations for vitamin D intake. *Zdrav Vestn.* 2013;82:1-145-51.
 93. Nacionalni inštitut za javno zdravje. Referenčne vrednosti za energijski vnos ter vnos hranil. Accessed December 21st, 2020 at: https://www.nijz.si/sites/www.nijz.si/files/uploaded/referencne_vrednosti_2020_3_2.pdf.
 94. Benedik E, Koroušič Seljak B, Hribar M, Rogelj I, Bratanič B, Orel R, et al. Comparison of a web-based dietary assessment tool with software for the evaluation of dietary records. *Zdr Varst.* 2015;54(2):91-7. doi: 10.1515/sjph-2015-0014.
 95. Koenig J, Elmadfa I. Status of calcium and vitamin D of different population groups in Austria. *Int J Vitam Nutr Res.* 2000;70(5):214-20. doi: 10.1024/0300-9831.70.5.214.
 96. Novaković R, Cavelaars AE, Bekkering GE, Roman-Viñas B, Ngo J, Gurinović M, et al. Micronutrient intake and status in Central and Eastern Europe compared with other European countries: results from the EURRECA network. *Public Health Nutr.* 2013;16(5):824-40. doi: 10.1017/s1368980012004077.
 97. Jenab M, Salvini S, van Gils CH, Brustad M, Shakya-Shrestha S, Buijsse B, et al. Dietary intakes of retinol, beta-carotene, vitamin D and vitamin E in the European prospective investigation into cancer and nutrition cohort. *Eur J Clin Nutr.* 2009;63(Suppl 4):S150-78. doi: 10.1038/ejcn.2009.79.
 98. Spiro A, Buttriss JL. Vitamin D: an overview of vitamin D status and intake in Europe. *Nutr Bull.* 2014;39(4):322-50. doi: 10.1111/nbu.12108.
 99. Kiely M, Black LJ. Dietary strategies to maintain adequacy of circulating 25-hydroxyvitamin D concentrations. *Scand J Clin Lab Invest Suppl.* 2012;243:14-23. doi: 10.3109/00365513.2012.681893.
 100. Elmadfa I, Meyer A, Nowak V, Hasenegger V, Putz P, Verstraeten R, et al. European nutrition and health report 2009. *Forum Nutr.* 2009;62:12-3. doi: 10.1159/000242367.
 101. Mensink GB, Fletcher R, Gurinovic M, Huybrechts I, Lafay L, Serra-Majem L, et al. Mapping low intake of micronutrients across Europe. *Br J Nutr.* 2013;110(4):755-73. doi: 10.1017/S000711451200565X.
 102. Braegger C, Campoy C, Colomb V, Decsi T, Domellof M, Fewtrell M, et al. Vitamin D in the healthy European paediatric population. *J Pediatr Gastroenterol Nutr.* 2013;56(6):692-701. doi: 10.1097/MPG.0b013e31828f3c05.
 103. de Benoist B, McLean E, Egli I, Cogswell M, editors. Worldwide prevalence of anaemia 1993-2005: WHO global database on anaemia. Geneva: World Health Organization, 2008.
 104. Vujasinović M, Kunst G, Breznikar B, Rožej B, Tepeš B, Rudolf S, et al. Is pancreatic exocrine insufficiency a cause of malabsorption in patients after bariatric surgery? *J Pancreas.* 2016;17(4):402-5.

Supplementary table 1. Review of reports for vitamin D status in the Slovenian population.

References	Data collection period	Study population	N	Serum 25(OH)D			
				Mean conc.±SD or (95% CI) [nmol/L]	Prevalence of conc. < 50 nmol/L	Prevalence of conc. > 50 nmol/L	Publ. type
Avberšek-Lužnik 2016 (23)	Jan 2014 - Dec 2015	Self-paying subjects from laboratory information system in Gorenjska region Age: 19-90; mean 49 years	73	65.1±33.0 Apr-Sept: 75.3±37.1 Oct-Mar: 55.7±25.7	<25 nmol/L: 9.6% 25-75 nmol/L: 49.3%	25-75 nmol/L: 49.3% >75 nmol/L: 41.1%	B
Blazina, Bratanič et al. 2010 (43)	Oct - May (year not provided)	Children and adolescents with celiac disease on strictly compliant to a gluten-free diet (D1) and not strictly compliant to a gluten-free diet (D2): D1 age: 11.4±4.3 years; 3.8-17.6 years D2 age: 13.4±4.8 years; 3.8-20.7 years	D1: 55 D2: 19	D1: 55.1 D2: 53.9	'Common' deficiency between Dec and Apr		A

References	Data collection period	Study population	N	Serum 25(OH)D			
				Mean conc.±SD or (95% CI) [nmol/L]	Prevalence of conc. < 50 nmol/L	Prevalence of conc. > 50 nmol/L	Publ. type
Dovnik, Mujezinović et al. 2014, Dovnik, Mujezinović et al. 2015, Dovnik, Mujezinović et al. 2015, Dovnik 2016, Dovnik, Mujezinović et al. 2017, Dovnik, Mujezinović et al. 2017 (33-38)	Sept 2013	Pregnant women and neonates Age: All: 29.6±4.7 years Sept: 29.1±4.2 years Dec: 29.3±5.2 years Mar: 29.6±4.6 years Jun: 30.3±4.9 years	398	Mothers:	Mothers: <25 nmol/L: 23.6%	Mothers 50-80 nmol/L: 27.9%	A, A, B, B, B, B, C1
	Dec 2013		Mothers:	43.4±23.8			
	Mar 2014		Sept 100	Sep: 54.3±25.2	Sept: 10.0%	Sept: 40.0%	
	Dec 99		Dec: 33.3±18.6	Dec: 41.4%	Dec: 16.2%		
	Jun 2014	Mar 10	Mar: 28.5±17.1	Mar: 34.0%	Mar: 12.0%		
	Jun 99	Jun: 54.8±24.1	Jun: 9.1%	Jun: 43.4%			
		401	Neonates:	Mothers: 25-50 nmol/L: 41.5%	Mothers >80 nmol/L: 7%		
		Neonates:	55.2±30.9				
		Sept 100	Sep: 72.9±31.7	Sept: 38.0%	Sept: 12.0%		
		Dec 100	Dec: 52.5±27.3	Dec: 40.4%	Dec: 2.0%		
	Mar 101	Mar: 36.4±22.6	Mar: 53.0%	Mar: 13.0%			
	Jun 100	Jun: 59.5±24.9	Jun: 34.3%	Jun: 13.1%			
				Neonates:	Neonates:		
				<50 nmol/L: 47.6%	50-75 nmol/L: 27.4%		
					>75 nmol/L: 24.9%		
Ekart, Vodošek Hojs et al. 2013 (45)		Outpatients with chronic kidney disease in predialysis clinic Age: 60.72±12.7 years	72	55.75±27.59			B
Ferant and Kozar 2012, Šikić Pogačar, Dolinšek et al. 2013, Ferant, Kozar et al. 2014 (40-42)	Mar 2011 - Apr 2012	Pediatrics patients with celiac disease (CD) and inflammatory bowel disease (IBD) Age: 10-18 years	CD: 35 IBD: 35 K: 34	CD: 32.5 IBD: 32.6 K: 49.9			A, B, C4,
Gradišnik 2017, Velnar, Gradišnik et al. 2018 (28, 29)	Feb - Mar 2016	Institutionalized residents (IR), controls living (K) at home Age: 51-93 years	80 IR: 42 K: 38	33.0±27.0 IR: 27.0 K: 39.5	IR: <7.5 nmol: 14% K: <7.5 nmol: 13%		A, C3
Gros 2008 (55)	2006 Jan, Feb, Mar, Aug, Sept, Oct	Participants in the laboratory for analysis of hormones and tumour markers	Jan M: 54 F: 120 Feb M: 48 F: 82 Mar M: 67 F: 140 Aug F: 34 Sept M: 54 F: 81 Oct M: 42 F: 118	Jan M: 31.9±16.2 F: 40.3±39.0 Feb M: 32.2±27.3 F: 29.5±21.7 Mar M: 27.6±21.8 F: 28.7±19.9 Aug F: 59.6±35.4 Sept M: 52.2±33.8 F: 55.9±26.3 Oct M: 55.8±23.9 F: 56.0±27.6			C3

References	Data collection period	Study population	N	Serum 25(OH)D			
				Mean conc.±SD or (95% CI) [nmol/L]	Prevalence of conc. < 50 nmol/L	Prevalence of conc. > 50 nmol/L	Publ. type
Hribar, Hristov et al. 2020 (20)	2017 - 2018 Summer: May-Oct, Winter: Nov-Apr	A nationally representative sample of adults and elderly Age adults: 18-64 years; mean 46.5±13.2 Age elderly: 65-75 years; mean 68.6±2.8	Adults: 125	Adults: 50.7 (45.4-56.0)	<30 nmol/L	<75 nmol/L	A
			Summer: 52	Summer: 70.4 (62.2-78.5)	Adults: 24.9%	Adults: 83.3%	
			Winter: 73	Winter: 36.7 (32.5-40.9)	Adults winter: 40.8%	Adults winter: 98.0%	
			Elderly: 155		Adults summer: 2.6%	Adults summer: 62.6%	
			Summer: 64		Elderly: 23.5%	Elderly: 84.4%	
			Winter: 91	Elderly: 47.7 (43.9-51.5)	Elderly winter: 7.8%	Elderly winter: 92.2%	
				Summer: 60.1 (54.0-66.2)	Elderly summer: 34.6%	Elderly summer: 73.4%	
				Winter: 39.0 (35.0-43.0)			
					<50 nmol/L		
					Adults: 58.2%		
Humar 2015, Osredkar, Humar et al. 2015 (49, 50)	2000 - 2010	Patients with IgA nephropathy before and after cholecalciferol supplementation Age: 47.1±10.1 years	18	Before supplementation: 40.5±17.6			B, C3
				After supplementation: 65.0±27.3			
Jakopin, Pečovnik-Balon et al. 2011, Jakopin 2013 (46, 47)	May 2008 - May 2010	Haemodialysis patients Age: 63.3±13.5 years	101	28.6±16.7	<12 nmol/L: 16.8%	40 - 75 nmol/L: 28.7%	A, C1
					12 - 37 nmol/L: 51.5%	>75nmol/L: 3%	
					40 - 75 nmol/L: 28.7%		
Jakše, Sekulič 2020 (39)	Apr (year not provided)	Indoor female athletes: athletes engaged in weight-bearing (artistic gymnastic (AG)) and non-weight-bearing (swimming (S)) sport Age: 16.9±4.4 years	AG: 17	67.6±36.2	<50 nmol/L: 35.5%	51-75 nmol/L: 32.3%	A
			S: 14		S: 35.7%	S: 21.4%	
				AG: 65.1±36.0	AG: 35.3%	AG: 41.2%	
				S: 70.6±37.5		>75 nmol/L: 32.3%	
						S: 42.9%	
Krajnc, Čokolič et al. 2018 (54)	2014 - 2017	Hospitalised and outpatient patients mostly with osteoporosis and vitamin D risk factors for insufficiency	2082	Median: 62.9		Most individuals lower than 75 nmol/L	B

References	Data collection period	Study population	N	Serum 25(OH)D			
				Mean conc.±SD or (95% CI) [nmol/L]	Prevalence of conc. < 50 nmol/L	Prevalence of conc. > 50 nmol/L	Publ. type
Krajnc, Čokolič et al. 2013 (51)	(year not provided)	Outpatient type II diabetics Age: 59±8 years, mean: 10±8	45	58.0±35.1			B
Kšela and Zavratnik 2013 (52)	Apr 2011	Ambulatory patients with type II diabetes mellitus Age: 65±8.7 years	47	33.8±17.1		2 patients above 75 nmol/L	B
Marc 1993 (18)	(year not provided)	Haemodialysis patients (DP) and continuous peritoneal dialysis patients (CAPD)	85 DP: 51 CAPD: 34	29.2±22.8 DP: 39.5 CAPD: 13.8			C2
Osredkar, Marc 1996 (22)	1994 - 1995	Individuals from general population included on periodic medical examinations	240	Feb: 61.3 Apr: 71.5 Jun: 92.8 Aug: 81.8 Oct: 74.5 Dec: 70.3			A
Pajek, Čuk et al. 2017, Šturm 2017 (24, 25)	2014 - 2015	Dialysis patients (DP) from nine Haemodialysis outpatient centres and controls (K) Age: 56.1±15.2 years	DP: 54 K: 81	68.0±28.4 DP: 73.1±35.4 K: 64.6±22			A, C2
Pajk 2011 (30)	Jan 2009 - Dec 2010	Postmenopausal women from endocrinology and nephrology outpatients centres Age: 55 - 90 years, mean: 72.7	457 55-69 years: 170 70-79 years: 166 80-90 years: 121	48.2±27.4 55-69 years: 51.6±27.0 70-79 years: 48.8±26.8 80-90 years: 42.7±28.3	<25 nmol/L: 25.8% 25-75 nmol/L: 56.7%	25-75 nmol/L: 56.7% >75 nmol/L: 17.5%	C3
Pečovnik-Balon, Jakopin et al. 2009 (48)	Sept 2005 - Sept 2007	Haemodialysis patients treated at the Department of Haemodialysis Age: 60.5±13.1 years	102	58±35.6	10.5-50 nmol/L: 48%	>50 nmol/L: 52%	A
Rakuša, Vrtovec et al. 2020 (57)	2009 - 2018	Heart transplantation recipients Median age: 57 years	123	73.5 (54.7-89.6)	<24.9 nmol/L: 5.1% 25.0-49.9 nmol/L: 16.2%	50.0-74.9 nmol/L: 33.4% >75.0 nmol/L: 45.3%	A
Rondaij, Rotovnik Kozjek et al. 2021 (59)	Jan 2017 - Dec 2018	Chronic intestinal failure patients on long-term home parenteral nutrition Median age: 65 years	63	41.3	<30 nmol/L: 24% 30-50 nmol/L: 48%	50-75 nmol/L: 26%	
Salkić 2016 (56)	Summer: Sept - Nov 2015 Winter: Jan - Mar 2015	Outpatient clinic laboratory Age: 64.5 years	Summer: 776 Winter: 434	Summer: 58.6 Winter: 52.3			C3

References	Data collection period	Study population	N	Serum 25(OH)D			
				Mean conc.±SD or (95% CI) [nmol/L]	Prevalence of conc. < 50 nmol/L	Prevalence of conc. > 50 nmol/L	Publ. type
Soltirovska Salamon, Benedik et al. 2015 (17)	Dec 2010 - Oct 2012	Pregnant Women in 3. trimester	132	74.7±27.5	<50 nmol/L: 14%	50-75 nmol/L: 41%	A
Smaller subsamples of same population (31, 32)		Age: 30.6±4.4 years					
Tesovnik, Kovac et al. 2015 (44)	(year not provided)	Juvenile population with type I diabetes Age: 4-14 years, median: 8.7					
Tomažič, Ul et al. 2007 (58)	Jan - Feb 2006	HIV-infected male population, on three different treatments: T1: antiretroviral treatment naïve; T2: treated with non-protease-inhibitor antiretroviral treatment; T3: treated with protease-inhibitor containing antiretroviral treatment Age: 43 years	96 T1: 24 T2: 37 T3: 35	T1: 35.8 T2: 26.5 T3: 36.5	<50 nmol/L: 82% T1: 75% T2: 89% T3: 82%		A
Trdan 2016 (26)	2014 - 2015	DM and controls age: 67.6 years K age: 55.4 years	818 DM: 266 K: 552	DM: 31.6±17.4 K: 49.2±29.6			C3
Troskot and Duhovnik 2013 (27)	Autumn 2011 - spring 2013	Patients treated with antiepileptic drugs carbamazepine and oxcarbazepine (P) and healthy controls (K) P age: 28-59 years K age: 29-60 years	P: 32 K: 22	P: 55.6±25.2 K: 50.6±20	P: <50 nmol/L: 46.9% K: <50 nmol/L: 59.1%	P: 50-70 nmol/L: 31.3% K: 50-70 nmol/L: 31.8% >75 nmol/L: 21% >75 nmol/L: 9.1%	C4
Vindišar, Goličnik et al. 2009 (53)	(year not provided)	Patients with hip fracture Age: 58- 95 years, mean 79 years	38	39.6±11.9	20-50 nmol/L: 85.8%	>50 nmol/L: 13.2%	B
Vujasinović, Kunst et al. 2016 (104)	(year not provided)	Patients 1 year after bariatric surgery Age: 42.0±9.2 years	22			<65 nmol/L: 95.4%	A
Žmitek, Hribar et al. 2020 (21)	Jan - Mar 2019	Healthy adults, mostly from Central Slovenia Age: 18-65 years; mean 37.7±11.4	238	44.0±17.0 M: 44.3±16.0 F: 43.8±18.0	<30 nmol/L: 21% 30 -50 nmol/L: 45%	50 -75 nmol/L: 31% >75 nmol/L: 3%	A

Notes: Jan - January, Feb - February, Mar - March, Apr - April, Jun - June, Aug - August, Sept - September, Oct - October, Nov - November, Dec - December, K - control group, CD - celiac disease, IR - Institutionalised residents, IBD - inflammatory bowel disease, DP - Dialysis patients, P - Patients treated with antiepileptic drugs carbamazepine and oxcarbazepine, DM - diabetes patients, T1 - antiretroviral treatment naïve; T2 - treated with non-protease-inhibitor antiretroviral treatment; T3 - treated with protease-inhibitor containing antiretroviral CAPD - treatment continuous peritoneal dialysis patients, AG- artistic gymnastic, S - swimming, D1 -strictly compliant to a gluten-free diet, D2 - not strictly compliant to a gluten-free diet, M - male, F - female, Publication type (A - scientific paper, B - conference paper, C - academic thesis; 1 - doctoral, 2 - master, 3 - diploma, 4 - other)

Supplementary table 2. Review of reports for vitamin D intake in the Slovenian population.

Authors	Population	Mean vitamin D intake [$\mu\text{g/day}$]	Data collection	Publication type
Fidler Mis, Kobe et al. 2012 (61)	Teenagers Age 15-16 years M: n=1,010 F: n=1,214	M: 4 \pm 4 F: 4 \pm 3	sq-FFQ (n=2,661) 3DDR for adjustment (n=197)	A
Gregorič 2019 (69, 70)	Elderly in residential home n=49 Age: 65-91 years	2 x 24h-recall: 1.1 \pm 0.6 3DDR: 1.3 \pm 0.7	2 x 24 h-recall 3DDR	A, C3
Gregorič 2015 (62)	Teenagers from 10 primary schools Age 11-15 years n=327 6. grade: age 11-12 years, mean 12.0 \pm 0.1 F: n=79 M: n=78 8. grade 13-15 years, mean 14.0 \pm 0.2 F: n=84 M: n=86	All: 1.7 6. grade F: 1.6 \pm 0.9 M: 1.9 \pm 1.3 8. grade F: 1.5 \pm 1.3 M: 1.9 \pm 1.3	2 x 24h recall	C1
Hribar 2021 (19)	Adolescents Age: 10-17 years M: n=238 F: n=230 Adults Age: 18-64 years M: n=173 F: n=191 Elderly Age: 65-74 years M: n=213 F: n=203	Adolescents All: 2.7 M: 3.0 F: 2.4 Adults All: 2.9 M: 3.4 F: 2.3 Elderly All: 2.5 M: 2.6 F: 2.3	2 x 24h recall adjusted with FPQ	A
Jakše 2020 (71)	Healthy, active adults after dietary intervention study Age: 18-78 years, mean 39.6 years M: n= 42 F: n=109	Food, supplements, and plant-based meal replacements: M: 7.1 \pm 5.8 F: 14.2 \pm 19.4 Food: M: 0.5 \pm 1.1 F: 0.6 \pm 1.0 Supplements, and plant-based meal replacements: M: 6.6 \pm 5.7 F: 13.6 \pm 19.3	3DDR	A
Jeretina 2019 (66)	Healthy peri- and postmenopausal women Age: 40 - 65 years, mean 57.1 \pm 4.7 years n=59	Food: 1.3 \pm 0.93 Food and supplements: 4.1 \pm 10.8	sq-FFQ for fish, milk, and milk products	C2
Juvan 1997 (68)	Students eating vegetarian diets n=14 Semi-vegetarians: 21% Lacto-ovo vegetarians: 58% Lacto vegetarians: 21%	2.6 Semi-vegetarians: 3.3 Lacto-ovo vegetarians: 2.8 Lacto vegetarians: 1.1	7DDR	C3
Koch and Gregorič 2009 (63)	Nationally representative data from secondary school students Age: mean 18 years M: n=216 F: n=237	Consumed with breakfast M: 0.3 \pm 0.5 F: 0.2 \pm 0.4	24h-recall	A

Authors	Population	Mean vitamin D intake [$\mu\text{g/day}$]	Data collection	Publication type
Kocuvan Mijatov and Mičetić-Turk 2016 (72)	Adult female coeliac disease patients Age: 23 - 76 years n=40	2.6	3DDR	A
Lichthammer, Nagy et al. 2015 (65)	Teenage and adult population Age: 15-75 years, mean 38.8 n=81	All: 2.57 ----- 14-18 years: 3.7 \pm 3.3 19-29 years: 3.0 \pm 2.1 30-69 years: 2.5 \pm 1.7 >70 years: 1.0 \pm 2.4 FFQ: 1.8 24h-recall: 1.4	sq-FFQ	A
Lopert 2019 (73)	Female patients with osteoporosis Age: mean 71.4 years n=43	FFQ: 1.8 24h-recall: 1.4	FFQ, 24h-recall	C2
Poličnik, Pokorn et al. 2013 (60)	Pre-school children in Central Slovenia Age: 2-6 years, average 4.2 n=129 M: n=68 F: n=61	All: 1.1 \pm 0.7 M: 1.0 \pm 0.7 F: 1.1 \pm 0.7	3DDR	A
Soltirovska Salamon, Benedik et al. 2015 (17)	Pregnant women in third trimester Age: 30.6 \pm 4.4 years n=132		4DDR	A
Urh, Babnik et al. 2017 (67)	Elderly in a residential home Age: mean 82.2 years n=48	2.6	7DDR	A
Zdešar-Kotnik 2019 (64)	Pupils of 1. years of Secondary school M: Age: 13.5 years; n=162 F: Age: 15.4 years; n=180	M: 1.4 (0.8 - 2.5) F: 1.0 (0.6 - 1.7)	2x24h-recall	C1

Notes: M - males, F - females, DDR - day dietary record, FFQ - food frequency questionnaire, FPQ - food propensity questionnaire, sq - semi-quantitative; Publication type (A - scientific paper, B - conference paper, C - academic thesis; 1 - doctoral, 2 - master, 3 - diploma, 4 - other)