



Review

The use of continuous glucose monitoring in people living with obesity, intermediate hyperglycemia or type 2 diabetes

Tadej Battelino^{a,*}, Nebojsa Lalic^b, Sufyan Hussain^{c,d,e}, Antonio Ceriello^f, Sanja Klobucar^g, Sarah J. Davies^h, Pinar Topseverⁱ, Julie Heverly^j, Francesca Ulivi^k, Kevin Brady^l, Tsvetalana Tankova^m, Júlia Galhardoⁿ, Kostas Tagkalos^o, Erik Werson^p, Chantal Mathieu^q, Peter Schwarz^{r,s}

^a University Medical Center Ljubljana, and University of Ljubljana, Faculty of Medicine, Ljubljana, Slovenia

^b Faculty of Medicine, University of Belgrade, Center for Diabetes and Lipid Disorders, Clinic for Endocrinology, Diabetes and Metabolic Diseases, University Clinical Centre of Serbia, Belgrade, Serbia

^c Department of Diabetes, School of Cardiovascular, Metabolic Medicine and Sciences, King's College London, London, UK

^d Department of Diabetes and Endocrinology, Guy's & St Thomas' NHS Foundation Trust, London, UK

^e Institute of Diabetes, Endocrinology and Obesity, King's Health Partners, London, UK

^f IRCCS MultiMedica, Milan, Italy

^g Department for Endocrinology, Diabetes and Metabolism, University Hospital Centre Rijeka, School of Medicine, University of Rijeka, Croatia

^h Woodlands Medical Centre, Cardiff, Wales, UK

ⁱ Acibadem Mehmet Ali Aydinlar University, School of Medicine, Department of Family Medicine, Istanbul, Türkiye

^j diaTribe Foundation, San Francisco, CA, United States

^k Fondazione Italiana Diabete, Legnano, Milan, Italy

^l diabetes Geneva, Avenue Cardinal-Mermillod 36, Carouge, Switzerland

^m Department of Endocrinology, Medical University - Sofia, Sofia, Bulgaria

ⁿ Dona Estefânia Hospital - ULSS José, Lisbon, Portugal

^o IDF Europe, the Netherlands

^p Beekbergen, the Netherlands

^q Department of Endocrinology, University Hospitals Leuven, Louvain, Belgium

^r Department for Prevention and Care of Diabetes, Faculty of Medicine Carl Gustav Carus at the Technische Universität/TU Dresden, Dresden, Germany

^s Paul Langerhans Institute Dresden of Helmholtz Zentrum München at University Hospital and Faculty of Medicine, TU Dresden 01307 Dresden, Germany

ARTICLE INFO

Keywords:

Prediabetes

Intermediate hyperglycemia

Obesity

Type 2 diabetes

Continuous glucose monitoring

Consensus statements

ABSTRACT

A global trend towards increased obesity, intermediate hyperglycemia (previously termed prediabetes) and type 2 diabetes, has prompted a range of international initiatives to proactively raise awareness and provide action-driven recommendations to prevent and manage these linked disease states. One approach, that has shown success in managing people already diagnosed with type 2 diabetes mellitus, is to use continuous glucose monitoring (CGM) devices to help them manage their chronic condition through understanding and treating their daily glucose fluctuations, in association with glucose-lowering medications, including insulin. However, much of the burden of type 2 diabetes mellitus is founded in the delayed detection both of type 2 diabetes mellitus itself, and the intermediate hyperglycemia that precedes it. In this review, we provide evidence that using CGM technology in people at-risk of intermediate hyperglycemia or type 2 diabetes mellitus can significantly improve the rate and timing of detection of dysglycemia. Earlier detection allows intervention, including through continued use of CGM to guide changes to diet and lifestyle, that can delay or prevent harmful progression of early dysglycemia. Although further research is needed to fully understand the cost-effectiveness of this intervention in people at-risk or with early dysglycemia, the proposition for use of CGM technology is clear.

* Corresponding author at: University Medical Center Ljubljana, Bohoriceva 20, 1000 Ljubljana, Slovenia.

E-mail address: tadej.battelino@mf.uni-lj.si (T. Battelino).

<https://doi.org/10.1016/j.diabres.2025.112111>

Received 17 February 2025; Received in revised form 12 March 2025; Accepted 17 March 2025

Available online 19 March 2025

0168-8227/© 2025 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Obesity, particularly abdominal obesity, is a significant risk factor for the development of insulin resistance, which is a precursor to both intermediate hyperglycemia and type 2 diabetes [1,2]. The terms prediabetes, non-diabetic hyperglycemia and intermediate hyperglycemia have been used interchangeably to indicate a state of increased risk of progressing to type 2 diabetes mellitus. For consistency, we will refer to only intermediate hyperglycemia throughout the rest of this discussion. Excess fat, especially in the abdominal area, leads to an increase in fatty acids and inflammation, which can impair the body's ability to use insulin effectively. Insulin resistance as a consequence of obesity is a common feature of intermediate hyperglycemia, with elevated blood glucose as the body struggles to use insulin effectively, eventually progressing in many individuals to type 2 diabetes mellitus if not managed. A series of major studies in the USA, Finland, and China have shown that interventions to increase dietary weight loss and improve physical activity, can prevent progression to type 2 diabetes mellitus in high-risk persons by 51–58 %, for at least 6 years following the original intervention [3–5], with continued protection of 27–43 % for up to 20 years [5–7]. The efficacy of successful intervention has been attributed to a multiplicity of factors, including: lifestyle coaches; contact with at-risk persons; education, and; networks for feedback and support [8]. The opportunity provided by proactive use of CGM systems to provide aspects of this support at lower resource intensity, for example by using remote versus in-person contacts with healthcare professionals (HCPs) must be considered here.

An important goal in the chain of metabolic dysregulation is to limit the risks associated with type 2 diabetes mellitus as early as possible, through interventions to promote weight loss and mitigate the adverse effects of hyperglycemia. A recent study has shown that modification of lifestyle behavior is important to prevent damage to kidney function among the overweight prediabetic population in men [9]. The attributes of continuous glucose monitoring (CGM) technologies, which are proven to improve glycemic control for adults with type 1 diabetes (T1D) [10–14], are increasingly proposed as an intervention for managing obesity, by enabling behavioral change that drives weight loss, and for glycemic control in intermediate hyperglycemia and type 2 diabetes mellitus. A small number of proof-of-concept studies have shown that adults with type 2 diabetes mellitus using CGM do choose lower glycemic index foods [15], increased physical activity [16], decreased caloric intake, weight loss, and reduced postprandial glucose levels [17]. Among people with intermediate hyperglycemia, there is only one published study addressing the role of CGM in promoting behavior change, and, while it showed greater dietary self-efficacy, neither weight nor glycemic measures were reported [18].

Ultimately, there is a significant unmet need to emphasize with at-risk persons and those with intermediate hyperglycemia that there are a significant number of modifiable factors within their own capabilities to change – and get them to act on them, while bearing in mind other contributing factors such as social deprivation. Through empowerment of these groups of people, the burden of type 2 diabetes mellitus can be restricted or delayed. Lifestyle and dietary interventions managed in primary care can restore normoglycemia for people with intermediate hyperglycemia [19]. Significantly, despite concerns about the negative connotations of being 'labelled' with intermediate hyperglycemia, at least one study has indicated that intermediate hyperglycemia was not considered as negative but an opportunity to engage with primary-care supported dietary interventions [20]. Equally, although people with lower social determinants of health (SDoH) are known to have reduced access to healthcare services, education has been shown to support improved glycemic performance and encourage healthy lifestyle changes in people with intermediate hyperglycemia from communities with lower socioeconomic status [21]. In each of these situations, there is a need to emphasize that the application of CGM can be part of continuous glucose education.

As CGM devices are becoming available without medical prescription in the USA, increasing random use is anticipated with likely additional burden to the primary care level physicians and teams, with no available guidance on how to instruct this populations.

2. The evidence for using CGM in normoglycemic persons living with obesity

The rationale for using CGM in obese persons without intermediate hyperglycemia or type 2 diabetes mellitus is to support lifestyle changes to diet and physical activity, by providing real-time biofeedback on postprandial glucose spikes and glycemic variability (GV), that have been proposed to drive hyperinsulinemia and increased fat storage [22,23]. However, research in this aspect of obesity management is a significant unmet need. One study using CGM in normoglycemic adults has shown that glucometric data measuring glycemic variability was not different between obese (BMI > 30 kg/m²) and non-obese persons [24]. A separate study on normoglycemic but severely (Class 3) obese people (BMI ≥ 40 kg/m²) did find significantly increased metrics of GV, compared to normoglycemic non-obese persons [25], similar to severely obese people diagnosed as having intermediate hyperglycemia. This suggests that the goal of influencing behavior using biofeedback from CGM devices may be more realistic in severely obese people, but no studies have examined this proposition.

2.1. Targeting and modifying behavior using CGM in persons living with obesity

Studies using CGM have confirmed that overweight and obesity in men without diabetes and with a 2-hour plasma glucose during oral glucose tolerance test (OGTT) < 200 mg/dL (11.1 mmol/L), is associated with eating and snacking behaviors that increase CGM-detected indices of hyperglycemia, including maximal glucose, time above range (TAR) between 140 mg/dL (7.8 mmol/L), and 200 mg/dL (11.1 mmol/L), as well as the frequency of postprandial hyperglycemia [26,27]. A low physical activity profile has also contributed to impaired glycemic control in this group. Notably, these associations were found to be independent of measured pancreatic β -cell function.

In this context, a qualitative study has concluded that successful weight-loss interventions for adults living with obesity but without a diagnosis of diabetes are dependent on regular monitoring and feedback to encourage ownership, accountability and self-efficacy, to develop behaviors adapted to long-term adherence with weight-loss goals [28]. Periodic use of CGM can be argued to support these aspects of behavior change, providing daily feedback to the user on healthy versus unhealthy activities, along with objective data on glycemic patterns that can be reviewed with their HCP. A small-scale pilot study using CGM to support people with intermediate hyperglycemia or type 2 diabetes mellitus has shown that using CGM to provide biofeedback was effective at promoting attendance at planned exercise sessions and registration for future exercise activities, compared to participants not using CGM [29]. Looking at dietary interventions, CGM has been used to monitor and support adult males (n = 15, mean age 55 yrs) taking part in a study on time-restricted eating (TRE) over one week [30]. The outcomes of this study showed a limited benefit on mean fasting glucose during TRE phase. Use of CGM in adolescents living with obesity has also provided some evidence that adherence with TRE strategies may be supported when participants wear CGM to monitor adherence with TRE activities [31]. Overall, the available studies support the proposition that, rather than simply being tools that help users to 'react' to hypoglycemia or hyperglycemia, CGM devices can help people with prediabetes or diabetes to proactively make preventive lifestyle behavior changes [32]. However, larger-scale studies are necessary to fully-explore these

associations.

Consensus statements. People living with obesity and at-risk of progression to type 2 diabetes mellitus.

- The goal of addressing modifiable risk factors in people living with obesity is realistic, such that progression to intermediate hyperglycemia and type 2 diabetes mellitus may be reduced.
- Benefits at a global population level will only be realised if access to CGM technology is improved to enable widespread adoption, especially in developing countries where burden and impact from diabetes remains highest.
- Periodic use of CGM should be considered in people living with obesity and at-risk of developing type 2 diabetes mellitus based on established risk factors [33], to assess glycemic risks and support early intervention.
- Measures of CGM-defined glycemic risk amongst people living with obesity centered on time above tight range (TATR) > 140 mg/dL (7.8 mmol/L) and glycemic variability may be used to identify persons at-risk of intermediate hyperglycemia.
- The application of CGM should be considered during proactive efforts to modify diet and physical activity in people living with obesity, in order to provide immediate feedback on the glycemic impact of dietary choices, as well as the glycemic benefits of physical activity and exercise.
- Any at-risk individual living with obesity and with normoglycemia, as assessed by FPG, plasma glucose during OGTT or standard HbA1c testing, should have periodic use of CGM to assess glycemic control patterns under free-living conditions.
- There is an unmet need for prospective studies to understand the value of CGM use in people living with obesity and at-risk of type 2 diabetes mellitus. This should include studies to investigate the real-time glycemic patterns that characterize these at-risk persons and also investigate the value of CGM to support dietary and lifestyle interventions.
- Further developments and integration of CGM data and providing artificial intelligence (AI) enabling Clinical Decision Support Systems (CDSS) is required to simplify interpretation for those with lower digital and health literacy.

3. The evidence for using CGM in intermediate hyperglycemia

Intermediate hyperglycemia is defined by either: HbA1c 5.7–6.4 % (39–47 mmol/mol), impaired glucose tolerance (IGT) with fasting plasma glucose (FPG) 100–125 mg/dL (5.6–6.9 mmol/L) or IGT with 2-hour plasma glucose 140–199 mg/dL (7.8–11.0 mmol/L) or a 1-hour plasma glucose $\geq$ 155 mg/dL (8.6 mmol/L) following a 75 g OGTT [34,35]. Each of these criteria provides a single measurement at a single timepoint, and are impractical to repeat serially. In contrast, the use of a single CGM sensor for a 10–15-day period can provide a wealth of glycemic data under real-world conditions, including at home and work, that can provide a definitive diagnosis of intermediate hyperglycemia, along with additional insights into day-to-day glycemia and how it is affected by meal planning and lifestyle. All of this can be managed with remote monitoring, such that a clinic visit may not be required in many cases.

People living with intermediate hyperglycemia have an 83 % increased risk of CVD [36], although non-glycemic health status indicators also contribute to this risk. A meta-analysis of 71 studies assessing the impact of GV on cardiometabolic risk factors in people without diabetes found that elevated GV in intermediate hyperglycemia is associated with risk of developing coronary atherosclerosis, independently of obesity or traditional risk factors [37]. Intermediate hyperglycemia is also associated with increased incidence of microvascular complications [38–42]. In progression to type 2 diabetes mellitus, the prediabetic period has been separated into stage 1 (early) and stage 2 (late) [43,44], and CGM can be an effective tool to identify the presence and severity of dysglycemia for persons living with intermediate hyperglycemia [44]. For example, GV [45,46] and time above tight range (TATR) > 140 mg/dL (7.8 mmol/L) [47] can be used separately or together to distinguish between people with or without type 2 diabetes mellitus. Using CGM in people without a diagnosis of type 2 diabetes mellitus indicates that TATR > 140 mg/dL (7.8 mmol/L) is uncommon [48,49], suggesting that %TATR > 140 mg/dL (7.8 mmol/L) may be clinically meaningful in assessing the risk of developing type 2 diabetes mellitus amongst people living with intermediate hyperglycemia [50]. This is further supported by a retrospective comparative analysis of CGM

data from 836 adults across 5 separate studies, including normoglycemic persons, those with intermediate hyperglycemia, people with type 2 diabetes mellitus and HbA1c < 6.5 % (<48 mmol/mol), <7.0 % (<53 mmol/mol) or $\geq$ 7.0 % ($\geq$ 53 mmol/mol) [51]. The outcomes showed a progressive deterioration in GV and %TATR > 140 mg/dL (7.8 mmol/L) from normoglycemia to dysglycemia, through to type 2 diabetes mellitus with HbA1c < 6.5 % (<48 mmol/mol) and thereafter to type 2 diabetes mellitus with HbA1c $\geq$ 7.0 % ($\geq$ 53 mmol/mol). Established CGM metrics also show a progression of dysglycemia from at-risk persons to those with intermediate hyperglycemia and those with type 2 diabetes mellitus [52], who demonstrated increased TATR 140–180 mg/dL (7.8 – 10.0 mmol/L), during the day and overnight. Stratifying CGM readings by time of day and the range 140–180 mg/dL (7.8 – 10.0 mmol/L), provides additional metrics to differentiate between the groups.

In people considered to be nondiabetic using the established criteria (i.e., HbA1c < 6.5 % [<48 mmol/mol] or 2-hour 75 g OGTT > 200 mg/dL), application of CGM has identified an additional 15 % of people with intermediate hyperglycemia [53], with significantly increased metrics of GV, which can differentiate between people with normal glucose tolerance and those with intermediate hyperglycemia, who are matched for HbA1c [54–56], indicating that CGM metrics of GV may be a more-sensitive indicator of intermediate hyperglycemia. The use of CGM metrics together with isolated abnormalities in OGTT testing has also been proposed as a way to identify persons at high-risk of intermediate hyperglycemia or type 2 diabetes mellitus, despite HbA1c < 5.7 % (<39 mmol/mol) [57]. It seems clear that, compared to HbA1c or OGTT testing, the key benefit of CGM is the stratification of intermediate hyperglycemia according to GV metrics, as this may provide stronger information of β -cell function and insulin sensitivity [58], and potentially to identify glucotypes [53,59] that may guide better personalized prevention strategies earlier in the disease process. Recently, the International Diabetes Federation has issued a position statement supporting the value of 1-hour glycemia $\geq$ 155 mg/dL (8.6 mmol/L) during a 75 g OGTT as a stronger tool for the early diagnosis of intermediate hyperglycemia, with a 1-hour plasma glucose of $\geq$ 209 mg/dL (11.6 mmol/L) indicating type 2 diabetes mellitus [35]. In this context, it is also important to note that 1-hour glycemia has been associated with a high risk for both macro- and microvascular complications [60]. Studies in this setting, involving CGM, are not yet available.

Although CGM-derived metrics of remission of intermediate hyperglycemia to normoglycemia have not been investigated formally, the evidence, as discussed above, suggests that optimizing T1TR 70–140 mg/dL (3.9–7.8 mmol/L) and reducing TATR > 140 mg/dL (>7.8 mmol/L) should be targeted. Although studies vary, individuals with intermediate hyperglycemia may have < 77 % T1TR 70–140 mg/dL (3.9–7.8 mmol/L), compared to > 87 % for normoglycemic individuals, with up to 17 % TATR > 140 mg/dL (>7.8 mmol/L) compared to 12 % [61]. Given the increased risks for macrovascular and microvascular disease for people with intermediate hyperglycemia, this is an important goal. Studies have shown that T1TR 70–140 mg/dL (3.9–7.8 mmol/mol) is inversely correlated with the occurrence of cerebrovascular accidents (CVAs) after adjustment for HbA1c [84]. Regression analysis concluded that for every 10 % increase in T1TR 70–140 mg/dL (3.9–7.8 mmol/mol) the risk of a CVA event was reduced by 45 % ($p < 0.01$). Similarly, each 10 % increase in T1TR 70–140 mg/dL (3.9–7.8 mmol/mol) has been associated with a 12 % rate reduction in the incidence of peripheral neuropathy [45] and a 56 % lower rate of diabetic retinopathy [72]. These data refer to study cohorts with a diagnosis of diabetes, and additional studies are required specifically in individuals with intermediate hyperglycemia.

With the availability of numerous CGM-derived metrics that allow detection and monitoring of intermediate hyperglycemia, it has been proposed that this is an opportunity to avoid therapeutic inertia in the progression to confirmed type 2 diabetes mellitus [62], since these glycemic changes will indicate ongoing cellular dysfunction that can generate insulin deficiency, insulin resistance or both. Given that changes in meal-planning and physical activity are more-effective than

pharmacotherapy in preventing progression from intermediate hyperglycemia to type 2 diabetes mellitus in adults [3], the use of CGM as a motivational tool for making lifestyle changes is of considerable importance. Only limited data is available from a small number of studies, but the data does support the proposition that using CGM can lead to improved PA, dietary choices and weight loss in people with intermediate hyperglycemia. Daily consumption of carbohydrate and refined grains have been found to be associated with higher GV, whereas whole grains and daily protein intake have been related to lower GV in people with IGT. These results can offer further insights into designing a more efficacious dietary intervention in people with intermediate hyperglycemia [63]. One small pilot study (n = 13) on exercise, including participants with intermediate hyperglycemia and diabetes, reported positive changes in body composition and increased fitness when using CGM, as well as improved goal setting and self-monitoring behavior [29]. A larger study (n = 168) in adults living with intermediate hyperglycemia [64] showed that a 12-week intervention with a smartphone app and CGM increased T1TR 70–140 mg/dL (3.9–7.8 mmol/L), accompanied by reduced carbohydrate intake and weight loss. A single-arm prospective study of CGM use in 32 people with intermediate hyperglycemia [65], reported satisfaction with CGM use was largely positive, and the majority agreed (60 %) or strongly agreed (22 %) that wearing a CGM sensor reminded them to stay healthy every day. Another study assessed the satisfaction and feasibility of using CGM in 15 adults with intermediate hyperglycemia in combination with a low-carbohydrate diet [66]. In addition to a high rate of satisfaction (93 %) among participants, investigators observed significant reductions in HbA1c and body weight.

3.1. Remission of type 2 diabetes mellitus to intermediate hyperglycemia

It is widely accepted that type 2 diabetes mellitus is a chronic disease, which can be controlled through lifestyle adaptations and drug treatment or metabolic surgery, but which is progressive and inevitable. Consequently, remission of type 2 diabetes mellitus is rarely addressed as a treatment goal. However, remission of type 2 diabetes mellitus to glucose levels in the range of intermediate hyperglycemia, or even normoglycemia, is now considered an achievable goal for certain persons [67,68]. A 2022 consensus has proposed that, from a glycemic standpoint, remission in type 2 diabetes mellitus can be defined as a return to an HbA1c < 6.5 % (48 mmol/mol), measured at least 3 months following the cessation of active glucose-lowering pharmacotherapy [69]. Although FPG < 126 mg/dL (7.0 mmol/L) in the absence of pharmacotherapy can be considered in some settings as an alternative criterion for remission of type 2 diabetes mellitus [70], the need for overnight fasting and the significant variation between repeated FPG measurements is a critical disadvantage. HbA1c itself is affected by a number of non-glycemic factors and repeated HbA1c testing is meaningful only every 3 months, which limits its value for the timely detection of a recurrence of hyperglycemia. Thus, use of CGM to monitor maintained euglycemia in remission of type 2 diabetes mellitus is of interest.

Remission of type 2 diabetes mellitus has become a realistic goal largely as a result of research showing the metabolic benefits of sustained weight loss, typically as a result of dietary restrictions, when insulin secretion and insulin sensitivity are not significantly impaired. The primary-care DiRECT trial in the UK showed remission of type 2 diabetes mellitus in 46 % of participants after 1 year following an intensive dietary replacement intervention [71]. After 2 years, and 12 months after the end of the intervention, remission was still evident in 36 % of participants, with loss of remission being associated with weight gain [67]. Notably, restoration of β -cell function was also evident, associated with reduced ectopic fat in the liver and pancreas. Similar outcomes using the DiRECT protocol have been observed in a primary-care study in Australia [72]. These outcomes were confirmed in the DIADEM-I RCT [73], showing remission of type 2 diabetes mellitus in

60 % of participants after a 12-month dietary intervention, including a return to normoglycemia in 30 % of participants. The long-term follow-up of the DiRECT trial also indicates that the metabolic signature associated with development of type 2 diabetes mellitus can be modified and reversed by weight-loss interventions [74].

The Look AHEAD study, which used a lifestyle intervention based on frequent counselling to reduce caloric intake and increase physical activity, reported a type 2 diabetes mellitus remission rate of 11.5 % among participants at 12 months, falling to 7 % at 2 years [75]. However, an important outcome for participants who showed evidence of remission of type 2 diabetes mellitus was a 33 % reduced risk of chronic kidney disease (CKD) and a 40 % reduced risk of CVD after 8 years of follow-up, compared to participants who did not exhibit remission [76].

Pharmacotherapy to support intensive weight-loss programs in obesity and type 2 diabetes mellitus has been proposed as a way to reduce the hunger associated with caloric restriction and to increase adherence with low-energy diets [77]. This has been evaluated in pilot studies on dulaglutide [78], a once-weekly glucagon-like peptide-1 receptor agonist (GLP-1 RA), which has an appetite-suppressant action [79,80]. A 12-week intervention showed that use of dulaglutide alongside a low-calorie diet promoted a mean 9.5 kg loss of weight and a reduction in HbA1c of -1.4 % (-15.8 mmol/mol). Although these reductions in weight and HbA1c are significant over a 3 month period, no comparator arm was included in this pilot investigation.

T1R does not appear to be helpful for assessing diabetes remission to the intermediate hyperglycemic state, in contrast to T1TR which is a more beneficial CGM metric for confirming normoglycemia, as has been recently suggested [81]. Discussion of CGM metrics of remission in cases of previously diagnosed type 2 diabetes mellitus are largely focused on glycemic changes following either dietary intervention or bariatric surgery in people with type 2 diabetes mellitus [82–85]. Although the 14–18 % T1TR > 140 mg/dL (7.8 mmol/L) values identified in these remission studies [82,84,85] are aligned with the investigations on intermediate hyperglycemia described earlier in this paper, the achievement of only 70 % T1TR is not consistent with restored normoglycemia [84,85]. The limited evidence does imply that a target of < 20 % T1TR > 140 mg/dL (7.8 mmol/L), for at least a 12-month duration in the absence of pharmacological therapy, may support maintenance of remission from overt type 2 diabetes mellitus but still indicate intermediate hyperglycemia. Additional studies are needed to better understand how CGM may be used in this group of people with dysglycemia. It is also important to identify a wider selection of CGM measures of glycemia in people with type 2 diabetes mellitus during periods of confirmed remission to intermediate hyperglycemia, with sustained HbA1c 6.5 % (<48 mmol/mol), including the correlation between the CGM-derived glucose management indicator (GMI), derived from mean glucose levels, and HbA1c during remission.

The potential value and suitability of using CGM systems, both to assess and manage intermediate hyperglycemia, is clearly indicated by these few studies, and underscores the need for further prospective studies on the application of CGM in persons living with intermediate hyperglycemia.

Consensus statements. People living with intermediate hyperglycemia and at-risk of progression to type 2 diabetes mellitus.

- The goal of reversing dysglycemia in people living with intermediate hyperglycemia is realistic, as is delaying progression to type 2 diabetes mellitus.
- Benefits at a global population level will only be realised if access to CGM technology is improved to enable widespread adoption, especially in developing countries where burden and impact from diabetes remains highest.
- Periodic use of CGM should be considered in people living with intermediate hyperglycemia, based on established risk factors for type 2 diabetes mellitus [33], to assess glycemic risks and support early intervention when needed.
- For persons with prior intermediate hyperglycemia and a recent diagnosis of type 2 diabetes mellitus and able to meet CGM targets for > 70 % T1TR, setting targets for T1TR can be considered to aim for near-normal glycemia or remission.
- Remission criteria for people being treated for type 2 diabetes mellitus should mirror diagnostic criteria, and include: FPG < 126 mg/dL (7.0 mmol/L), mean

(continued on next page)

(continued)

Consensus statements. People living with intermediate hyperglycemia and at-risk of progression to type 2 diabetes mellitus.

- glucose levels < 100 mg/dL (5.6 mmol/L), GMI < 6.5 % (<48 mmol/mol), sustained over 3 months in the absence of pharmacotherapy.
- Use of CGM should be considered as part of proactive efforts to modify diet and physical activity in people living with intermediate hyperglycemia, in order to provide immediate feedback on the glycemic impact of dietary choices and physical activity or exercise.
- There is an unmet need for prospective studies to understand the value of CGM use in people living with intermediate hyperglycemia and at-risk of progression to type 2 diabetes mellitus. This should include studies to investigate the real-time glycemic patterns that characterize intermediate hyperglycemia and to investigate the value of CGM to support dietary and lifestyle interventions in intermediate hyperglycemia.
- Further developments and integration of CGM data is required to simplify interpretation for those with lower digital and health literacy.

4. The evidence for using CGM in people with type 2 diabetes mellitus

Compared to the very limited data on the use of CGM in at-risk, normoglycemic obese adults and those with a dysglycemic profile characteristic of intermediate hyperglycemia, the data on outcomes following initiation of CGM in people with diagnosed type 2 diabetes mellitus are substantial, including those on non-insulin therapies. Besides the glycemic benefits summarized below, there are several studies that have demonstrated the benefits of CGM for supporting behavioral change, including weight loss, lower carbohydrate intake and improved physical activity [15–17,86–89]. One of the more important goals of using CGM in type 2 diabetes mellitus is to combat therapeutic inertia, in which escalation of drug therapy is delayed well beyond recommended guidelines. Although therapeutic inertia is a consequence of several intersecting factors [90–93], fear of hypoglycemia is a key problem [94,95], because of its association with adverse outcomes [96,97]. Use of CGM can make this a modifiable factor and several studies have shown that using CGM in T1D is associated with reduced fear of hypoglycemia [98,99]. More significantly, retrospective analysis of large healthcare claims datasets indicate that application of CGM in type 2 diabetes mellitus is associated with more-timely treatment intensification, compared with self-monitored blood glucose (SMBG) testing [100].

4.1. Persons with type 2 diabetes mellitus on intensive insulin therapy

There is considerable evidence that people with type 2 diabetes mellitus on intensive insulin therapy (either multiple daily injections [MDI] with insulin or continuous subcutaneous insulin infusion [CSII]) can benefit from CGM devices in the same way as has been demonstrated for people with T1D. This includes lower HbA1c [101,102] and reduced hypoglycemia [102,103], as well as fewer hospital admissions for acute diabetes events (ADEs), such as diabetic ketoacidosis (DKA) or severe hypoglycemia [104–106], or for long-term microvascular and macrovascular complications [107]. Consequently, guidelines recommend CGM in type 2 diabetes mellitus on intensive insulin therapy [108,109], given the clinical benefits and cost effectiveness [110,111].

4.2. Persons with type 2 diabetes mellitus on basal insulin therapy

Initiation and titration of basal insulin therapy can be associated with episodes of problematic hypoglycemia, particularly in older people [112], and application of CGM can help to reduce such episodes following the start of basal insulin, by allowing the persons with type 2 diabetes mellitus to see their glucose levels in real time, as well as whether they are falling and how fast, using the trend arrows. Similarly, persistent TAR can be recognized using CGM and addressed by adjusting basal insulin doses upwards.

The MOBILE RCT has shown that use of CGM can significantly reduce

HbA1c, time in hyperglycemia and the rate of hypoglycemia events over an 8-month period, compared to a control group using SMBG testing alone [113]. These data are consistent with results of other randomized controlled trials (RCTs) [114,115] and retrospective studies [116,117] demonstrating significant reduction in HbA1c for people with type 2 diabetes mellitus on basal insulin therapy. Equally, use of CGM in type 2 diabetes mellitus treated with basal insulin is associated with reduced hospital admissions, both for ADEs [118] and for cardiovascular complications [107]. Overall, use of CGM in type 2 diabetes mellitus treated with basal insulin has been shown to be cost effective, compared to routine SMBG testing [119].

4.3. Persons with type 2 diabetes mellitus on non-insulin therapies

A number of RCTs and real-world studies have shown that the use of CGM in non-insulin treated type 2 diabetes mellitus can significantly reduce HbA1c [120–124], particularly for those with higher HbA1c levels [125]. Glycemic variability, which is associated with adverse clinical outcomes [126–129], is also reduced in people with type 2 diabetes mellitus on non-insulin therapy using CGM [121]. Similarly, use of CGM in type 2 diabetes mellitus on non-insulin therapy is associated with reduced ADEs requiring hospital attendance or admission [125]. Notably, CGM has been shown to act as a motivational tool for helping people with inadequately controlled type 2 diabetes mellitus on non-insulin therapy to establish and adhere to lifestyle changes, and thereby reduce glycemia [130,131].

The use of CGM on a periodic basis can be a viable option at regular intervals or during treatment intensification or deintensification, particularly if oral insulin-secretagogue drugs (such as sulphonylureas) have been prescribed and where a risk of hypoglycemia is evident [132]. Blinded CGM has revealed that approximately 50 % of people with type 2 diabetes mellitus, including those on non-insulin therapy, experience frequent mild or clinically significant hypoglycemia [133], which is typically asymptomatic.

Together, these studies show that intermittent use of CGM systems provides glycemic information of value both to the person with type 2 diabetes mellitus and to their healthcare professional, which can facilitate improved glycemic control through changes to lifestyle and periodic medication adjustments. An important goal for this group of people with type 2 diabetes mellitus is to delay progression to insulin therapy. Since obesity is a major factor in metabolic decompensation leading to insulin treatment, it can be proposed that CGM can be used to monitor hyperglycemia for people with type 2 diabetes mellitus on weight-loss medications [134,135], during this period.

4.4. The newly diagnosed person with type 2 diabetes mellitus

The heterogeneity of disease for people diagnosed with type 2 diabetes mellitus is significant and can be mapped to a number of glucometric profiles, including for people newly diagnosed with type 2 diabetes mellitus [53,59,136]. The UKPDS study population consisted of 5,102 people with newly diagnosed type 2 diabetes mellitus, and demonstrated the importance of early and proactive glucose control for reducing long-term diabetes complications [137–139]. Therefore, CGM can be an effective option during the period following diagnosis of type 2 diabetes mellitus, to establish baseline glycemic profiles for each individual, against which subsequent treatment decisions may be compared and disease progression monitored.

4.5. Targeting and modifying behavior using CGM in people living with type 2 diabetes mellitus

The goal of addressing unhealthy dietary and lifestyle factors that may contribute to a diagnosis of type 2 diabetes mellitus may be productively addressed in people newly diagnosed, when motivation may be high to make positive changes to self-management behaviors related

to diet and exercise. Dietary interventions have been shown to be successful in adults with newly diagnosed type 2 diabetes mellitus, as measured by blinded CGM that is not used as a motivational tool [140]. A 2022 pilot study [82] specifically used CGM as a part of a 3-month intervention, focused on minimizing postprandial excursions in 17 adults with recently-diagnosed type 2 diabetes mellitus and mean HbA1c 8.0 % (64 mmol/mol), through educated use of CGM to understand the glycemic impact of food choices and exercise on PPE. After 3 months, 67 % of the participants exhibited remission of type 2 diabetes mellitus, with HbA1c < 6.5 % (<48 mmol/mol), TATR > 140 mg/dL (7.8 mmol/L) was significantly reduced, and consumption of carbohydrates and BMI were significantly lower. Although the study conclusions are limited by the small study size, similar outcomes for remission of type 2 diabetes mellitus were achieved as in the 12-month DiRECT study (see above), but over a 3-month period and without whole-scale dietary replacement. The CGM-directed intervention also involved regular motivational text messaging, thus leveraging multiple technologies, but participants reported that CGM use was the most significant element of the intervention. This pilot study did not have a control arm but the outcomes were improved compared to similar studies using education to minimize PPE and using standard SMBG to monitor glucose fluctuations [141,142]. Use of CGM actively to support behavior change through education aimed at reducing PPE has also shown significant reductions in HbA1c in adults with a type 2 diabetes mellitus duration of up to 11 years, compared to a control group using SMBG [123]. Significantly reduced glycemia > 180 mg/dL (>10.0 mmol/L), lowered mean HbA1c (−1.0 % [−11 mmol/mol]) and weight loss (≥4 lbs) has also been demonstrated among 72 people with type 2 diabetes mellitus not on insulin using CGM, with or without a food-logging app, over 90 days [143].

Consensus statements. People living with type 2 diabetes mellitus.

- The goal of reversing dysglycemia in people newly diagnosed with type 2 diabetes mellitus is achievable in many cases.
- Benefits at a global population level will only be realised if access to CGM technology is improved to enable widespread adoption, especially in developing countries where burden and impact from diabetes remains highest.
- For people with newly diagnosed type 2 diabetes mellitus, CGM should be applied to establish their baseline glycemic profile and deciding and monitoring their initial treatment approach. Where restoration of normoglycemia is assessed as achievable, use of CGM should be used as part of this approach.
- For people with type 2 diabetes mellitus on non-insulin therapy, CGM use should be considered as an educational and motivational tool during diabetes self-management education and support (DSMES) interventions targeted at diet, physical activity and exercise.
- CGM should be applied as standard of care in people living with type 2 diabetes mellitus on non-insulin therapies, at any point where treatment intensification or deintensification is required.
- For people with type 2 diabetes mellitus on insulin therapy, we endorse existing international recommendations on the access and use of CGM technology [144]
- Further developments and integration of CGM data is required to simplify interpretation for those with lower digital and health literacy.

5. The value of CGM for predicting long term complications of type 2 diabetes mellitus

To date, HbA1c is the established reference metric for the assessment of glycemia in people with type 2 diabetes mellitus, based on its correlation with the incidence of microvascular and long-term macrovascular disease [137], although this assumption has been recently questioned [145]. The increasing use of CGM in type 2 diabetes mellitus has prompted a recent systematic review [146] of eleven studies, including a total of 13,987 people with type 2 diabetes mellitus, that evaluated the relationship between CGM measured TIR and diabetes complications. In each of these, the study cohort was not differentiated by treatment type. Four studies examined TIR in diabetic retinopathy and diabetic neuropathy, while seven studies evaluated TIR in diabetic peripheral neuropathy (DPN). A 10 % increase in TIR was associated with significant reductions in albuminuria, severity of retinopathy, and prevalence

of DPN and cardiac autonomic neuropathy (CAN).

Subsequent cross-sectional studies have confirmed that CGM-defined TIR is emerging as a relevant surrogate endpoint for microvascular complications, with increased TIR being associated with decreased rates of retinopathy [147,148], painful diabetic neuropathy [149] and preserved peripheral nerve function [150]. Both intra-day GV and time in hypoglycemia have been associated with retinal nerve-fiber thinning in retinopathy and neuropathy in type 2 diabetes mellitus [38,151], and GV has also been associated with cardiovascular autonomic neuropathy [152].

In terms of macrovascular outcomes, lower TIR is associated with surrogate vascular risk markers [153,154], increased risk of all-cause and CVD mortality [155], as well as peripheral artery disease [156] and diabetic foot ulcers [157]. Overall, in type 2 diabetes mellitus there is a consistent association between higher TIR and fewer macro- and microvascular complications [150,153,158–162]. Since the use of CGM is a driver for increased TIR in type 2 diabetes mellitus, the goal of reducing complications in type 2 diabetes mellitus can be associated with wider application of CGM systems, although further prospective longitudinal studies are necessary to validate these metrics for comprehensive risk profiling.

6. Primary care is a critical resource for management of obesity, intermediate hyperglycemia and type 2 diabetes mellitus

The majority of people living with obesity or intermediate hyperglycemia, who are at-risk of developing type 2 diabetes mellitus, and people with type 2 diabetes mellitus who are not on insulin therapy, are typically managed within primary care. The key goal for primary care teams must be the health promotion for at-risk populations and prevention of type 2 diabetes mellitus, especially amongst people living with obesity or intermediate hyperglycemia. This is made more-complex by the multitude of different populations of people at risk or with type 2 diabetes mellitus, the task of managing the multiple long-term conditions of associated macro- and microvascular disease, as well as their diverse ethnic and SDoH profiles. It is important that the landscape of digital health application evolves to meet this challenge. A 2018 survey reported that 89 % of primary care physicians (PCPs) across the EU did not use telemedicine with their patients and 81 % did not use it with other HCPs [163]. However, from 2020 onwards, the COVID-19 pandemic significantly changed the delivery of diabetes care to emphasize telemedicine in diabetes consultations [164], with evidence that this was not inferior to in-clinic consultations [164,165]. A small number of studies have also indicated that primary care teams and patients have embraced telemedicine as a consequence of the pandemic [166,167].

This trend is accompanied by evidence that use of CGM is increasing rapidly in adults with type 2 diabetes mellitus, with most growth in primary care. A review of e-health records from 30,585 adults with type 2 diabetes mellitus in Tennessee in the US [168] revealed a prevalence of 13 % of CGM use among this cohort, of whom 12 % were not on insulin. Twenty three percent of CGM users reported to only have visited their PCP during the previous year. Between 2020 and 2021, this study reported an overall monthly increase in new prescription rates for CGM of 36 % across all services, whereas the growth rate in primary care prescriptions was 125 % in the same period. New CGM users in primary care were mostly on intensive insulin regimen (49 %), whereas, 28 % were not using insulin.

The acknowledged efficacy of using CGM in people with dysglycemia should give primary care teams a significant tool to manage these populations, if access can be provided. However, experience with using and interpreting CGM systems, and the glucometric data that they provide, is a significant unmet need within primary care. Limited resources, with multiple competing priorities, are a barrier to uptake in primary care. A systematic review of the few studies available indicates that using CGM in the primary care setting for the management of adults

with type 2 diabetes mellitus is more effective than relying on SMBG for reducing HbA1c [169]. Just as important, people with type 2 diabetes mellitus reported that CGM improved treatment satisfaction, with better understanding of self-management needs and how therapy of type 2 diabetes mellitus works. PCPs reported that CGM and the glycemic reports generated assisted in effective communication with their patients and that they were willing to continue using CGM.

Using CGM and CGM-derived glucose data facilitates cooperation between PCPs and people with diabetes for clinically relevant, person-centered, achievable glycemic goal setting [170]. This empowers people with diabetes to manage the challenges of daily self-management and may even lead to healthy lifestyle behavior changes. Small-scale training for PCPs in the application and use of CGM by family-practice teams is well received and may be the best way to meet the needs of people with diabetes in their clinics [171]. Interest in applying CGM in their practices amongst PCPs has been assessed as higher among those with practices located further from endocrinology centers [172]. As with all settings, the management of type 2 diabetes mellitus in primary care services requires that access to CGM technology is improved, including reimbursement, to enable widespread adoption [173].

A further consideration that can support PCPs in using CGM devices and data with people who have intermediate hyperglycemia or are diagnosed with type 2 diabetes mellitus, is the use of AI-CDSS. Such tools are a natural development, since the management of people with dysglycemia and diabetes is laid out in comprehensive guidelines that are driven by clear treatment algorithms [108]. The development of AI-CDSS, that marry the decision paths laid out in guidelines with objective data on short and long-term glycemic performance, will create significant benefits for PCPs and people with intermediate hyperglycemia or type 2 diabetes mellitus in their care. At least one small-scale survey has indicated that PCPs have a generally positive view of AI, dependent on the context in which AI was adopted [174], and the use of AI-CDSS in the context of diabetes care is suited to the primary care setting, leveraging PCP expertise with knowledge-based machine learning.

Consensus statements. Primary-care management of CGM use in people living with obesity, intermediate hyperglycemia or type 2 diabetes mellitus.

- There is an unmet need for prospective studies in the primary-care setting, centered on the application of CGM technology in the management of people living with obesity, intermediate hyperglycemia or type 2 diabetes mellitus.
- Training and education on the effective application of CGM and interpretation of CGM data must be developed and delivered as part of continuing professional development for all primary-care team members.
- Access to support from specialist diabetes teams on the use of CGM and data interpretation is vital to establishing CGM use in the primary care setting.
- Development of and access to AI-enabled Clinical Decision Support Systems (AI-CDSS) for CGM data interpretation is critical for increasing time efficiency and wider adoption of CGM on the primary care level.
- Clear pathways and guidelines on patient eligibility and prescribing in obesity and intermediate hyperglycemia need to be developed.
- Support for proactive identification of eligible persons in the primary care setting will help to avoid worsening health inequalities.

7. Cost effectiveness of implementing CGM in the management of people living with obesity, intermediate hyperglycemia or type 2 diabetes mellitus

Application of CGM in the management of obesity, intermediate hyperglycemia and type 2 diabetes mellitus has the potential to significantly reduce the burden of disease for people living with each of these chronic conditions and the health services tasked with their care. The cost effectiveness of this intervention needs to be carefully analyzed. To date, there has been very limited cost-effectiveness analysis for use of CGM systems in type 2 diabetes mellitus, and less in either obesity or intermediate hyperglycemia. Of note, when looking at diabetes prevention strategies, for example in obesity or intermediate hyperglycemia, strategies that target populations from more-disadvantaged SDOH

groups may be cost-effective at higher input costs and lower efficacy compared to untargeted policies [175].

7.1. Type 2 diabetes mellitus on intensive insulin therapy

Use of CGM for people with type 2 diabetes mellitus on either MDI or CSII therapy has been assessed as cost-effective compared to standard care with SMBG, based on a 40-year horizon [110,111,176]. This includes total intervention costs and direct medical costs, as well as hospital admission costs for ADEs, in two of these analyses [111,176]. Productivity loss was included in the third study [110]. Use of CGM improves quality-adjusted life years (QALYs) for people with type 2 diabetes mellitus on intensive insulin therapy, which supports a favorable incremental cost-effectiveness ratio (ICER) [110,111,176].

7.2. Type 2 diabetes mellitus on basal insulin therapy

Intermittent use of CGM in four cycles over 3 months, followed up to 52 weeks, compared to using SMBG has been shown to be cost effective [119]. Direct medical costs, including treatment for depression and for diabetes complications were included. Improved Life expectancy and quality-adjusted life expectancy (QALE) outcomes for the CGM cohort generated gains for ICER and QALY. Notably, the cost-effectiveness in this model was attributed to users making informed behavioral choices without clinician guidance [119].

7.3. Type 2 diabetes mellitus on any insulin therapy

Using CGM in people with type 2 diabetes mellitus on intensive or basal insulin therapy was associated with increased QALYs and improved ICER over a lifetime horizon, compared with SMBG, based only on direct healthcare costs in the UK. Drivers included HbA1c reduction and reduced SMBG strip-testing [177].

7.4. Type 2 diabetes mellitus in the primary care setting

A randomized, 6-month prospective US trial was conducted using CGM compared to SMBG testing for participants receiving usual care in primary care clinics [178]. Of the 99 participants, 93 had a diagnosis of type 2 diabetes mellitus but were selected without consideration of their current dietary, oral medication or injectable therapeutic regimens. After 6 months, CGM users had reduced costs overall for primary care visits, emergency department attendance and laboratory investigations. Savings were not universal and depended on the health insurance provider.

Consensus statements. Cost-effectiveness of CGM use in people living with obesity, intermediate hyperglycemia or type 2 diabetes mellitus.

- There is an unmet need for cost-effectiveness analysis of using CGM technology as part of intervention strategies to modify diet and physical activity in people living with obesity, intermediate hyperglycemia or type 2 diabetes mellitus.
- Future health-economic models should incorporate measures of SDOH to improve targeting of CGM as an intervention strategy for prevention or delayed progression of type 2 diabetes mellitus.

8. Summary

Based on available evidence, CGM technology can be used to support behavioral change and improved glycemic control for people at-risk of type 2 diabetes mellitus, including people living with obesity or intermediate hyperglycemia, and for people diagnosed with type 2 diabetes mellitus. The perceived and demonstrated benefits include: (1) to actively engage people with their glycemic status, (2) to evaluate treatment responses and achievement of goals; (3) to reinforce education and self-management skills in people with intermediate hyperglycemia or with type 2 diabetes mellitus; (4) adjust therapy as needed for

people with type 2 diabetes mellitus. Significant further research is needed to fully evaluate the value proposition for CGM technology in normoglycemic obese at-risk persons and in those with intermediate hyperglycemia. It is important to state that the benefits of the application of CGM technology will only be fully realized with significant widening of access globally, to avoid regional disparities and inequities, along with further refinements to improve simplicity and interpretation of CGM data to include people with lower levels of digital and health literacy, where AI-CDSS may play a critical role.

CRedit authorship contribution statement

Tadej Battelino: Writing – review & editing, Writing – original draft, Conceptualization. **Nebojsa Lalic:** Writing – review & editing, Writing – original draft, Conceptualization. **Sufyan Hussain:** Writing – review & editing, Writing – original draft, Conceptualization. **Antonio Ceriello:** Writing – review & editing, Writing – original draft, Conceptualization. **Sanja Klobucar:** Writing – review & editing, Writing – original draft, Conceptualization. **Sarah J. Davies:** Writing – review & editing, Writing – original draft, Conceptualization. **Pinar Topsever:** Writing – review & editing, Conceptualization. **Julie Heverly:** Writing – review & editing, Conceptualization. **Francesca Ulivi:** Writing – review & editing, Conceptualization. **Kevin Brady:** Writing – review & editing, Conceptualization. **Tsvetalana Tankova:** Writing – review & editing, Writing – original draft, Conceptualization. **Júlia Galhardo:** Writing – review & editing, Conceptualization. **Kostas Tagkalos:** Writing – review & editing, Conceptualization. **Erik Werson:** Writing – review & editing, Conceptualization. **Chantal Mathieu:** Writing – review & editing, Conceptualization. **Peter Schwarz:** Writing – review & editing, Writing – original draft, Conceptualization.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: TB served on advisory panels of Novo Nordisk, Sanofi, Eli Lilly, Boehringer, Medtronic, Abbott, Indigo Diabetes. TB received honoraria for participating on the speaker's bureau of Eli Lilly, Novo Nordisk, Medtronic, Abbott, Sanofi, Dexcom, Aventis, Astra Zeneca, and Roche. TB's Institution received research grant support from Abbott, Medtronic, Novo Nordisk, Sanofi, Novartis, Sandoz, and Zealand Pharma, Slovenian Research and Innovation Agency, the National Institutes of Health, and the European Union. SH has served on the advisory board for Tandem, Dexcom, Medtronic, Sanofi, Vertex; undertaken non-promotional educational and/or consultancy work for Abbott UK, Insulet, Dexcom, and Roche; SH's institution has received research grant support from Abbott and Insulet. PT served on advisory boards of Eli Lilly, Boehringer Ingelheim, Novo Nordisk, Lifescan and Astra Zeneca. PT participated in CME's supported by Astra Zeneca, Boehringer Ingelheim, Eli Lilly and Novo Nordisk. PT is the vice chair of PCDE, which has received corporate sponsorship from AstraZeneca, Eli Lilly, Novo Nordisk, and Roche Diagnostics, but the companies had no input in the study protocols. TT has served on advisory panels of Boehringer Ingelheim, Astra Zeneca, NovoNordisk, EliLilly, Medtronic. TT has received honoraria for participating in speaker's bureau of Boehringer Ingelheim, Astra Zeneca, NovoNordisk, Eli Lilly, Merck, Sanofi, MSD, Servier. CM serves or has served on the advisory panel for Dexcom, Roche, Medtronic, Insulet. Financial compensation for these activities has been received by KU Leuven; KU Leuven has received research support for CM from Medtronic. CM serves or has served on the speakers bureau for Medtronic. Financial compensation for these activities has been received by KU Leuven. CM is president of EASD. All external support of EASD is to be found on <https://www.easd.org>.

Acknowledgements

The author group wish to thank The International Diabetes Federation Europe (IDF-Eu) for inviting and organizing the author working group and for providing funding to Robert Brines, Bite Medical Consulting, who supported the author group by collating and compiling author revisions during the manuscript drafting process. TB was supported in part by the Slovenian Research and Innovation Agency grant # P3-0343. SH is supported by the Medical Research Council Clinical Academic Partnership award (MR/W030004/1).

References

- [1] Zheng Y, Manson JE, Yuan C, Liang MH, Grodstein F, Stampfer MJ, et al. Associations of weight gain from early to middle adulthood with major health outcomes later in life. *JAMA* 2017;318:255–69. <https://doi.org/10.1001/jama.2017.7092>.
- [2] de Mutser R, Sun Q, Willett WC, Hu FB, van Dam RM. Overweight in early adulthood, adult weight change, and risk of type 2 diabetes, cardiovascular diseases, and certain cancers in men: a cohort study. *Am J Epidemiol* 2014;179: 1353–65. <https://doi.org/10.1093/aje/kwu052>.
- [3] Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403. <https://doi.org/10.1056/nejmoa012512>.
- [4] Li G, Zhang P, Wang J, Gregg EW, Yang W, Gong Q, et al. The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study. *Lancet* 2008;371:1783–9. [https://doi.org/10.1016/S0140-6736\(08\)60766-7](https://doi.org/10.1016/S0140-6736(08)60766-7).
- [5] Nathan DM, Barrett-Connor E, Crandall JP, Edelstein SL, Goldberg RB, Horton ES, et al. Long-term effects of lifestyle intervention or metformin on diabetes development and microvascular complications over 15-year follow-up: the Diabetes Prevention Program Outcomes Study. *Lancet Diabetes Endocrinol* 2015; 3:866–75. [https://doi.org/10.1016/S2213-8587\(15\)00291-0](https://doi.org/10.1016/S2213-8587(15)00291-0).
- [6] Lindström J, Peltonen M, Eriksson JG, Ilanne-Parikka P, Aunola S, Keinänen-Kiukaanniemi S, et al. Improved lifestyle and decreased diabetes risk over 13 years: long-term follow-up of the randomised Finnish Diabetes Prevention Study (DPS). *Diabetologia* 2013;56:284–93. <https://doi.org/10.1007/s00125-012-2752-5>.
- [7] Li G, Hu Y, Yang W, Jiang Y, Wang J, Xiao J, et al. Effects of insulin resistance and insulin secretion on the efficacy of interventions to retard development of type 2 diabetes mellitus: the DA Qing IGT and Diabetes Study. *Diabetes Res Clin Pract* 2002;58:193–200. [https://doi.org/10.1016/S0168-8227\(02\)00175-4](https://doi.org/10.1016/S0168-8227(02)00175-4).
- [8] Diabetes Prevention Program (DPP) Research Group. The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes Care* 2002;25: 2165–71. <https://doi.org/10.2337/diacare.25.12.2165>.
- [9] Chen J-Y, Deng S, Wagatsuma Y. Combination of risks of BMI and health-related lifestyles on kidney function in the prediabetic Japanese population: a prospective cohort study. *Int J Environ Res Public Health* 2023;20:5338. <https://doi.org/10.3390/ijerph20075338>.
- [10] Lind M, Polonsky W, Hirsch IB, Heise T, Bolinder J, Dahlqvist S, et al. Continuous glucose monitoring vs conventional therapy for glycemic control in adults with type 1 diabetes treated with multiple daily insulin injections: the GOLD Randomized Clinical Trial. *JAMA* 2017;317:379–87. <https://doi.org/10.1001/jama.2016.19976>.
- [11] Beck RW, Riddlesworth T, Ruedy K, Ahmann A, Bergenstal R, Haller S, et al. Effect of continuous glucose monitoring on glycemic control in adults with type 1 diabetes using insulin injections: the DIAMOND Randomized Clinical Trial. *JAMA* 2017;317:371–8. <https://doi.org/10.1001/jama.2016.19975>.
- [12] Leelarathna L, Evans ML, Neupane S, Rayman G, Lumley S, Cranston I, et al. Intermittently scanned continuous glucose monitoring for type 1 diabetes. *N Engl J Med* 2022;387:1477–87. <https://doi.org/10.1056/nejmoa2205650>.
- [13] Bolinder J, Antuna R, Geelhoed-Duijvestijn P, Kröger J, Weitgasser R. Novel glucose-sensing technology and hypoglycaemia in type 1 diabetes: a multicentre, non-masked, randomised controlled trial. *Lancet* 2016;388:2254–63. [https://doi.org/10.1016/S0140-6736\(16\)31535-5](https://doi.org/10.1016/S0140-6736(16)31535-5).
- [14] Campbell FM, Murphy NP, Stewart C, Biester T, Kordonouri O. Outcomes of using flash glucose monitoring technology by children and young people with type 1 diabetes in a single arm study. *Pediatr Diabetes* 2018;19:1294–301. <https://doi.org/10.1111/pedi.12735>.
- [15] Cox DJ, Taylor AG, Moncrief M, Diamond A, Yancy WS, Hegde S, et al. Continuous glucose monitoring in the self-management of type 2 diabetes: a paradigm shift. *Diabetes Care* 2016;39:e71–3. <https://doi.org/10.2337/dc15-2836>.
- [16] Allen NA, Fain JA, Braun B, Chipkin SR. Continuous glucose monitoring counseling improves physical activity behaviors of individuals with type 2 diabetes: a randomized clinical trial. *Diabetes Res Clin Pract* 2008;80:371–9. <https://doi.org/10.1016/j.diabres.2008.01.006>.
- [17] Yoo HJ, An HG, Park SY, Ryu OH, Kim HY, Seo JA, et al. Use of a real time continuous glucose monitoring system as a motivational device for poorly controlled type 2 diabetes. *Diabetes Res Clin Pract* 2008;82:73–9. <https://doi.org/10.1016/j.diabres.2008.06.015>.

- [18] Ahn Y-C, Kim YS, Kim B, Ryu JM, Kim MS, Kang M, et al. Effectiveness of non-contact dietary coaching in adults with diabetes or prediabetes using a continuous glucose monitoring device: a Randomized Controlled Trial. *Healthcare* 2023;11:252. <https://doi.org/10.3390/healthcare11020252>.
- [19] Liu Y, Guo H, Wang Q, Chen J, Xuan Y, Xu J, et al. Short-term effects of lifestyle intervention in the reversion to normoglycemia in people with prediabetes. *Prim Care Diabetes* 2022;16:168–72. <https://doi.org/10.1016/j.pcd.2021.12.009>.
- [20] Coppel KJ, Abel S, Whitehead LC, Tangiora A, Spedding T, Tipene-Leach D. A diagnosis of prediabetes when combined with lifestyle advice and support is considered helpful rather than a negative label by a demographically diverse group: a qualitative study. *Prim Care Diabetes* 2022;16:301–6. <https://doi.org/10.1016/j.pcd.2021.10.003>.
- [21] Thomas F, Pannier B, Meghrief K, Galtier D, Lamande J, Raison J, et al. Adapted educational health program among deprived subjects with prediabetes. *Prim Care Diabetes* 2022;16:664–9. <https://doi.org/10.1016/j.pcd.2022.06.003>.
- [22] Ludwig DS, Ebbeling CB. The carbohydrate-insulin model of obesity: beyond “calories in, calories out.”. *JAMA Intern Med* 2018;178:1098. <https://doi.org/10.1001/jamainternmed.2018.2933>.
- [23] Hall KD. A review of the carbohydrate-insulin model of obesity. *Eur J Clin Nutr* 2017;71:323–6. <https://doi.org/10.1038/ejcn.2016.260>.
- [24] Cooper DJ, Zarabi S, Farrand B, Becker A, Roslin M. Continuous glucose monitoring reveals similar glycemic variability in individuals with obesity despite increased HOMA-IR. *Front Nutr* 2022;9:1070187. <https://doi.org/10.3389/fnut.2022.1070187>.
- [25] Salkind SJ, Huizenga R, Fonda SJ, Walker MS, Vigorsky RA. Glycemic variability in nondiabetic morbidly obese persons. *J Diabetes Sci Technol* 2014;8:1042–7. <https://doi.org/10.1177/1932296814537039>.
- [26] Kishimoto I, Ohashi A. Impact of lifestyle behaviors on postprandial hyperglycemia during continuous glucose monitoring in adult males with overweight/obesity but without diabetes. *Nutrients* 2021;13:3092. <https://doi.org/10.3390/nu13093092>.
- [27] Kashiwagi K, Inaishi J, Kinoshita S, Wada Y, Hanashiro S, Shiga K, et al. Assessment of glycemic variability and lifestyle behaviors in healthy nondiabetic individuals according to the categories of body mass index. *PLoS One* 2023;18:e0291923. <https://doi.org/10.1371/journal.pone.0291923>.
- [28] Liu K, Choi TST, Zhao L, Teong XT, Hutchison AT, Heilbronn LK. A qualitative exploration of behaviour change and maintenance experience in people with overweight or obesity in a dietary intervention. *Nutr Diet* 2023;296–305. <https://doi.org/10.1111/1747-0080.12855>.
- [29] Bailey KJ, Little JP, Jung ME. Self-monitoring using continuous glucose monitors with real-time feedback improves exercise adherence in individuals with impaired blood glucose: a pilot study. *Diabetes Technol Ther* 2016;18:185–93. <https://doi.org/10.1089/dia.2015.0285>.
- [30] Hutchison AT, Regmi P, Manogian ENC, Fleischer JG, Wittert GA, Panda S, et al. Time-restricted feeding improves glucose tolerance in men at risk for type 2 diabetes: a randomized crossover trial. *Obesity* 2019;27:724–32. <https://doi.org/10.1002/oby.22449>.
- [31] Naguib MN, Hegedus E, Raymond JK, Goran MI, Salvy S-J, Wee CP, et al. Continuous glucose monitoring in adolescents with obesity: monitoring of glucose profiles, glucose excursions, and adherence to time restricted eating programs. *Front Endocrinol* 2022;13:841838. <https://doi.org/10.3389/fendo.2022.841838>.
- [32] Ehrhardt N, Zaghaf EA. Behavior modification in prediabetes and diabetes: potential use of real-time continuous glucose monitoring. *J Diabetes Sci Technol* 2019;13:271–5. <https://doi.org/10.1177/1932296818790994>.
- [33] Bellou V, Belbasis L, Tzoulaki I, Evangelou E. Risk factors for type 2 diabetes mellitus: an exposure-wide umbrella review of meta-analyses. *PLoS One* 2018;13:e0194127. <https://doi.org/10.1371/journal.pone.0194127>.
- [34] American Diabetes Association Professional Practice Committee. 2. Diagnosis and Classification of Diabetes: Standards of Care in Diabetes—2024. *Diabetes Care* 2024;47:S20–42. <https://doi.org/10.2337/dc24-s002>.
- [35] Bergman M, Manco M, Satman I, Chan J, Schmidt MI, Sesti G, et al. International Diabetes Federation Position Statement on the 1-hour post-load plasma glucose for the diagnosis of intermediate hyperglycaemia and type 2 diabetes. *Diabetes Res Clin Pract* 2024;209:111589. <https://doi.org/10.1016/j.diabres.2024.111589>.
- [36] Welsh C, Welsh P, Celis-Morales CA, Mark PB, Mackay D, Ghouri N, et al. Glycated hemoglobin, prediabetes, and the links to cardiovascular disease: data from UK Biobank. *Diabetes Care* 2020;43:440–5. <https://doi.org/10.2337/dc19-1683>.
- [37] Hjort A, Iggman D, Rosqvist F. Glycemic variability assessed using continuous glucose monitoring in individuals without diabetes and associations with cardiometabolic risk markers: a systematic review and meta-analysis. *Clin Nutr* 2024;43:915–25. <https://doi.org/10.1016/j.clnu.2024.02.014>.
- [38] van der Heide FCT, Foreman YD, Franken IWM, Henry RMA, Kroon AA, Dagnelie PC, et al. (Pre)diabetes, glycemia, and daily glucose variability are associated with retinal nerve fiber layer thickness in The Maastricht Study. *Sci Rep UK* 2022;12:17750. <https://doi.org/10.1038/s41598-022-22748-2>.
- [39] Tapp RJ, Tikellis G, Wong TY, Harper CA, Zimmet PZ, Shaw JE, et al. Longitudinal association of glucose metabolism with retinopathy. *Diabetes Care* 2008;31:1349–54. <https://doi.org/10.2337/dc07-1707>.
- [40] Plantinga LC, Crews DC, Coresh J, Miller ER, Saran R, Yee J, et al. Prevalence of chronic kidney disease in US adults with undiagnosed diabetes or prediabetes. *Clin J Am Soc Nephrol* 2010;5:673–82. <https://doi.org/10.2215/cjn.07891109>.
- [41] White NH, Pan Q, Knowler WC, Schroeder EB, Dabelea D, Chew EY, et al. Risk factors for the development of retinopathy in prediabetes and type 2 diabetes: the diabetes prevention program experience. *Diabetes Care* 2022;45:2653–61. <https://doi.org/10.2337/dc22-0860>.
- [42] Sörensen BM, Houben AJHM, Berendschot TTJM, Schouten JSAG, Kroon AA, van der Kallen CJH, et al. Prediabetes and type 2 diabetes are associated with generalized microvascular dysfunction. *Circulation* 2016;134:1339–52. <https://doi.org/10.1161/circulationaha.116.023446>.
- [43] Monnier L, Colette C, Dunseath GJ, Owens DR. The loss of postprandial glycemic control precedes stepwise deterioration of fasting with worsening diabetes. *Diabetes Care* 2007;30:263–9. <https://doi.org/10.2337/dc06-1612>.
- [44] Miller E, Polonsky WH, Miller K. What role might there be for continuous glucose monitoring in the assessment of diabetes risk? *Diabetes Technol Ther* 2023;25:S14–20. <https://doi.org/10.1089/dia.2023.0136>.
- [45] Acciaroli G, Sparacino G, Hakaste L, Facchinetti A, Nunzio GMD, Palombi A, et al. Diabetes and prediabetes classification using glycemic variability indices from continuous glucose monitoring data. *J Diabetes Sci Technol* 2018;12:105–13. <https://doi.org/10.1177/1932296817710478>.
- [46] Peyser TA, Balo AK, Buckingham BA, Hirsch IB, Garcia A. Glycemic variability percentage: a novel method for assessing glycemic variability from continuous glucose monitor data. *Diabetes Technol Ther* 2018;20:6–16. <https://doi.org/10.1089/dia.2017.0187>.
- [47] Frank S, Hames KC, Jbaily A, Park JH, Stroyeck C, Price D. Feasibility of using a factory-calibrated continuous glucose monitoring system to diagnose type 2 diabetes. *Diabetes Technol Ther* 2022;24:907–14. <https://doi.org/10.1089/dia.2022.0189>.
- [48] Shah VN, DuBose SN, Li Z, Beck RW, Peters AL, Weinstock RS, et al. Continuous glucose monitoring profiles in healthy nondiabetic participants: a multicenter prospective study. *J Clin Endocrinol Metab* 2019;104:4356–64. <https://doi.org/10.1210/je.2018-02763>.
- [49] Group JDRFGMS, Fox LA, Beck RW, Xing D. Variation of interstitial glucose measurements assessed by continuous glucose monitors in healthy, nondiabetic individuals. *Diabetes Care* 2010;33:1297–9. <https://doi.org/10.2337/dc09-1971>.
- [50] Marco A, Pazos-Couselo M, Moreno-Fernandez J, Díez-Fernández A, Alonso-Sampedro M, Fernández-Merino C, et al. Time above range for predicting the development of type 2 diabetes. *Front Public Health* 2022;10:1005513. <https://doi.org/10.3389/fpubh.2022.1005513>.
- [51] Cichosz SL, Kronborg T, Laugesen E, Hangaard S, Fleischer J, Hansen TK, et al. From stability to variability: classification of healthy individuals, prediabetes, and type 2 diabetes using glycemic variability indices from continuous glucose monitoring data. *Diabetes Technol Ther* 2025;27:34–44. <https://doi.org/10.1089/dia.2024.0226>.
- [52] Barua S, Sabharwal A, Glantz N, Conneely C, Larez A, Bevier W, et al. Dysglycemia in adults at risk for or living with non-insulin treated type 2 diabetes: Insights from continuous glucose monitoring. *Eclinicalmedicine* 2021;35:100853. <https://doi.org/10.1016/j.eclinm.2021.100853>.
- [53] Hall H, Perelman D, Breschi A, Limcaoco P, Kellogg R, McLaughlin T, et al. Glucotypes reveal new patterns of glucose dysregulation. *PLoS Biol* 2018;16:e2005143. <https://doi.org/10.1371/journal.pbio.2005143>.
- [54] Chakarova N, Dimova R, Grozeva G, Tankova T. Assessment of glucose variability in subjects with prediabetes. *Diabetes Res Clin Pract* 2019;151:56–64. <https://doi.org/10.1016/j.diabres.2019.03.038>.
- [55] Rizos EC, Kanellopoulou A, Filis P, Markozannes G, Chaliasos K, Ntzani EE, Tzamouranou A, Tentolouris N, Tsilidis KK. Difference on glucose profile from continuous glucose monitoring in people with prediabetes vs. normoglycemic individuals: a matched-pair analysis. *J Diabetes Sci Technol* 2024;18(2):414–22. <https://doi.org/10.1177/19322968221123530>.
- [56] Hanefeld M, Sulk S, Helbig M, Thomas A, Köhler C. Differences in glycemic variability between normoglycemic and prediabetic subjects. *J Diabetes Sci Technol* 2014;8:286–90. <https://doi.org/10.1177/1932296814522739>.
- [57] Dorcelly B, Sifonte E, Popp C, Divakaran A, Katz K, Musleh S, et al. Continuous glucose monitoring and 1-h plasma glucose identifies glycemic variability and dysglycemia in high-risk individuals with HbA1c < 5.7%: a pilot study. *Endocrine* 2022;77:403–7. <https://doi.org/10.1007/s12020-022-03109-5>.
- [58] Dimova R, Chakarova N, Daniele G, Bianchi C, Dardano A, Prato SD, et al. Insulin secretion and action affect glucose variability in the early stages of glucose intolerance. *Diabetes Metab Res Rev* 2022;38:e3531. <https://doi.org/10.1002/dmrr.3531>.
- [59] Schön M, Prystupa K, Mori T, Zaharia OP, Bódis K, Bombrich M, et al. Analysis of type 2 diabetes heterogeneity with a tree-like representation: insights from the prospective German Diabetes Study and the LURIC cohort. *Lancet Diabetes Endocrinol* 2024;12:119–31. [https://doi.org/10.1016/s2213-8587\(23\)00329-7](https://doi.org/10.1016/s2213-8587(23)00329-7).
- [60] Peng M, He S, Wang J, An Y, Qian X, Zhang B, et al. Efficacy of 1-hour postload plasma glucose as a suitable measurement in predicting type 2 diabetes and diabetes-related complications: a post hoc analysis of the 30-year follow-up of the Da Qing IGT and Diabetes Study. *Diabetes Obes Metab* 2024;26:2329–38. <https://doi.org/10.1111/dom.15547>.
- [61] Spartano NL, Sultana N, Lin H, Cheng H, Lu S, Fei D, Murabito JM, Walker ME, Wolpert HA, Steenkamp DW. Defining Continuous glucose monitor time in range in a large, community-based cohort without diabetes. *J Clin Endocrinol Metab* 2025;110(4):1128–34. <https://doi.org/10.1210/clinem/dgae626>.
- [62] Miller E, Miller K. Detection and intervention: use of continuous glucose monitoring in the early stages of type 2 diabetes. *Clin Diabetes* 2024;42:398–407. <https://doi.org/10.2337/cd23-0077>.
- [63] Dimova R, Chakarova N, Prato SD, Tankova T. The relationship between dietary patterns and glycemic variability in people with impaired glucose tolerance. *J Nutr* 2023;153:1427–38. <https://doi.org/10.1016/j.tjnut.2023.03.007>.

- [64] Kitazawa M, Takeda Y, Hatta M, Horikawa C, Sato T, Osawa T, et al. Lifestyle intervention with smartphone app and isCGM for people at high risk of type 2 diabetes: randomized trial. *J Clin Endocrinol Metab* 2024;109:1060–70. <https://doi.org/10.1210/clinem/dgad639>.
- [65] Lee JY, Nguyen JT, Arroyo J, Tran T, Hanami D, Mayorga J. Feasibility and acceptability of using flash glucose monitoring system sensors to empower lifestyle changes in people with prediabetes. *Diabetes Care* 2023;46:e10–1. <https://doi.org/10.2337/dc22-0612>.
- [66] Yost O, DeJonckheere M, Stonebraker S, Ling G, Buis L, Pop-Busui R, et al. Continuous glucose monitoring with low-carbohydrate diet coaching in adults with prediabetes: mixed methods pilot study. *JMIR Diabetes* 2020;5:e21551. <https://doi.org/10.2196/21551>.
- [67] Lean MEJ, Leslie WS, Barnes AC, Brosnahan N, Thom G, McCombie L, et al. Durability of a primary care-led weight-management intervention for remission of type 2 diabetes: 2-year results of the DIRECT open-label, cluster-randomised trial. *Lancet Diabetes Endocrinol* 2019;7:344–55. [https://doi.org/10.1016/s2213-8587\(19\)30068-3](https://doi.org/10.1016/s2213-8587(19)30068-3).
- [68] Rothberg A, Lean M, Laferrière B. Remission of type 2 diabetes: always more questions, but enough answers for action. *Diabetologia* 2024;67:602–10. <https://doi.org/10.1007/s00125-023-06069-1>.
- [69] Riddle MC, Cefalu WT, Evans PH, Gerstein HC, Nauck MA, Oh WK, et al. Consensus report: definition and interpretation of remission in type 2 diabetes. *Diabetes Care* 2021;44:2438–44. <https://doi.org/10.2337/dci21-0034>.
- [70] Nagi D, Hambling C, Taylor R. Remission of type 2 diabetes: a position statement from the Association of British Clinical Diabetologists (ABCD) and the primary care diabetes society (PCDS). *Br J Diabetes* 2019;19:73–6.
- [71] Lean ME, Leslie WS, Barnes AC, Brosnahan N, Thom G, McCombie L, et al. Primary care-led weight management for remission of type 2 diabetes (DIRECT): an open-label, cluster-randomised trial. *Lancet* 2018;391:541–51. [https://doi.org/10.1016/s0140-6736\(17\)33102-1](https://doi.org/10.1016/s0140-6736(17)33102-1).
- [72] Hocking SL, Markovic TP, Lee CMY, Picone TJ, Gudorf KE, Colagiuri S. Intensive lifestyle intervention for remission of early type 2 diabetes in primary care in australia: DIRECT-Aus. *Diabetes Care* 2024;47:66–70. <https://doi.org/10.2337/dc23-0781>.
- [73] Taheri S, Zaghoul H, Chagoury O, Elhadad S, Ahmed SH, Khatib NE, et al. Effect of intensive lifestyle intervention on bodyweight and glycaemia in early type 2 diabetes (DIADeM-1): an open-label, parallel-group, randomised controlled trial. *Lancet Diabetes Endocrinol* 2020;8:477–89. [https://doi.org/10.1016/s2213-8587\(20\)30117-0](https://doi.org/10.1016/s2213-8587(20)30117-0).
- [74] Corbin LJ, Hughes DA, Bull CJ, Vincent EE, Smith ML, McConnachie A, et al. The metabolomic signature of weight loss and remission in the Diabetes Remission Clinical Trial (DIRECT). *Diabetologia* 2024;67:74–87. <https://doi.org/10.1007/s00125-023-06019-x>.
- [75] Gregg EW, Chen H, Wagenknecht LE, Clark JM, Delahanty LM, Bantle J, et al. Association of an intensive lifestyle intervention with remission of type 2 diabetes. *JAMA* 2012;308:2489–96. <https://doi.org/10.1001/jama.2012.67929>.
- [76] Gregg EW, Chen H, Bancks MP, Manalac R, Maruthur N, Munshi M, et al. Impact of remission from type 2 diabetes on long-term health outcomes: findings from the Look AHEAD study. *Diabetologia* 2024;67:459–69. <https://doi.org/10.1007/s00125-023-06048-6>.
- [77] Gibson AA, Sainsbury A. Strategies to improve adherence to dietary weight loss interventions in research and real-world settings. *Behav Sci* 2017;7:44. <https://doi.org/10.3390/bs7030044>.
- [78] Shand JAD, Young S, Verster F, Peters C. Pilot study to test the safety, tolerability and feasibility of dulaglutide during a low-energy diet for weight loss and improved glycaemic control. *BMJ Nutr, Prev Health* 2023;6:341–6. <https://doi.org/10.1136/bmjnp-2023-000733>.
- [79] van Bloemendaal L, IJzerman RG, Ten Kulve JS, Barkhof F, Konrad RJ, Drent ML, Veltman DJ, Diamant M. GLP-1 receptor activation modulates appetite and reward-related brain areas in humans. *Diabetes* 2014;63:4186–96. <https://doi.org/10.2337/db14-0849>.
- [80] van Bloemendaal L, ten Kulve JS, la Fleur SE, IJzerman RG, Diamant M. Effects of glucagon-like peptide 1 on appetite and body weight: focus on the CNS. *J Endocrinol* 2014;221:T1–16. <https://doi.org/10.1530/joe-13-0414>.
- [81] Dunn TC, Ajan RA, Bergenstal RM, Xu Y. Is it time to move beyond TIR to TITR? Real-world data from over 20,000 users of continuous glucose monitoring in patients with type 1 and type 2 diabetes. *Diabetes Technol Ther* 2024;26:203–10. <https://doi.org/10.1089/dia.2023.0565>.
- [82] Oser TK, Cucuzzella M, Stasinopoulos M, Moncrief M, McCall A, Cox DJ. An innovative, paradigm-shifting lifestyle intervention to reduce glucose excursions with the use of continuous glucose monitoring to educate, motivate, and activate adults with newly diagnosed type 2 diabetes: pilot feasibility study. *JMIR Diabetes* 2022;7:e34465. <https://doi.org/10.2196/34465>.
- [83] Jiménez A, Ceriello A, Casamitjana R, Flores L, Viaplana-Masclans J, Vidal J. Remission of type 2 diabetes after Roux-en-Y Gastric Bypass or Sleeve gastrectomy is associated with a distinct glycemic profile. *Ann Surg* 2015;261:316–22. <https://doi.org/10.1097/sla.0000000000000586>.
- [84] Sawada S, Kodama S, Tsuchiya S, Kurosawa S, Endo A, Sugawara H, et al. Continuous glucose monitoring in patients with remission of type 2 diabetes after laparoscopic sleeve gastrectomy without or with duodenojejunal bypass. *Clin Obes* 2020;10:e12409. <https://doi.org/10.1111/cob.12409>.
- [85] Turquetil A, Morello R, Joubert M, Roux YL, Reznik Y. Early continuous glucose monitoring for predicting remission of type 2 diabetes 1 year after bariatric surgery. *Diabetes Metab* 2021;47:101255. <https://doi.org/10.1016/j.diabet.2021.101255>.
- [86] Taylor PJ, Thompson CH, Luscombe-Marsh ND, Wycherley TP, Wittert G, Brinkworth GD. Efficacy of real-time continuous glucose monitoring to improve effects of a prescriptive lifestyle intervention in type 2 diabetes: a pilot study. *Diabetes Ther* 2019;10:509–22. <https://doi.org/10.1007/s13300-019-0572-z>.
- [87] Taylor PJ, Thompson CH, Brinkworth GD. Effectiveness and acceptability of continuous glucose monitoring for type 2 diabetes management: a narrative review. *J Diabetes Investig* 2018;9:713–25. <https://doi.org/10.1111/jdi.12807>.
- [88] Allen N, Whittemore R, Melkus G. A continuous glucose monitoring and problem-solving intervention to change physical activity behavior in women with type 2 diabetes: a pilot study. *Diabetes Technol Ther* 2011;13:1091–9. <https://doi.org/10.1089/dia.2011.0088>.
- [89] Ida S, Kaneko R, Imataka K, Okubo K, Shirakura Y, Azuma K, et al. Effects of flash glucose monitoring on dietary variety, physical activity, and self-care behaviors in patients with diabetes. *J Diabetes Res* 2020;2020:9463648. <https://doi.org/10.1155/2020/9463648>.
- [90] Khunti K, Gomes MB, Pocock S, Shestakova MV, Pintat S, Fenici P, et al. Therapeutic inertia in the treatment of hyperglycaemia in patients with type 2 diabetes: a systematic review. *Diabetes Obes Metab* 2018;20:427–37. <https://doi.org/10.1111/dom.13088>.
- [91] Reach G, Pechter V, Gentilella R, Corcos A, Ceriello A. Clinical inertia and its impact on treatment intensification in people with type 2 diabetes mellitus. *Diabetes Metab* 2017;43:501–11. <https://doi.org/10.1016/j.diabet.2017.06.003>.
- [92] Giugliano D, Maiorino MI, Bellastella G, Esposito K. Clinical inertia, reverse clinical inertia, and medication non-adherence in type 2 diabetes. *J Endocrinol Invest* 2019;42:495–503. <https://doi.org/10.1007/s40618-018-0951-8>.
- [93] Russell-Jones D, Pouwer F, Khunti K. Identification of barriers to insulin therapy and approaches to overcoming them. *Diabetes Obes Metabol* 2018;20:488–96. <https://doi.org/10.1111/dom.13132>.
- [94] Polonsky WH, Fisher L, Guzman S, Villa-Caballero L, Edelman SV. Psychological insulin resistance in patients with type 2 diabetes. *Diabetes Care* 2005;28:2543–5. <https://doi.org/10.2337/diacare.28.10.2543>.
- [95] Peyrot M, Barnett AH, Meneghini LF, Schumm-Draeger P-M. Insulin adherence behaviours and barriers in the multinational Global Attitudes of Patients and Physicians in Insulin Therapy study. *Diabet Med* 2012;29:682–9. <https://doi.org/10.1111/j.1464-5491.2012.03605.x>.
- [96] Group AS, Gerstein HC, Miller ME, Genuth S, Ismail-Beigi F, Buse JB, et al. Long-term effects of intensive glucose lowering on cardiovascular outcomes. *N Engl J Med* 2011;364:818–28. <https://doi.org/10.1056/nejmoa1006524>.
- [97] Zoungas S, Patel A, Chalmers J, de Galan BE, Li Q, Billot L, et al. Severe hypoglycemia and risks of vascular events and death. *N Engl J Med* 2010;363:1410–8. <https://doi.org/10.1056/nejmoa1003795>.
- [98] Hohendorff J, Witek P, Kania M, Sudol M, Hajduk K, Stepien A, et al. Higher scanning frequency is correlated with less fear of hypoglycemia in type 1 diabetes patients using isCGM. *Front Endocrinol* 2022;13:996933. <https://doi.org/10.3389/fendo.2022.996933>.
- [99] Teasdale SL, Griffin A, Barrett HL, Coutts C, Vitanza M, Headey A. Continuous glucose monitoring in young adults with type 1 diabetes: impact on hypoglycemia confidence and fear. *Diabetes Spectr* 2022;35:322–6. <https://doi.org/10.2337/ds21-0066>.
- [100] Harris SB, Levrat-Guillen F. Use of the FreeStyle Libre system and diabetes treatment progression in T2DM: Results from a retrospective cohort study using a Canadian private payer claims database. *Diabetes Obes Metab* 2023;25:1704–13. <https://doi.org/10.1111/dom.15025>.
- [101] Beck RW, Riddleworth TD, Ruedy K, Ahmann A, Haller S, Kruger D, et al. Continuous glucose monitoring versus usual care in patients with type 2 diabetes receiving multiple daily insulin injections. *Ann Intern Med* 2017;167:365–74. <https://doi.org/10.7326/m16-2855>.
- [102] Yaron M, Roitman E, Aharon-Hananel G, Landau Z, Ganz T, Yanuv I, et al. Effect of flash glucose monitoring technology on glycemic control and treatment satisfaction in patients with type 2 diabetes. *Diabetes Care* 2019;42:1178–84. <https://doi.org/10.2337/dc18-0166>.
- [103] Haak T, Hanaire H, Ajjan R, Hermanns N, Riveline J-P, Rayman G. Flash glucose-sensing technology as a replacement for blood glucose monitoring for the management of insulin-treated type 2 diabetes: a multicenter. Open-Label Randomized Controlled Trial *Diabetes Ther* 2017;8:55–73. <https://doi.org/10.1007/s13300-016-0223-6>.
- [104] Roussel R, Riveline J-P, Vicaute E, de Pouvourville G, Detournay B, Emery C, et al. Important drop in the rate of acute diabetes complications in people with type 1 or type 2 diabetes after initiation of flash glucose monitoring in france: the RELIEF study. *Diabetes Care* 2021;44:1368–76. <https://doi.org/10.2337/dc20-1690>.
- [105] Bergenstal RM, Kerr MSD, Roberts GJ, Souto D, Nabutovsky Y, Hirsch IB. Flash CGM is associated with reduced diabetes events and hospitalizations in insulin-treated type 2 diabetes. *J Endocr Soc* 2021;5:bvab013. <https://doi.org/10.1210/jendso/bvab013>.
- [106] Riveline J-P, Roussel R, Vicaute E, de Pouvourville G, Detournay B, Emery C, et al. Reduced rate of acute diabetes events with flash glucose monitoring is sustained for two-years after initiation: extended outcomes from the RELIEF study. *Diabetes Technology Ther* 2022;24:611–8. <https://doi.org/10.1089/dia.2022.0085>.
- [107] Nathanson D, Eeg-Olofsson K, Spelman T, Bülow E, Kyhlstedt M, Levrat-Guillen F, et al. Intermittently scanned continuous glucose monitoring compared with blood glucose monitoring is associated with lower HbA1c and a reduced risk of hospitalisation for diabetes-related complications in adults with type 2 diabetes on insulin therapies. *Diabetologia* 2025;68:41–51. <https://doi.org/10.1007/s00125-024-06289-z>.

- [108] Davies MJ, Arora VR, Collins BS, Gabbay RA, Green J, Maruthur NM, et al. Management of hyperglycaemia in type 2 diabetes, 2022. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia* 2022;65:1925–66. <https://doi.org/10.1007/s00125-022-05787-2>.
- [109] NICE. National Institute for Health and Care Excellence (NICE). Type 2 diabetes in adults: management. NICE guideline [NG28] 2022.
- [110] Jendle J, Eeg-Olofsson K, Svensson A-M, Franzen S, Lamotte M, Levrat-Guillen F. Cost-effectiveness of the freestyle libre® system versus blood glucose self-monitoring in individuals with type 2 diabetes on insulin treatment in Sweden. *Diabetes Ther* 2021;12:3137–52. <https://doi.org/10.1007/s13300-021-01172-1>.
- [111] Ajjan R, Bilir SP, Hellmund R, Souto D. Cost-effectiveness analysis of flash glucose monitoring system for people with type 2 diabetes receiving intensive insulin treatment. *Diabetes Ther* 2022;13:1933–45. <https://doi.org/10.1007/s13300-022-01325-w>.
- [112] Bouillet B, Tschertcher P, Vaillard L, Nonciaux C, Hourdain P, Ravier A, et al. Frequent and severe hypoglycaemia detected with continuous glucose monitoring in older institutionalised patients with diabetes. *Age Ageing* 2021;50:2088–93. <https://doi.org/10.1093/ageing/afab128>.
- [113] Martens T, Beck RW, Bailey R, Ruedy KJ, Calhoun P, Peters AL, et al. Effect of continuous glucose monitoring on glycemic control in patients with type 2 diabetes treated with basal insulin. *JAMA* 2021;325:2262–72. <https://doi.org/10.1001/jama.2021.7444>.
- [114] Ehrhardt NM, Chellappa M, Walker MS, Fonda SJ, Vigersky RA. The effect of real-time continuous glucose monitoring on glycemic control in patients with type 2 diabetes mellitus. *J Diabetes Sci Technol* 2011;5:668–75. <https://doi.org/10.1177/193229681100500320>.
- [115] Vigersky RA, Fonda SJ, Chellappa M, Walker MS, Ehrhardt NM. Short- and long-term effects of real-time continuous glucose monitoring in patients with type 2 diabetes. *Diabetes Care* 2012;35:32–8. <https://doi.org/10.2337/dc11-1438>.
- [116] Carlson AL, Daniel TD, DeSantis A, Jabbour S, French EK, Kruger D, et al. Flash glucose monitoring in type 2 diabetes managed with basal insulin in the USA: a retrospective real-world chart review study and meta-analysis. *BMJ Open Diabetes Res Care* 2022;10:e002590. <https://doi.org/10.1136/bmjdr-2021-002590>.
- [117] Elliott T, Beca S, Beharry R, Tsoukas MA, Zarruk A, Abitbol A. The impact of flash glucose monitoring on glycosylated hemoglobin in type 2 diabetes managed with basal insulin in Canada: a retrospective real-world chart review study. *Diab Vasc Dis Res* 2021;18:14791641211021374. <https://doi.org/10.1177/14791641211021374>.
- [118] Guerci B, Roussel R, Levrat-Guillen F, Detournay B, Vicaud E, de Pourville G, et al. Important decrease in hospitalizations for acute diabetes events following FreeStyle Libre® system initiation in people with type 2 diabetes on basal insulin therapy in France. *Diabetes Technol Ther* 2023;25:20–30. <https://doi.org/10.1089/dia.2022.0271>.
- [119] Fonda SJ, Graham C, Munakata J, Powers JM, Price D, Vigersky RA. The cost-effectiveness of real-time continuous glucose monitoring (RT-CGM) in type 2 diabetes. *J Diabetes Sci Technol* 2016;10:898–904. <https://doi.org/10.1177/1932296816628547>.
- [120] Wright EE, Kerr MSD, Reyes IJ, Nabutovsky Y, Miller E. Use of flash continuous glucose monitoring is associated with A1C reduction in people with type 2 diabetes treated with basal insulin or noninsulin therapy. *Diabetes Spectr* 2021;34:184–9. <https://doi.org/10.2337/ds20-0069>.
- [121] Wada E, Onoue T, Kobayashi T, Handa T, Hayase A, Ito M, et al. Flash glucose monitoring helps achieve better glycemic control than conventional self-monitoring of blood glucose in non-insulin-treated type 2 diabetes: a randomized controlled trial. *BMJ Open Diabetes Res Care* 2020;8:e001115. <https://doi.org/10.1136/bmjdr-2019-001115>.
- [122] Grace T, Salyer J. Use of real-time continuous glucose monitoring improves glycemic control and other clinical outcomes in type 2 diabetes patients treated with less intensive therapy. *Diabetes Technol Ther* 2022;24:26–31. <https://doi.org/10.1089/dia.2021.0212>.
- [123] Cox DJ, Banton T, Moncrief M, Conaway M, Diamond A, McCall AL. Minimizing glucose excursions (GEM) with continuous glucose monitoring in type 2 diabetes: a randomized clinical trial. *J Endocr Soc* 2020;4:bvaa118. <https://doi.org/10.1210/jendso/bvaa118>.
- [124] Aronson R, Brown RE, Chu L, Bajaj HS, Khandwala H, Abitbol A, et al. Impact of flash glucose monitoring in people with type 2 Diabetes Inadequately controlled with non-insulin Antihyperglycaemic Therapy (IMMEDIATE): a randomized controlled trial. *Diabetes Obes Metab* 2023;25:1024–31. <https://doi.org/10.1111/dom.14949>.
- [125] Miller E, Kerr MSD, Roberts GJ, Nabutovsky Y, Wright E. Flash CGM associated with event reduction in nonintensive diabetes therapy. *Am J Manag Care* 2021;27:e372–7. <https://doi.org/10.37765/ajmc.2021.88780>.
- [126] Tang X, Li S, Wang M, Yin Q, Mu P, et al. Glycemic variability evaluated by continuous glucose monitoring system is associated with the 10-y cardiovascular risk of diabetic patients with well-controlled HbA1c. *Clin Chim Acta* 2016;461:146–50. <https://doi.org/10.1016/j.cca.2016.08.004>.
- [127] Echouffo-Tcheugui JB, Zhao S, Brock G, Matsouka RA, Kline D, Joseph JJ. Visit-to-visit glycemic variability and risks of cardiovascular events and all-cause mortality: the ALLHAT study. *Diabetes Care* 2019;42:486–93. <https://doi.org/10.2337/dc18-1430>.
- [128] Nyiraty S, Pesei F, Orosz A, Coluzzi S, Vági OE, Lengyel C, et al. Cardiovascular autonomic neuropathy and glucose variability in patients with type 1 diabetes: is there an association? *Front Endocrinol* 2018;9:174. <https://doi.org/10.3389/fendo.2018.00174>.
- [129] Wakasugi S, Mita T, Katakami N, Okada Y, Yoshii H, Osonoi T, et al. Associations between continuous glucose monitoring-derived metrics and diabetic retinopathy and albuminuria in patients with type 2 diabetes. *BMJ Open Diabetes Res Care* 2021;9:e001923. <https://doi.org/10.1136/bmjdr-2020-001923>.
- [130] Farhan HA, Bukhari K, Grewal N, Devarasetty S, Munir K. Use of continuous glucose monitor as a motivational device for lifestyle modifications to improve glycaemic control in patients with type 2 diabetes treated with non-insulin therapies. *BMJ Case Rep* 2022;15:e248579. <https://doi.org/10.1136/bcr-2021-248579>.
- [131] Schubert-Olesen O, Kröger J, Siegmund T, Thurm U, Halle M. Continuous glucose monitoring and physical activity. *Int J Environ Res Public Health* 2022;19:12296. <https://doi.org/10.3390/ijerph191912296>.
- [132] Palmer SC, Mavridis D, Nicolucci A, Johnson DW, Tonelli M, Craig JC, et al. Comparison of clinical outcomes and adverse events associated with glucose-lowering drugs in patients with type 2 diabetes: a meta-analysis. *JAMA* 2016;316:313–24. <https://doi.org/10.1001/jama.2016.9400>.
- [133] Gehlert RR, Dogbey GY, Schwartz FL, Marling CR, Shubbrook JH. Hypoglycemia in type 2 diabetes - more common than you think. *J Diabetes Sci Technol* 2015;9:999–1005. <https://doi.org/10.1177/1932296815581052>.
- [134] Heise T, DeVries JH, Urva S, Li J, Pratt EJ, Thomas MK, et al. Tirzepatide reduces appetite, energy intake, and fat mass in people with type 2 diabetes. *Diabetes Care* 2023;46:998–1004. <https://doi.org/10.2337/dc22-1710>.
- [135] Wharton S, Batterham RL, Bhatta M, Buscemi S, Christensen LN, Frias JP, et al. Two-year effect of semaglutide 2.4 mg on control of eating in adults with overweight/obesity: STEP 5. *Obesity* 2023;31:703–15. <https://doi.org/10.1002/oby.23673>.
- [136] Foreman YD, Brouwers MCGJ, van der Kallen CJH, Pagen DME, van Greevenbroek MMJ, Henry RMA, et al. Glucose variability assessed with continuous glucose monitoring: reliability, reference values, and correlations with established glycemic indices—the maastricht study. *Diabetes Technol Ther* 2020;22:395–403. <https://doi.org/10.1089/dia.2019.0385>.
- [137] Holman RR, Paul SK, Bethel MA, Matthews DR, Neil HAW. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* 2008;359:1577–89. <https://doi.org/10.1056/nejmoa0806470>.
- [138] Stratton IM, Adler AI, Neil HAW, Matthews DR, Manley SE, Cull CA, et al. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ* 2000;321:405–12. <https://doi.org/10.1136/bmj.321.7258.405>.
- [139] UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *UK Prospective Diabetes Study (UKPDS) Group. Lancet* 1998;352:837–53. [https://doi.org/10.1016/s0140-6736\(98\)07019-6](https://doi.org/10.1016/s0140-6736(98)07019-6).
- [140] Rein M, Ben-Yacov O, Godneva A, Shilo S, Zmora N, Kolobkov D, et al. Effects of personalized diets by prediction of glycemic responses on glycemic control and metabolic health in newly diagnosed T2DM: a randomized dietary intervention pilot trial. *BMC Med* 2022;20:56. <https://doi.org/10.1186/s12916-022-02254-y>.
- [141] Cox DJ, Oser T, Moncrief M, Conaway M, McCall A. Long-term follow-up of a randomized clinical trial comparing glycemic excursion minimization (GEM) to weight loss (WL) in the management of type 2 diabetes. *BMJ Open Diabetes Res Care* 2021;9:e002403. <https://doi.org/10.1136/bmjdr-2021-002403>.
- [142] Cox DJ, Banton T, Moncrief M, Conaway M, Diamond A, Holmes V, et al. Glycemic excursion minimization in the management of type 2 diabetes: a novel intervention tested in a randomized clinical trial. *BMJ Open Diabetes Res Care* 2020;8:e001795. <https://doi.org/10.1136/bmjdr-2020-001795>.
- [143] Martens TW, Willis HJ, Kruger DF, Karlioglu-French E, Steenkamp DW. A randomized controlled trial using continuous glucose monitoring (CGM) to guide food choices and diabetes self-care in people with type 2 diabetes not taking insulin. *Diabetes* 2024;73. <https://doi.org/10.2337/db24-356-or>.
- [144] Davies MJ, Arora VR, Collins BS, Gabbay RA, Green J, Maruthur NM, et al. Management of hyperglycemia in type 2 diabetes, 2022. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 2022;45:2753–86. <https://doi.org/10.2337/dci22-0034>.
- [145] Ceriello A, Praticchizzo F, Phillip M, Hirsch IB, Mathieu C, Battelino T. Glycaemic management in diabetes: old and new approaches. *Lancet Diabetes Endocrinol* 2022;10:75–84. [https://doi.org/10.1016/s2213-8587\(21\)00245-x](https://doi.org/10.1016/s2213-8587(21)00245-x).
- [146] Raj R, Mishra R, Jha N, Joshi V, Correa R, Kern PA. Time in range, as measured by continuous glucose monitor, as a predictor of microvascular complications in type 2 diabetes: a systematic review. *BMJ Open Diabetes Res Care* 2022;10:e002573. <https://doi.org/10.1136/bmjdr-2021-002573>.
- [147] Lu J, Ma X, Zhou J, Zhang L, Mo Y, Ying L, et al. Association of time in range, as assessed by continuous glucose monitoring, with diabetic retinopathy in type 2 diabetes. *Diabetes Care* 2018;41:2370–6. <https://doi.org/10.2337/dc18-1131>.
- [148] Wang Y, Lu J, Shen Y, Ni J, Zhang L, Lu W, et al. Comparison of glucose time in range and area under curve in relation to risk of diabetic retinopathy in type 2 diabetes patients. *J Diabetes Investig* 2022;13:1543–50. <https://doi.org/10.1111/jdi.13811>.
- [149] Yang J, Yang X, Zhao D, Wang X, Wei W, Yuan H. Association of time in range, as assessed by continuous glucose monitoring, with painful diabetic polyneuropathy. *J Diabetes Investig* 2021;12:828–36. <https://doi.org/10.1111/jdi.13394>.
- [150] Li F, Zhang Y, Li H, Lu J, Jiang L, Vigersky RA, et al. TIR generated by continuous glucose monitoring is associated with peripheral nerve function in type 2 diabetes. *Diabetes Res Clin Pract* 2020;166:108289. <https://doi.org/10.1016/j.diabres.2020.108289>.

- [151] Gad H, Elgassim E, Mohammed I, Alhaddad AY, Aly HAHZ, Cabibihan JJ, et al. Continuous glucose monitoring reveals a novel association between duration and severity of hypoglycemia, and small nerve fiber injury in patients with diabetes. *Endocr Connect* 2022;11:e220352. <https://doi.org/10.1530/ec-22-0352>.
- [152] Gad H, Elgassim E, Mohammed I, Alhaddad AY, Aly HAHZ, Cabibihan J-J, et al. Cardiovascular autonomic neuropathy is associated with increased glycemic variability driven by hyperglycemia rather than hypoglycemia in patients with diabetes. *Diabetes Res Clin Pract* 2023;200:110670. <https://doi.org/10.1016/j.diabres.2023.110670>.
- [153] Lu J, Ma X, Shen Y, Wu Q, Wang R, Zhang L, et al. Time in range is associated with carotid intima-media thickness in type 2 diabetes. *Diabetes Technol Ther* 2020;22:72–8. <https://doi.org/10.1089/dia.2019.0251>.
- [154] Zhou H, Wang W, Shen Q, Feng Z, Zhang Z, Lei H, et al. Time in range, assessed with continuous glucose monitoring, is associated with brachial-ankle pulse wave velocity in type 2 diabetes: a retrospective single-center analysis. *Front Endocrinol* 2022;13:1014568. <https://doi.org/10.3389/fendo.2022.1014568>.
- [155] Lu J, Wang C, Shen Y, Chen L, Zhang L, Cai J, et al. Time in range in relation to all-cause and cardiovascular mortality in patients with type 2 diabetes: a prospective cohort study. *Diabetes Care* 2021;44:549–55. <https://doi.org/10.2337/dc20-1862>.
- [156] Li J, Li Y, Ma W, Liu Y, Yin X, Xie C, et al. Association of time in range levels with lower extremity arterial disease in patients with type 2 diabetes. *Diabetes Metab Syndr* 2020;14:2081–5. <https://doi.org/10.1016/j.dsx.2020.09.028>.
- [157] Xie P, Deng B, Zhang X, Li Y, Du C, Rui S, et al. Time in range in relation to amputation and all-cause mortality in hospitalised patients with diabetic foot ulcers. *Diabetes Metabol Res Rev* 2022;38:e3498.
- [158] Mayeda L, Katz R, Ahmad I, Bansal N, Batacchi Z, Hirsch IB, et al. Glucose time in range and peripheral neuropathy in type 2 diabetes mellitus and chronic kidney disease. *BMJ Open Diabetes Res Care* 2020;8:e000991. <https://doi.org/10.1136/bmjopen-2019-000991>.
- [159] Guo Q-Y, Lu B, Guo Z-H, Feng Z-Q, Yuan Y-Y, Jin X-G, et al. Continuous glucose monitoring defined time-in-range is associated with sudomotor dysfunction in type 2 diabetes. *World J Diabetes* 2020;11:489–500. <https://doi.org/10.4239/wjd.v11.i11.489>.
- [160] Kim MY, Kim G, Park JY, Choi MS, Jun JE, Lee Y-B, et al. The association between continuous glucose monitoring-derived metrics and cardiovascular autonomic neuropathy in outpatients with type 2 diabetes. *Diabetes Technol Ther* 2021;23:434–42. <https://doi.org/10.1089/dia.2020.0599>.
- [161] Guo Q, Zang P, Xu S, Song W, Zhang Z, Liu C, et al. Time in range, as a novel metric of glycemic control, is reversely associated with presence of diabetic cardiovascular autonomic neuropathy independent of HbA1c in chinese type 2 diabetes. *J Diabetes Res* 2020;2020:5817074. <https://doi.org/10.1155/2020/5817074>.
- [162] Yoo JH, Choi MS, Ahn J, Park SW, Kim Y, Hur KY, et al. Association between continuous glucose monitoring-derived time in range, other core metrics, and albuminuria in type 2 diabetes. *Diabetes Technol Ther* 2020;22:768–76. <https://doi.org/10.1089/dia.2019.0499>.
- [163] Commission E. Benchmarking deployment of eHealth among. *General Practitioners* 2018.
- [164] Danne T, Limbert C, Domingo MP, Prato SD, Renard E, Choudhary P, et al. Telemonitoring, telemedicine and time in range during the pandemic: paradigm change for diabetes risk management in the post-COVID future. *Diabetes Ther* 2021;12:2289–310. <https://doi.org/10.1007/s13300-021-01114-x>.
- [165] Silva-Tinoco R, de la Torre-Saldaña V. The imperious need for telemedicine for the care of diabetes during the COVID-19 pandemic. a comprehensive approach study. *Gac Med Mex* 2021;157:309–12. <https://doi.org/10.24875/gmm.m21000563>.
- [166] Pogorzelska K, Marcinowicz L, Chlabicz S. A qualitative study of primary care physicians' experiences with telemedicine during the COVID-19 pandemic in North-Eastern Poland. *Int J Environ Res Public Health* 2023;20:1963. <https://doi.org/10.3390/ijerph20031963>.
- [167] Poitras M-E, Poirier M-D, Couturier Y, Vaillancourt VT, Cormier C, Gauthier G, et al. Chronic conditions patient's perception of post-COVID-19 pandemic teleconsulting continuation in primary care clinics: a qualitative descriptive study. *BMJ Open* 2022;12:e066871. <https://doi.org/10.1136/bmjopen-2022-066871>.
- [168] Mayberry LS, Guy C, Hendrickson CD, McCoy AB, Elasy T. Rates and correlates of uptake of continuous glucose monitors among adults with type 2 diabetes in primary care and endocrinology settings. *J Gen Intern Med* 2023;38:2546–52. <https://doi.org/10.1007/s11606-023-08222-3>.
- [169] Kieu A, King J, Govender RD, Östlundh L. The benefits of utilizing continuous glucose monitoring of diabetes mellitus in primary care: a systematic review. *J Diabetes Sci Technol* 2023;17:762–74. <https://doi.org/10.1177/19322968211070855>.
- [170] Edelman SV, Cavaola TS, Boeder S, Pettus J. Utilizing continuous glucose monitoring in primary care practice: what the numbers mean. *Prim Care Diabetes* 2021;15:199–207. <https://doi.org/10.1016/j.pcd.2020.10.013>.
- [171] Filippi MK, Lanigan AM, Oser SM, Alai J, Brooks-Greisen A, Oser TK. A team-based training for continuous glucose monitoring in diabetes care: mixed methods pilot implementation study in primary care practices. *JMIR Form Res* 2023;7:e45189. <https://doi.org/10.2196/45189>.
- [172] Oser TK, Hall TL, Dickinson LM, Callen E, Carroll JK, Nease DE, et al. Continuous glucose monitoring in primary care: understanding and supporting clinicians' use to enhance diabetes care. *Ann Fam Med* 2022;20:541–7. <https://doi.org/10.1370/afm.2876>.
- [173] Martens TW. Continuous glucose monitoring in primary care – are we there? *Curr Opin Endocrinol Diabetes Obes* 2022;29(1):10–6. <https://doi.org/10.1097/med.0000000000000689>.
- [174] Allen MR, Webb S, Mandvi A, Frieden M, Tai-Seale M, Kallenberg G. Navigating the doctor-patient-AI relationship - a mixed-methods study of physician attitudes toward artificial intelligence in primary care. *BMC Prim Care* 2024;25:42. <https://doi.org/10.1186/s12875-024-02282-y>.
- [175] Morton JI, Marquina C, Magliano DJ, Shaw JE, Ademi Z. Targeting diabetes prevention to more disadvantaged groups improves cost-effectiveness: implications of inequality in type 2 diabetes from theoretical interventions. *Value Health* 2023;26:974–83. <https://doi.org/10.1016/j.jval.2023.02.003>.
- [176] Bilir SP, Hellmund R, Wehler E, Li H, Munakata J, Lamotte M. The Cost-effectiveness of a flash glucose monitoring system for management of patients with type 2 diabetes receiving intensive insulin treatment in Sweden. *Eur Endocrinol* 2018;14:80–5. <https://doi.org/10.17925/ee.2018.14.2.80>.
- [177] Isitt JJ, Roze S, Sharland H, Cogswell G, Alshannaq H, Norman GJ, et al. Cost-effectiveness of a real-time continuous glucose monitoring system versus self-monitoring of blood glucose in people with type 2 diabetes on insulin therapy in the UK. *Diabetes Ther* 2022;13:1875–90. <https://doi.org/10.1007/s13300-022-01324-x>.
- [178] Isaacson B, Kaufusi S, Sorensen J, Joy E, Jones C, Ingram V, et al. Demonstrating the clinical impact of continuous glucose monitoring within an integrated healthcare delivery system. *J Diabetes Sci Technol* 2022;16:383–9. <https://doi.org/10.1177/1932296820955228>.

Glossary

ADE: Acute diabetes event
 BMI: Artificial intelligence
 CAN: Cardiac autonomic neuropathy
 CGM: Continuous glucose monitoring
 CKD: Chronic kidney disease
 DPN: Diabetic peripheral neuropathy
 DSMES: Diabetes self-management education and support
 FPG: Fasting plasma glucose
 GLP-1 RA: Glucagon-like peptide 1 receptor agonists
 GMI: Glucose management indicator
 GV: Glycemic variability
 HCP: Healthcare professional
 ICER: Incremental cost-effectiveness ratio
 IDF-Eu: International Diabetes Federation Europe
 IGT: Impaired glucose tolerance
 OGTT: Oral glucose tolerance test
 PCDE: Primary Care Diabetes Europe
 PCP: Primary care physician
 QALE: Quality adjusted life expectancy
 QALY: Quality adjusted life year
 RCT: Randomized controlled trial
 SDoH: Social determinants of health
 SMBG: Selfmonitored blood glucose
 T1D: Type 1 diabetes
 TAR: Time above range
 TIR: Time in range
 T1TR: Time in tight range
 TRE: Time restricted eating