

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 35, 2025, 2





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 35, 2025, 2

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 35, leto 2025 številka 2

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Založnika/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran /Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 6. 12. 2025.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.

Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>

VSEBINA / *INDICE GENERALE* / CONTENTSSREDOZEMSKJE HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
*MEDITERRANEAN SHARKS AND RAYS***Hakan KABASAKAL**

Analysis of Confirmed Shark Attacks
in the Eastern Mediterranean Sea and
the Sea of Marmara (1827–2025) 169
Analiza potrjenih napadov morskih psov
v vzhodnem Sredozemskem morju
in Marmarskem morju (1827–2025)

Alen SOLDO

Observations of a Juvenile Basking Shark
Cetorhinus maximus in the Adriatic Sea
Support the Hypothesis of a Distinct
Mediterranean Population 187
Opazovanja mladega morskega psa
orjaka (Cetorhinus maximus)
v Jadranskem morju podpirajo
hipotezo o ločeni sredozemski
populaciji

**Jacopo BERNARDI, Pero UGARKOVIĆ &
Ilija ČETKOVIĆ**

An Unusual Encounter with a Juvenile Kitefin
Shark, *Dalatias licha*, in Shallow Coastal
Waters of Montenegro (Adriatic Sea) 197
Nenavadno srečanje z mladim primerkom
klinoplavutega morskega psa, Dalatias licha,
v plitvih obalnih vodah Črne gore
(Jadransko morje)

**Cemal TURAN, Alen SOLDO, Servet A. DOĞDU,
Funda TURAN, Aysegül ERGENLER & Ali UYAN**
Identification of a Potential Nursery Ground
of the Spiny Butterfly Ray, *Gymnura altavela*,
in the Northeastern Mediterranean Sea, Türkiye 205
Identifikacija potencialnih jaslčic za
metuljastega skata, Gymnura
altavela, v severovzhodnem
Sredozemskem morju,
Turčija

**Alen SOLDO, Eleonora DE SABATA &
Simona CLO**

Recent Records of Critically Endangered
Common Guitarfish *Rhinobatos rhinobatos*
(Linnaeus, 1758) in the Northern
Mediterranean 213
Nedavni zapiski o pojavljanju kritično
ogroženega navadnega goslaša
Rhinobatos rhinobatos
(Linnaeus, 1758) v
severnem Sredozemlju

Hakan KABASAKAL & F. Saadet KARAKULAK

Demersal Elasmobranchs of the Sea of Marmara:
Updated Inventory, Taxonomic Issues and
Environmental Implications 221
Pridnene hrustančnice v Marmarskem morju:
posodobljena inventarizacija, taksonomska
vprišanja in okoljske posledice

IHTIOLOGIJA

ITTIOLOGIA

ICHTHYOLOGY

**Jamila RIZGALLA, Amani FITORI, Francesco
TIRALONGO & Abdalrh BEN ABDALAH**

First Records of Blennies (Suborder
Blennioidea) off the Coast of Libya 245
Prvi zapisi o pojavljanju babic
(podred Blennioidea) ob
libijski obali

**Farid HEMIDA, Sara LADOUL, Christian
REYNAUD & Christian CAPAPÉ**

Confirmed Occurrence of the Mediterranean
Spearfish *Tetrapturus belone* (Osteichthyes:
Istiophoridae) from the Algerian Coast
(Southwestern Mediterranean Sea) 257
Potrjeno pojavljanje sredozemske
jadrovnice, Tetrapturus belone
(Osteichthyes: Istiophoridae) iz
alžirske obale (jugovzhodno
Sredozemsko morje)

Lana KHREMA, Adib SAAD & Christian CAPAPÉ

First Substantiated Record of Blackfish
Centrolophus niger (Centrolophidae)
from the Syrian Coast (Eastern
Mediterranean Sea) 263
Prvi potrjen zapis črnuha, Centrolophus niger
(*Centrolophidae*), s sirske obale
(vzhodno Sredozemsko morje)

Amir IBRAHIM, Chirine HUSSEIN, Firas ALSHAWY & Alaa ALCHEIKH AHMAD

Marine Fishes (Teleostei / Osteichthyes)
of Syria (Eastern Mediterranean):
an Updated Checklist 269
Morske ribe (Teleostei / Osteichthyes)
Sirije (vzhodno Sredozemsko morje):
posodobljeni seznam vrst

BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
BIOTIC GLOBALIZATION

Paola LEOTTA, Rocco CICCARELLA, Ruben GARRANO, Enrico LA SPINA, Daniele TIBULLO & Francesco TIRALONGO

Cassiopea andromeda at the Southernmost
Tip of Italy: a Recent Arrival or an
Overlooked Resident? 287
Tujerodni klobučnjak Cassiopea andromeda
na najjužnejši konici Italije: nedavni
prišlek ali spregledan prebivalec?

Mourad CHÉRIF, Rimel BENMESSAOUD, Sihem RAFRAFI-NOUIRA, Mohamed Nouri AYADI & Christian CAPAPÉ

First Substantiated Record of the Golden-Banded
Goatfish *Upeneus moluccensis* (Osteichthyes:
Mullidae) from the Coast of Tunisia
(Central Mediterranean Sea) 295
Prvi potrjen zapis o zlatoprogem bradaču
Upeneus moluccensis (Osteichthyes: Mullidae)
z obale Tunizije (osrednje Sredozemsko morje)

Nikola DJORDJEVIĆ, Slavica PETOVIĆ, Ilija ČETKOVIĆ, Borut MAVRIČ & Lovrenc LIPEJ

New Record of the Long-Jawed Squirrelfish,
Holocentrus adscensionis (Osbeck, 1765),
in the Adriatic Sea 303
Novi zapis o pojavljanju veveričjaka vrste
Holocentrus adscensionis (Osbeck, 1765)
v Jadranskem morju

MORSKA FAVNA
FAUNA MARINA
MARINE FAUNA

Nicola BETTOSO, Lisa FARESI, Saul CIRIACO, Marco FANTIN & Marco SEGARICH

New Findings of the Cup-Shaped Demosponge
Calyx nicaeensis (Risso, 1826) on the Rocky
Outcrops in the Gulf of Trieste
(Northern Adriatic Sea) 311
Nove najdbe morskega keliha Calyx nicaeensis
(*Risso, 1826*) *na skalnih osamelcih v Tržaškem*
zalivu (severno Jadransko morje)

Aleksandra V. BORODINA & Yuri O. VELIAEV

Features of Lipid Accumulation in Striped Venus
Clam *Chamelea gallina* in the Sublittoral
Zone of the Crimean Coast (Black Sea) 319
Značilnosti kopičenja lipidov v navadni
venerici (Chamelea gallina) v obrežnem
pasu krimske obale (Črno morje)

Okan AKYOL & Oğuzhan TAKICAK

Recent Observations on *Monachus monachus*
(Phocidae) at Sea-Cage Fish Farms in Izmir
(Turkish Aegean Sea) 329
Nedavna opažanja primerkov Monachus monachus
(*Phocidae*) *v ribogojnicah z morskimi kletkami v*
Izmirju (turško Egejsko morje)

FLORA
FLORA
FLORA

Amelio PEZZETTA & Marco PAOLUCCI

La Flora di Palena (Parco Nazionale della
Majella): Aggiornamento Floristico 337
Flora Palene (Nacionalni park Majella):
floristična posodobitev

DELO NAŠIH ZAVODOV IN DRUŠTEV
ATTIVITÀ DEI NOSTRI ISTITUTI E SOCIETÀ
ACTIVITIES BY OUR INSTITUTIONS AND
ASSOCIATIONS

Bojana LIPEJ

Reviving Landscapes, Connecting Species:
Lessons from the ReCo Project 383

IN MEMORIAM

Tom TURK

V spomin Marjanu Richterju (1935–2025) 389

Kazalo k slikam na ovitku 391

Index to images on the cover 391

received: 2025-05-16

DOI 10.19233/ASHN.2025.23

IDENTIFICATION OF A POTENTIAL NURSERY GROUND OF THE SPINY BUTTERFLY RAY, *GYMNURA ALTAVELA*, IN THE NORTHEASTERN MEDITERRANEAN SEA, TÜRKİYE

Cemal TURAN

Iskenderun Technical University, Faculty of Marine Sciences and Technology, Molecular Ecology and Fisheries Genetics Laboratory,
31220, Iskenderun, Hatay, Türkiye
Nature and Science Society, Modernevler Neighborhood 303 St. No:9/1, Iskenderun, Hatay
e-mail: cemal.turan@iste.edu.tr

Alen SOLDÓ

Department of Marine Studies, University of Split, Split 21000, Croatia

Servet A. DOĞDU

Nature and Science Society, Modernevler Neighborhood 303 St. No:9/1, Iskenderun, Hatay
Iskenderun Technical University, Maritime Technology Vocational School of Higher Education, Underwater Technologies, 31200
Iskenderun, Hatay, Türkiye

Funda TURAN, Ayşegül ERGENLER & Ali UYAN

Iskenderun Technical University, Faculty of Marine Sciences and Technology, Molecular Ecology and Fisheries Genetics Laboratory,
31220, Iskenderun, Hatay, Türkiye
Nature and Science Society, Modernevler Neighborhood 303 St. No:9/1, Iskenderun, Hatay

ABSTRACT

*This study reports the first identified potential nursery area for the critically endangered *Gymnura altavela* off the coast of Samandağ in the northeastern Mediterranean. Two divers conducted seasonal surveys across a total area of 750 m² at three depth intervals (5–10 m, 11–20 m, 21–30 m) using the standard underwater visual census method. Additionally, bycatch data from nearby commercial trawling operations were collected. The highest individual density recorded was 0.051 n/m² in autumn, the lowest 0.02 n/m² in summer. The findings suggest that the area functions as a nursery habitat for the *G. altavela* population.*

Key words: *Gymnura altavela*, nursery site, breeding area, Samandağ coasts, northeastern Mediterranean

IDENTIFICAZIONE DI UN POTENZIALE LUOGO DI RIPRODUZIONE DI *GYMNURA ALTAVELA* NEL MAR MEDITERRANEO NORD-ORIENTALE, TURCHIA

SINTESI

*Questo studio riporta la prima area potenziale identificata come zona di riproduzione per la specie *Gymnura altavela*, gravemente minacciata di estinzione, al largo della costa di Samandağ, nel Mediterraneo nord-orientale. Due subacquei hanno condotto indagini stagionali su un'area totale di 750 m² a tre intervalli di profondità (5-10 m, 11-20 m, 21-30 m) utilizzando il metodo standard di censimento visivo subacqueo. Sono stati inoltre raccolti dati sulle catture accessorie delle vicine attività di pesca commerciale con reti a strascico. La densità individuale più alta registrata è stata di 0,051 n/m² in autunno, la più bassa di 0,02 n/m² in estate. I risultati suggeriscono che l'area funge da habitat di riproduzione per la popolazione di *G. altavela*.*

Parole chiave: *Gymnura altavela*, sito di nursery, area di riproduzione, coste di Samandağ, Mediterraneo nord-orientale

INTRODUCTION

Butterfly rays (Elasmobranchii: Myliobatiformes: Gymnuridae) are demersal batoids with a global distribution in the inshore coastal waters of tropical and temperate seas, typically found over soft sandy or muddy bottoms but also occurring over rocky reefs (Compagno & Last, 1999; McEachran & Carvalho, 2002; Ebert & Dando, 2020; Barone *et al.*, 2022).

The family is represented by a single genus, *Gymnura*, which comprises 16 species worldwide. *Gymnura altavela* (Linnaeus, 1758) is the only member of this family present in the Mediterranean Sea (Ebert & Dando, 2020; Barone *et al.*, 2022). The spiny butterfly ray *G. altavela* is the largest member of the Gymnuridae family, reaching a maximum disc width (DW) of approximately 2600 mm (Ebert & Dando, 2020). It is widely distributed in the western and eastern Atlantic Ocean, the Mediterranean Sea, and the Black Sea (Weigmann, 2016), inhabiting shallow inshore waters in the bathymetric range from 10 to 69 m (Ebert & Stehmann, 2013). Its diet consists of all kinds of benthic animals, but primarily fish and cephalopods (Bauchot, 1987). The species is aplacental viviparous, with an annual reproductive cycle that includes a six-month gestation period and results in a litter of 2–7 pups (McEachran & Capapé 1984; Bradai *et al.*, 2012). Although generally harmless to humans, *G. altavela* possesses a caudal spine capable of inflicting a painful wound if stepped on. It is also listed as a game fish in some regions (Conrath & Scarbrough, 2009).

G. altavela is caught as bycatch in trawl, trammel net, and longline fisheries (Bauchot 1987; Yağlıoğlu *et al.*, 2015). The wings are marketed fresh, chilled, or frozen in Sicily and Morocco, rarely elsewhere (Bauchot, 1987). The IUCN Red List of Threatened Species classifies the species as Endangered globally, and Critically Endangered in the Mediterranean, due to a suspected population decline exceeding 80% over the past three generations (Dulvy *et al.*, 2021). This drastic decline, documented in catch data (Abdul Malak *et al.*, 2011; Özbek *et al.*, 2016), is driven by intense fishing pressure and the species' slow reproductive rate.

The spiny butterfly ray was historically moderately abundant and commonly caught throughout the Mediterranean region. For much of the last century, up until the 1980s, it frequently appeared in the catches of demersal trawl and set net fisheries, particularly along the southern shores. In areas such as the Sicilian Channel, where it was once regularly captured, the species has now become very rare or is absent from local catch records (Bauchot, 1987). Its decline is further evidenced by its absence from the International Trawl Survey in the Mediterranean (MEDITS) records since 1994. The occasional contemporary reports of

its presence are typically based on the incidental capture of individual specimens in demersal fisheries. An exception to this overall decline is the Levant coast, where the species remains relatively abundant and is regularly caught using bottom trawls, fixed nets, and longlines (Dulvy *et al.*, 2021).

Identification of critical habitats is an essential and well-established component of sustainable resource management (Gonçalves Silva and Vianna, 2018; Ergüden *et al.*, 2025; Turan *et al.*, 2025), and nursery sites are an important category of such habitats (Medeiros *et al.*, 2015; Rangel *et al.*, 2018). They can be used by many species, separated in space and time, and it is important to understand the mechanisms and drivers behind their use (Heithaus, 2007; Rosenfelder *et al.*, 2012; Cengiz *et al.*, 2024; Kabasakal *et al.*, 2024; Turan *et al.*, 2024). Heupel *et al.* (2007) established three key criteria for identifying an elasmobranch nursery habitat: (1) juvenile elasmobranchs are more commonly encountered there than in other areas, (2) they remain in the area for extended periods, and (3) the area is used repeatedly across years. Therefore, the main aim of this study is to evaluate a potential nursery ground



Fig. 1: Map of the study area.

Sl. 1: Zemljevid obravnavanega območja.

for the spiny butterfly ray *G. altavela* in the north-eastern Mediterranean. If confirmed, this site would represent the second identified critical habitat for this species along the Turkish Mediterranean coast (Bilgili & Kabasakal, 2023).

MATERIAL AND METHODS

This study, carried out over a one-year research period within the framework of a project supported by the Turquoise Coast Environment Fund–Turkey (ElasmoKAP-TCEF-2024), utilized a standardized underwater visual census (UVC) method (Whitfield et al., 2007; Murphy et al., 2010; Turan & Doğdu, 2022) to assess the density, abundance, distribution, and reproductive status–based on observed abdominal swellings–of *G. altavela* in the Ray field region of Hatay-Samandağ coast. Two divers monitored three depth ranges: shallow (5–10 m),




Fig. 2: Mean density of *G. altavela* across all depths and seasons within the identified nursery area.

Sl. 2: Povprečna gostota primerkov vrste *G. altavela* v vseh globinah in letnih časih znotraj opredeljenega območja jaslic.




Fig. 3: Dorsal view of egg sacs from pregnant females *G. altavela* (A and B) (Photo: Cemal Turan).

Sl. 3: Pogled na hrbtno stran jajčnih vrečk brejih samic vrste *G. altavela* (A in B) (Foto: Cemal Turan).




Fig. 4: Individuals of *G. altavela* in the investigating area: (A) a pregnant female and (B) an adult male (Photo: Cemal Turan).

Sl. 4: Osebk vrste *G. altavela* na raziskovanem območju: (A) breja samica in (B) odrasel samec (Foto: Cemal Turan).

intermediate (11–20 m), and deep (21–30 m). At each depth range, a 5 × 50 m transect was surveyed, resulting in a total examined area of 750 m² along the Samandağ coast (Fig. 1). Surveys were conducted four times over a year, once per season. The estimate of the mean density (D) on a transect is expressed as:

$$D = \frac{\sum_{i=1} n_i}{a}$$

where n is the number of *G. altavela* individuals observed, and a the census area (Whitfield *et al.*, 2007). In addition to UVC, data on *G. altavela* bycatch from commercial trawling activities near the study area were monitored for one year, and biological data were collected from captured individuals. All captured spiny butterfly rays were handled carefully and retained under appropriate conditions to minimize post-release mortality. All statistical analyses were performed using R-Studio.

RESULTS

G. altavela was observed on the Samandağ coast in all seasons, which suggests its regular occurrence in the area. The mean density across all depths and seasons was 0.038 n/m². Densities varied by depth range, with 0.0 n/m² at 5–10 m, 0.076 n/m² at 10.1–20 m, and 0.052 n/m² at 20.1–30 m. Seasonally, the highest density was recorded in autumn (0.051 n/m²), the lowest in summer (0.02 n/m²) (Fig. 2).

A single visual count in autumn along a 150 m transect in the 11–20 m depth range was documented. There were pregnant females and males were resting on the sandy seabed that a dorsal view of egg sacs from a pregnant females *G. altavela* was given in the Fig. 3A and 3B. Eight pregnant females and three males with long claspers resting on the sandy seabed (Fig. 4A and 4B). Conversely, during a summer survey in the same area and depth range, three non-pregnant females and one male were counted.

During the study, female butterfly ray–pregnant individuals in particular–were observed moving away from the area as divers approached (Fig. 5), a behavior not exhibited by males.

The UVC findings were further corroborated by trawl observations carried out in an adjacent area where commercial fishing is permitted. Trawl bycatch data collected during the local fishing season confirm that *G. altavela*–including mature individuals, juveniles, and newborns–are commonly encountered (Fig. 6) from September to May. The species is less frequently observed during the remaining, rest period.

DISCUSSION

The present study identifies a potential and rare nursery site for *Gymnura altavela* in the Mediterranean. The area is predominantly sandy, with a few small rocky structures, exhibiting similar oceanographic features as other known aggregation sites (Silva & Vianna, 2018; Castro & Meyers, 2022; Espino Ruano *et al.*, 2023).



Fig. 5: Ventral surface of a pregnant female *G. altavela*, showing the egg sac and open cloaca (Photo: Cemal Turan).
Sl. 5: Trebušna površina breje samice vrste *G. altavela*, na kateri je prikazana jajčna vrečka in odprta kloaka (Foto: Cemal Turan).




Fig. 6: Specimens of *G. altavela* on a trawler deck, captured as bycatch near the nursery area: (A) adult and (B) newborn individuals (Photo: Cemal Turan).

Sl. 6: Primerki vrste *G. altavela* na palubi vlečne mreže, ujeti kot prilov v bližini območja jaslic: (A) odrasli in (B) novokoteni primerki (foto: Cemal Turan).

The average water temperature shows considerable seasonal variations: 30–32°C in summer (July), 20–24°C in autumn (November), 16–20°C in winter (February), and 20–22°C in spring (May). Salinity levels fluctuate between 33.6 and 37.1‰. Prior to this study, Bilgili and Kabasakal (2023) identified another aggregation site and potential breeding ground for *G. altavela* in Güllük Bay (SE Aegean Sea). Based on eight years of observation, the authors reported that aggregations of large pregnant females in that area peaked in mid-summer and were subsequently replaced by juveniles in the fall—a pattern consistent with a reproductive aggregation.

Applying the three established criteria for identifying nursery habitats (Heupel *et al.*, 2007) to our findings, we can confirm that the Samandağ Hırlavuk area functions as a nursery ground for *G. altavela*. The highest number of pregnant females was recorded in early autumn, aligning with the known fecundation period (Espino Ruano *et al.*, 2023). Furthermore, year-round observations suggest that this area serves multiple purposes, including feeding, mating, and parturition. The designation of a site as a nursery area is contingent on the presence of newborns, juveniles, and gravid females (Castro, 1993; Heupel *et al.*, 2007). Such areas typically support a higher proportion of juveniles, thereby contributing more significantly to adult recruitment compared to other habitats (Gunter, 1967; Beck *et al.*, 2001). Beck *et al.* (2001) further emphasized the essential role of juvenile survival in driving population growth. Also, in the spawning area they examined, females significantly outnumbered males among the adult population, a pattern consistent with our findings. Capapé *et al.* (2003) suggested that females migrate to shallow coastal waters to find suitable hydrobiological conditions for parturition, thereby minimizing risks to free-swimming neonates from both

intra- and interspecific competition, including cannibalism. This behavior underscores the ecological importance of shallow nursery grounds in enhancing juvenile survival.

Given that *G. altavela* is classified as a critically endangered species in the Mediterranean, the identification and protection of its nursery habitats are crucial for effective conservation and population sustainability. However, research on its reproductive biology and life history remains limited, both in Türkiye and globally (Alkusaairy *et al.*, 2014; Silva & Vianna, 2018; Yeldan *et al.*, 2018; Taylan *et al.*, 2019). The findings of this study contribute valuable insights into the species' reproductive ecology and habitat preferences, emphasizing the need for further research and conservation efforts.

In conclusion, the data presented in this study indicate that a sustainable population of *G. altavela* is established in the Samandağ Hırlavuk area, which serves as a critical nursery habitat. This finding adds Samandağ Hırlavuk to the list of other essential habitats for this species, alongside the Aegean coast of Türkiye (Taylan *et al.*, 2019) and Tunisian and Syrian waters (Alkusaairy *et al.*, 2014; Capapé *et al.*, 1992; El Kamel *et al.*, 2009). Given the species' endangered status, it is imperative to implement conservation strategies that prioritize the protection of these nursery sites to support population recovery and long-term sustainability.

ACKNOWLEDGEMENTS

This study was supported by Turquoise Coast Environment Fund-Turkey with the project “Restoration, Conservation and Monitoring of Shark and Stingray Species Habitats in Hatay Samandağ Hırlavuk Coast-TCEF-2024”, carried out by the Nature and Science Society (www.dogavebilim.com).

IDENTIFIKACIJA POTENCIALNIH JASLIC ZA METULJASTEGA SKATA, *GYMNURA ALTAVELA*, V SEVEROVZHODNEM SREDOZEMSKEM MORJU, TURČIJA

Cemal TURAN

Iskenderun Technical University, Faculty of Marine Sciences and Technology, Molecular Ecology and Fisheries Genetics Laboratory,
31220, Iskenderun, Hatay, Türkiye
Nature and Science Society, Modernevlar Neighborhood 303 St. No:9/1, Iskenderun, Hatay
e-mail: cemal.turan@iste.edu.tr

Alen SOLDÓ

Department of Marine Studies, University of Split, Split 21000, Croatia

Servet A. DOĐDU

Nature and Science Society, Modernevlar Neighborhood 303 St. No:9/1, Iskenderun, Hatay
Iskenderun Technical University, Maritime Technology Vocational School of Higher Education, Underwater Technologies, 31200
Iskenderun, Hatay, Türkiye

Funda TURAN, Ayşegöl ERGENLER & Ali UYAN

Iskenderun Technical University, Faculty of Marine Sciences and Technology, Molecular Ecology and Fisheries Genetics Laboratory,
31220, Iskenderun, Hatay, Türkiye
Nature and Science Society, Modernevlar Neighborhood 303 St. No:9/1, Iskenderun, Hatay

POVZETEK

*Avtorji poročajo o prvem prepoznanem potencialnem območju za razmnoževanje kritično ogrožene vrste *Gymnura altavela* ob obali Samandağ v severovzhodnem Sredozemlju. Dva potapljača sta izvedla sezone raziskave na skupni površini 750 m² v treh globinskih intervalih (5–10 m, 11–20 m, 21–30 m) z uporabo standardne metode podvodnega opazovalnega popisa. Poleg tega so bili zbrani podatki o prilovu iz bližnjih komercialnih vlečnih mrež. Najvišja zabeležena gostota primerkov je bila 0,051 n/m² jeseni, najnižja pa 0,02 n/m² poleti. Ugotovitve kažejo, da območje deluje kot jaslice za populacijo vrste *G. altavela*.*

Ključne besede: metuljasti skat, *Gymnura altavela*, jaslice, razmnoževalno okolje, obale Samandağ, severovzhodno Sredozemlje

REFERENCES

- Abdul Malak, D., S.R. Livingstone, D. Pollard, B.A. Polidoro, A. Cuttelod, M. Bariche, M. Bilecenoglu, K.E. Carpenter, B.B. Collette, P. Francour, M. Goren, M. Hichem Kara, E. Massutí, C. Papaconstantinou & L. Tunesi (2011): Overview of the conservation status of the marine fishes of the Mediterranean Sea. IUCN, Gland and Spain, 61 pp.
- Alkusaïry, H., M. Ali, A. Saad, C. Reynaud & C. Capapé (2014): Maturity, reproductive cycle, and fecundity of spiny butterfly ray, *Gymnura altavela* (Elasmobranchii: Rajiformes: Gymnuridae), from the coast of Syria (eastern Mediterranean). Acta Ichthyol. Piscat., 44, 229-240.
- Barone, M., C. Mazzoldi & F. Serena (2022): Sharks, rays and chimaeras in the Mediterranean and Black Seas – Key to identification. FAO, Rome.
- Bauchot, M.L. (1987): Raies et autres batoides. In: Fischer, W., M.L. Bauchot & M. Schneider (eds.): Fiches FAO d'identification pour les besoins de la pêche, Vol. 2. Commission des Communautés Européennes and FAO, Rome, pp. 845-886.
- Beck, M.W., K.L. Heck, K.W. Able, D.L. Childers, D.B. Eggleston, B.M. Gillanders, B. Halpern, C.G. Hays, K. Hoshino, T.J. Minello, R.J. Orth, P.F. Sheridan & M.P. Weinstein (2001): The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates: a better understanding of the habitats that serve as nurseries for marine species and the factors that create site-specific variability in nursery quality will improve conservation and management of these areas. Bioscience, 51(8), 633-641.
- Bilgili, A. & H. Kabasakal (2023): Encounters with threatened batoids from the perspective of a spearfisherman suggesting an aggregation site in southeastern Aegean Sea, Turkey. Regional Studies in Marine Science, 61, 102894.
- Bradai, M.N., B. Saidi & S. Enajjar (2012): Elasmobranchs of the Mediterranean and Black Sea: status, ecology and biology, Bibliographic analysis, Studies and Reviews. FAO, Rome, 103 pp.
- Capapé, C., O. Guélorget, C. Reynaud, A. Marquès, J.L. Bouchereau & J. Zaouali (2003): Effects of reproductive factors on interrelationships among three deep water sharks from northern Tunisia (Central Mediterranean). Annales, Ser. Hist. Nat., 13(2), 109-120.
- Capapé, C., J. Zaouali, J.A. Tomasini & J.L. Bouchereau (1992): Reproductive biology of the spiny butterfly ray, *Gymnura altavela* (Linnaeus, 1758) (Pisces: Gymnuridae) from off the Tunisian coasts. Sci. Mar., 56(4), 347-355.
- Castro, J.I. (1993): The shark nursery of Bulls Bay, South Carolina, with a review of the shark nurseries of the southeastern coast of the United States. Environ. Biol. Fishes., 38, 37-48.
- Castro, J.J. & E.K. Meyers (2022): Aggregative Behaviour of Spiny Butterfly Rays (*Gymnura altavela*, Linnaeus, 1758) in the Shallow Coastal Zones of Gran Canaria in the Eastern Central Atlantic. Animals, 13(9), 1455.
- Cengiz, Ç.C., C.M. Aydın, A. Ünlüoğlu & S. Akalin (2024): Length-weight relationship of the blackmouth catshark (*Galeus melastomus* Rafinesque, 1810) on the Turkish coasts of the Mediterranean Sea. Tethys Env. Sci., 1(3), 135-143.
- Compagno, L.J.V. & P.R. Last (1999): *Gymnuridae* Butterfly rays. In: Carpenter, K. E. & V. Neim (eds.): FAO Identification Guide for Fishery Purposes. The Living Marine Resources of the Western Pacific, Vol. 3. FAO, Rome, pp. 1506-1510.
- Conrath, C. & R. Scarbrough (2009): Biological Profiles: Spiny Butterfly Ray. <https://www.floridamuseum.ufl.edu/> (Last accession, 4 March 2025).
- Dulvy, N.K., P. Charvet, J. Carlson, L. Badji, M.P. Blanco-Parra, E. Chartrain, G. De Bruyne, D. Derrick, M. Dia, P. Doherty, J. Dossa, M. Ducrocq, G.H.L. Leurs, G. Notarbartolo di Sciarra, J.C. Pérez Jiménez, J.D. Pires, I. Seidu, F. Serena, A.-L. Soares, A. Tamo, M. Vacchi, R.H.L. Walls & A.B. Williams (2021): *Gymnura altavela*. The IUCN Red List of Threatened Species 2021: e.T63153A3123409. <https://www.iucnredlist.org/species/63153/3123409> (Last accession, 19 September 2024).
- Ebert, D.A. & M. Stehmann (2013): Sharks, batoids, and chimaeras of the North Atlantic. FAO Species Catalogue for Fishery Purposes. No. 7. FAO, Rome. 523 pp.
- Ebert, D.A. & M. Dando (2020): Field guide to sharks, rays & chimaeras of Europe and the Mediterranean, Vol. 20. Princeton University Press, New Jersey, 384 pp.
- El Kamel, O., N. Mnasri, M. Boumaïza, M.M.B. Amor, C. Reynaud & C. Capapé (2010): Additional records of the bull ray *Pteromyiaeus bovinus* (Chondrichthyes: Myliobatidae), in the Lagoon of Bizerte (Northern Tunisia, central Mediterranean). Annales, Ser. Hist. Nat., 20(2), 169-174.
- Ergüden, D., A. Uyan & S.A. Doğdu (2025): Rare occurrence of the rough ray *Raja radula* (Family: Rajidae) in the northeastern Mediterranean. Tethys Env. Sci., 2(2), 68-76.
- Espino Ruano, A., J.J. Castro, A. Guerra-Marrero, L. Couce-Montero, E.K.M. Meyers, A. Santana-del-Pino & D. Jimenez-Alvarado (2023): Aggregative Behaviour of Spiny Butterfly Rays (*Gymnura altavela*, Linnaeus, 1758) in the Shallow Coastal Zones of Gran Canaria in the Eastern Central Atlantic. Animals, 13(9), 1455.
- Gonçalves Silva, F. & M. Vianna (2018): Diet and reproductive aspects of the endangered butterfly ray *Gymnura altavela* raising the discussion of a possible nursery area in a highly impacted environment. Braz. J. Oceanogr., 66(3), 315-324.

- Gunter, G. (1967):** Some relationships of estuaries to the fisheries of the Gulf of Mexico. In: Lauff, G. H. (ed.): Estuaries. American Association for the Advancement of Science, Washington, pp. 621-638.
- Heithaus, M.R. (2007):** Nursery Areas as Essential Shark Habitats: A Theoretical Perspective. Am. Fish. Soc. Symp., 50, 3-13.
- Heupel, M.R., K.J. Carlson & C.A. Simpfendorfer (2007):** Shark nursery areas: concepts, definition, characterization and assumptions. Mar. Ecol. Prog. Ser., 337, 287-297.
- Kabasakal, H., U. Uzer, O. Gönülal & F.S. Karakulak (2024):** Bioecological lessons learned from the neonate longnose spurdogs, *Squalus blainville* (Squaliformes: Squalidae) suggest a potential nursery ground in the Marmara Sea. Acta Adriatica, 65.
- McEachran, J.D. & C. Capapé (1984):** Gymnuriidae. In: Carpenter K. E. (ed.): FAO species identification guide for fishery purposes. The living marine resources of the Western Central Atlantic, Vol. 1: Introduction, molluscs, crustaceans, hagfishes, sharks, batoid fishes and chimaeras. FAO, Rome, pp. 575-577.
- McEachran, J.D. & M.R. Carvalho (2002):** Rays. In: Carpenter K. E. (ed.): The living marine resources of the Western Central Atlantic, Vol. 1: Introduction, molluscs, crustaceans, hagfishes, sharks, batoid fishes and chimaeras. FAO, Rome, pp. 508-530.
- Medeiros, A.M., O.J. Luiz & C. Domit (2015):** Occurrence and use of an estuarine habitat by giant manta ray *Manta birostris*. J. Fish Biol., 86, 1830-1838.
- Murphy, H.M. & G.P. Jenkins (2010):** Observational methods used in marine spatial monitoring of fishes and associated habitats: a review. Mar. Freshw. Res., 61(2), 236-252.
- Özbek, E.Ö., M. Çardak & T. Kebapçioğlu (2016):** Spatio-temporal patterns of abundance, biomass and length-weight relationships of *Gymnura altavela* (Linnaeus, 1758) (Pisces: Gymnuridae) in the Gulf of Antalya, Turkey (Levantine Sea). J. Black Sea/Medit. Environ., 22, 16-34.
- Rangel, B.S., A. Rodrigues & R.G. Moreira (2018):** Use of a nursery area by cownose rays (Rhinopteridae) in southeastern Brazil. Neotrop. Ichthyol., 16, e170089.
- Rosenfelder, N., K. Lehnert, S. Kaffarnik, J.P.M. Torres, M. Vianna & W. Vetter (2012):** Thorough analysis of polyhalogenated compounds in ray liver samples off the coast of Rio de Janeiro, Brazil. Environ. Sci. Pollut. Res., 19, 379-389.
- Silva, F.G. & M. Vianna (2018):** Diet and reproductive aspects of the endangered butterfly ray *Gymnura altavela* raising the discussion of a possible nursery area in a highly impacted environment. Braz. J. Oceanogr., 66(3), 315-324.
- Simpfendorfer, C.A. & N.E. Milward (1993):** Utilisation of a tropical bay as a nursery area by sharks of the families Carcharhinidae and Sphyrnidae. Environ. Biol. Fishes., 37, 337-345.
- Taylan, B., B. Bayhan, C. Sağlam & A. Kara (2019):** First observation of the embryos of spiny butterfly ray, *Gymnura altavela* (Linnaeus, 1758) (Chondrichthyes: Gymnuridae) from Eastern Mediterranean, a species critically endangered. Fresenius Environ. Bull., 28, 3147-3152.
- Turan, C. & S.A. Doğdu (2022):** Preliminary assessment of invasive lionfish *Pterois miles* using underwater visual census method in the Northeastern Mediterranean. Croat. J. Fish., 80(1), 38-46.
- Turan, C., S.A. Doğdu & A. Soldo (2024):** Evidence of a nursery area and length-weight relationships for longnose spurdog *Squalus blainville* in the eastern Mediterranean Sea. Tethys Env. Sci., 1(4), 200-210.
- Turan, C., A. Uyan, A. Soldo, S.A. Doğdu & D. Ergüden (2025):** Checklist of cartilaginous species with current status and conservation strategies in Turkish marine waters. Tethys Env. Sci., 2(1), 31-61.
- Weigmann, S. (2016):** Annotated checklist of the living sharks, batoids and chimaeras (Chondrichthyes) of the world, with a focus on biogeographical diversity. J. Fish Biol., 88(3), 837-1037.
- Whitfield, P.E., J.A. Hare, A.W. David, S.L. Harter, R.C. Muñoz & C.M. Addison (2007):** Abundance estimates of the Indo-Pacific lionfish *Pterois volitans/miles* complex in the Western North Atlantic. Biol. Invasions, 9(1), 53-64.
- Yağlıoğlu, D., T. Deniz, M. Gürlek, D. Ergüden & C. Turan (2015):** Elasmobranch bycatch in a bottom trawl fishery in the Iskenderun Bay, northeastern Mediterranean. Cah. Biol. Mar., 56(3), 237-243.
- Yeldan, H. (2018):** Estimating some population parameters and stock assessment of spiny butterfly ray, *Gymnura altavela* (Linnaeus, 1758) the Levant Basin Coast (Northeastern Mediterranean). Indian J. Anim. Res., 52(12), 1790-1796.