

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 33, 2023, 2





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 33, 2023, 2

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 33, leto 2023, številka 2

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačič (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / *Società storica del Litorale - Capodistria*®Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / *Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente*®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / *Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran*
SI-6330 Piran / *Pirano*, Fornace/*Fornace* 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 23. 12. 2023.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS) in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / *La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in:* BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).To delo je objavljeno pod licenco / *Quest'opera è distribuita con Licenza / This work is licensed under a* Creative Commons BY-NC 4.0.Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/> / *The submission guidelines and all scientific journals and articles are available free of charge on the website https://zdjp.si/en/p/annalesshn/* / *Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito https://zdjp.si/en/p/annalesshn/*

VSEBINA / *INDICE GENERALE* / CONTENTSSREDOZEMSKÉ HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
*MEDITERRANEAN SHARKS AND RAYS***Christian CAPAPÉ, Christian REYNAUD & Farid HEMIDA**

The First Substantiated Records of Smoothback Angelshark *Squatina oculata* (Squatinae) from the Algerian Coast (Southwestern Mediterranean Sea) 143
Prvi utemeljeni zapis o pojavljanju pegastega sklata Squatina oculata (Squatinae) iz alžirske obale (jugozahodno Sredozemsko morje)

Tanguy CARPAYE-TAILAMEE & Mattéo MAUREL

Perspective on Great White Sharks (*Carcharodon carcharias*) in the Northwestern Mediterranean and Recommendations for Further Field Research 151
Pogled na velikega belega morskega volka (Carcharodon carcharias) v severozahodnem Sredozemlju in priporočila za nadaljnje terenske raziskave

Hakan KABASAKAL

A Preliminary Social Media Survey of Sharks and Batoids Captured in North Aegean Sea Commercial Fisheries 165
Preliminarna raziskava o morskih psih in skatih, ujetih v komercialnem ribištvu severnega Egejskega morja na osnovi podatkov iz socialnih medijev

Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ

On The Occurrence of Norwegian Skate, *Dipturus nidarosiensis* (Rajidae) on the Algerian Coast (Southwestern Mediterranean Sea) 187
O pojavljanju norveške raže, Dipturus nidarosiensis (Rajidae), ob alžirski obali (jugozahodno Sredozemsko morje)

Alen SOLDÓ

The First Record of Complete Albinism in Common Stingray *Dasyatis pastinaca* (Linnaeus, 1758) 193
Prvi zapis o najdbi popolnega albinističnega primerka navadnega morskega biča, Dasyatis pastinaca (Linnaeus, 1758)

Christian CAPAPÉ, Christian REYNAUD & Farid HEMIDA

Capture of a Giant Round Fantail Stingray *Taeniurops grabatus* (Dasyatidae) from the Algerian Coast (Southwestern Mediterranean Sea) 199
Ulov okroglega morskega biča (Taeniurops grabatus) (Dasyatidae) iz alžirske obale (jugozahodno Sredozemsko morje)

IHTIOFAUNA**ITTIOFAUNA****ICHTHYOFAUNA****Nicola BETTOSO & Diego BORME**

Recent Record of the Atlantic Pomfret *Brama brama* (Bonnaterre, 1788) (Scombriformes: Bramidae) in the Gulf of Trieste (Northern Adriatic Sea) 207
Recentni zapis o pojavljanju kostanjevke Brama brama (Bonnaterre, 1788) (Scombriformes: Bramidae) v Tržaškem zalivu (severno Jadransko morje)

Alan DEIDUN, Bruno ZAVA, Alessio MARRONE, Johann GALTIES, Arnold SCIBERRAS & Maria CORSINI-FOKA

The Confirmed Occurrence of *Schedophilus medusophagus* (Cocco, 1839) and *Petromyzon marinus* Linnaeus, 1758 in Maltese Waters, Central Mediterranean 213
Potrjeno pojavljanje meduzojedca, Schedophilus medusophagus (Cocco, 1839), in morskega piškurja, Petromyzon marinus Linnaeus, 1758, v malteških vodah, osrednje Sredozemsko morje

Gianni INSACCO, Gildo GAVANELLI, Bruno ZAVA & Maria CORSINI-FOKA

An Overlooked Finding of *Mola alexandrini* (Ranzani, 1839) in the Adriatic Sea 221
Sprejeto najdba vrste Mola alexandrini (Ranzani, 1839) v Jadranskem morju

Borut MAVRIČ, Lovrenc LIPEJ, Jelena BELAMARIĆ, Dule BULAJA, Matea ŠPIK & Petar KRUŽIĆ

Additional Data on the Bump-Head Sunfish, *Mola alexandrini* (Ranzani, 1839) in the Adriatic Sea 229
Dodatni podatki o pojavljanju grbastega morskega meseca, Mola alexandrini (Ranzani, 1839) v Jadranskem morju

Lana KHREMA, Amina ALNESSER, Adib SAAD & Christian CAPAPÉ

First Substantiated Record of Painted Eel
Echelus myrus (Ophichthidae) from the Syrian
 Marine Waters (Eastern Mediterranean Sea) 235
Prvi utemeljeni zapis o pojavljanju pisane
jegulje, Echelus myrus (Ophichthidae),
iz morskih voda Sirije (vzhodno
Sredozemsko morje)

BIOTSKA GLOBALIZACIJA
 GLOBALIZZAZIONE BIOTICA
 BIOTIC GLOBALIZATION

Deniz ERGUDEN, Deniz AYAS & Zafer KUŞATAN

The Presence of *Hippocampus fuscus* Rüppell,
 1838, in the Northeastern Mediterranean Sea 243
Pojavljanje morskega konjička vrste
Hippocampus fuscus Rüppell, 1838,
v severovzhodnem Sredozemskem morju

Christian CAPAPÉ & Adib SAAD

Confirmed Occurrence of Pharaoh
 Cardinal Fish *Apogonichthyoides*
pharaonis (Osteichthyes: Apogonidae)
 from the Syrian Coast
 (Eastern Mediterranean Sea) 249
Potrjeno pojavljanje faraonskega
kraljička Apogonichthyoides
pharaonis (Osteichthyes: Apogonidae)
iz sirske obale (vzhodno
Sredozemsko morje)

Deniz ERGUDEN, Deniz AYAS & Cemal TURAN

First Record of *Epinephelus areolatus*
 (Epinephelidae) from the South-Eastern
 Mediterranean, Turkey 255
Prvi zapis o pojavljanju rdečepikčaste
kirnje Epinephelus areolatus
(Epinephelidae) v jugovzhodnem
Sredozemskem morju (Turčija)

FAVNA
 FAUNA
 FAUNA

Andrea LOMBARDO & Giuliana MARLETTA

First Record of the Marine Heterobranch
Spinoaglaja wildpretii (Ortea, Bacallado &
 Moro, 2003) (Cephalaspidea: Aglajidae)
 in Sicily (Ionian Sea) with Notes on
 Its Biology and Ecology 263
Prvi zapis o pojavljanju morskega zaškrgarja
vrste Spinoaglaja wildpretii (Ortea,
Bacallado & Moro, 2003) (Cephalaspidea:
Aglajidae) na Siciliji (Jonsko morje) z
zapiski o njeni biologiji in ekologiji

Marco FANTIN, Saul CIRIACO, Lisa FARESI, Chiara SCRIGNER, Juri VECCHI, Domen TRKOV & Lovrenc LIPEJ

First Evidence of the Presence of *Okenia*
picoensis Paz-Sedano, Ortigosa & Pola,
 2017 (Gastropoda: Nudibranchia)
 in the Adriatic Sea 271
Prvi zapis o pojavljanju vrste gološkrgarja
Okenia picoensis Paz-Sedano, Ortigosa &
Pola, 2017 (Gastropoda: Nudibranchia)
iz Jadranskega morja

FLORA
 FLORA
 FLORA

Amelio PEZZETTA & MARCO PAOLUCCI

La flora di Lama dei Peligni (Abruzzo,
 Italia): aggiornamento sistematico e
 nuove segnalazioni 279
Flora občine Lama dei Peligni
(Abruci, Italija): sistematična
posodobitev in nove najdbe

MISCELLANEA**Al VREZEC**

Bird (Aves) Descriptions of Joannes
 Antonius Scopoli (1723-1788):
 General Overview 327
Opisi ptic (Aves) Joannesa Antoniusa Scopolija
(1723-1788): osnovni pregled

OCENE IN POROČILA
 RECENSIONI E RELAZIONI
 REVIEWS AND REPORTS

Mitja KALIGARIČ

Recenzija knjige:
 Podobe iz modrine 365

IN MEMORIAM**Elide CATALFAMO**

Ricordo del professor Guido Bressan 369

Kazalo k slikam na ovitku 371

Index to images on the cover 371

received: 2023-09-11

DOI 10.19233/ASHN.2023.20

A PRELIMINARY SOCIAL MEDIA-BASED SURVEY OF SHARKS AND BATOIDS CAPTURED IN COMMERCIAL FISHERIES OF THE NORTHERN AEGEAN SEA

Hakan KABASAKAL

Istanbul University, Institute of Science, Fisheries Technologies and Management Program, Süleymaniye, Esnaf Hastanesi 4. Kat,
34116 Fatih, İstanbul, Türkiye
WWF Türkiye
e-mail: kabasakal.hakan@gmail.com

ABSTRACT

In recent years, the use of social media films has been increasingly considered a promising method in elasmobranch research. The present study provides a regional example of good practice in using social media films for gathering additional and complementary information on the sharks and batoids in the northern Aegean Sea. A total of 67 shark and batoid individuals representing 8 orders, 15 families, and 21 species appeared in the examined video footages. The majority of the identified elasmobranchs consisted of sharks. Incidental captures of identified elasmobranchs mostly occurred in association with demersal fishing gears, frequently in demersal long-line fishery. The incidental captures of newborn specimens of blue shark and common angel shark support the suggestion that the northern Aegean Sea may serve as a parturition and/or nursery ground for both species.

Keywords: Elasmobranchs, captures, conservation, social media, opportunistic sampling

INDAGINE PRELIMINARE BASATA SUI SOCIAL MEDIA SU SQUALI E BATOIDI CATTURATI NELLA PESCA COMMERCIALE DEL MAR EGEO SETTENTRIONALE

SINTESI

Negli ultimi anni, l'uso dei filmati pubblicati sui social media è stato sempre più considerato un metodo promettente nella ricerca sugli elasmobranchi. Il presente studio fornisce un esempio regionale di buona pratica nell'uso dei filmati dei social media per raccogliere informazioni aggiuntive e complementari su squali e batoidi dell'Egeo settentrionale. Nei filmati esaminati sono apparsi un totale di 67 individui di squali e batoidi che rappresentano 8 ordini, 15 famiglie e 21 specie. La maggior parte degli elasmobranchi identificati era costituita da squali. Le catture accidentali degli elasmobranchi identificati si sono verificate soprattutto in associazione con attrezzature da pesca demersali, principalmente durante la pesca con palangari. Le catture accidentali di esemplari neonati di verdesca e pesce angelo supportano l'ipotesi che l'Egeo settentrionale possa fungere da area di nascita e/o crescita per entrambe le specie.

Parole chiave: elasmobranchi, catture, conservazione, social media, campionamento opportunistico

INTRODUCTION

Assessment of species diversity and abundance in chondrichthyans is an important step in the evaluation of their conservation status (Serena *et al.*, 2020). In the Mediterranean Sea, chondrichthyans, particularly large predatory sharks, have experienced a dramatic decline in numbers over the last two centuries (Ferretti *et al.*, 2008; Bargnesi *et al.*, 2020); therefore, every effort devoted to contributing to a better understanding of their life histories, distribution, etc., is valuable. For over 50 years, research programs have investigated the occurrence and distribution of chondrichthyans throughout the Mediterranean Sea, enhancing our in-depth understanding of their richness in the region (e.g., Capapé, 1989; Başusta *et al.*, 1998; Barrull *et al.*, 1999; De Maddalena & Pisticelli, 2001; Storai, 2004; Kabasakal & Kabasakal, 2004; Capapé *et al.*, 2006; Storai *et al.*, 2006; İşmen *et al.*, 2009; Damalas & Megalofonou, 2012; Sperone *et al.*, 2012; Ragonese *et al.*, 2013; Rafrafi-Nouira *et al.*, 2015; Ennajar *et al.*, 2022). Tracing the material and methods sections of these references, it becomes evident that the majority of the sampling has been carried out either using conventional methods of systematic scientific sampling or through opportunistic sampling in commercial fishing operations. Undoubtedly, monitoring the bycatch of commercial fisheries is a prolific source of data in chondrichthyan research (Kabasakal & Kabasakal, 2004; Kabasakal *et al.*, 2017; Mancusi *et al.*, 2020; Bonanomi *et al.*, 2018, 2022; Gallo *et al.*, 2022). However, the mining of social media has become an increasingly applied alternative method for collecting data in shark and batoid research (Boldrocchi & Storai, 2020; Kabasakal & Bilecenoğlu, 2020; Mancusi *et al.*, 2020; Taklis *et al.*, 2020; Kesici *et al.*, 2021; Gallo *et al.*, 2022; Bargnesi *et al.*, 2020, 2022).

The chondrichthyan fauna of the northeastern Aegean Sea has been investigated both in general ichthyological inventory studies (Eryılmaz, 2003; İşmen *et al.*, 2009; Cengiz *et al.*, 2011; Gönülal, 2016), and in a shark-specific survey (Kabasakal & Kabasakal, 2004). In this paper, the author reports the incidental captures of chondrichthyans in commercial fisheries operating in the northern Aegean Sea between September 2021 and August 2023, mainly based on data mined from mainstream social media channels. As a secondary objective, the author provides notes on the sharks and batoids that appeared in the examined footages.

MATERIAL AND METHODS

Area of investigation and type of fishery

The present study mostly covers the marine area surrounding the Gökçeada island and the mouth of

Saroz Bay in the northeastern Aegean Sea (Fig. 1). In this region, artisanal fishermen deploy trammel nets with a 42 mm knot-to-knot mesh opening (locally known as “marya” nets), along with demersal long-lines. In these fisheries, the primary targeted species include bony fishes such as the common dentex, *Dentex dentex* (Linnaeus, 1758), angler fish, *Lophius piscatorius* (Linnaeus, 1758), tub gurnard, *Chelidonichthys lucerna* (Linnaeus, 1758), and hake, *Merluccius merluccius* (Linnaeus, 1758). Commercial species of secondary importance include crustaceans, such as the European lobster, *Homarus gammarus* (Linnaeus, 1758) and European spiny lobster, *Palinurus vulgaris* Latreille, 1804, and cephalopods, such as *Octopus* sp. and *Sepia* sp. On rare occasions, pelagic teleosts such as swordfish, *Xiphias gladius* Linnaeus, 1758, and little tunny, *Euthynnus alletteratus* (Rafinesque, 1810) may also be captured.

Sampling methodology

The sampling approach in the present study was a typical example of opportunistic research (Jessup 2003), with data obtained solely from social media platforms Facebook, Instagram, and YouTube. To extract data from these platforms, a regular data mining survey was carried out once a week using the “Date Posted” filter in order to identify the most recent posts (Taklis *et al.*, 2020). The data search covers the period from September 2021 to August 2023. Furthermore, video footages recorded by local fishers between September 2021 and August 2023, documenting their catches and including scenes of incidentally captured elasmobranchs, were shared with the author for species identification. A total of 31 video footages (with a combined duration of 9 hours, 10 minutes, and 5 seconds) were analysed. An individual elasmobranch record was considered valid if the specimen depicted in a digital photograph was clearly visible in a side view; in the case of video footage, the specimen had to be visible for approximately 5 seconds, allowing for the capture of a still image for species identification (Kabasakal & Bilecenoğlu, 2020). To achieve this, the “Image Capture” function of the VLC Media Player was used. To facilitate the search and filter the information from the web, we used the following hashtags in Turkish (Kim *et al.*, 2016): “köpekbalıkları [sharks], yakalandı [captured], camgöz [dogfish], domuz köpekbalığı [angular rough shark], keler [angel shark], sapan balığı [thresher shark], mavi köpekbalığı [blue shark], büyük beyaz [great white shark], canavar (sea monster), kuzey Ege [north Aegean], Gökçeada [Gökçeada island], Saroz [Saroz Bay]”. To avoid repetition or duplication of records and to confirm their provenance, the owner of the original post was contacted for every incidence.




Fig. 1: Maps showing the approximate locations of capture of sharks (a) and batoids (b) in the northern Aegean Sea. Circles on the small maps indicate the locality of investigation in the Aegean Sea, as well as on the globe. Sl. 1: Zemljevidi s prikazi lokacij ulova morskih psov (a) in skatov (b) v severnem Egejskem morju. Krogci na manjših zemljevidih označujejo obravnavano območje v Egejskem morju, katerega lega je prikazana na planetarnem nivoju.




Fig. 2: Numerical occurrence of identified sharks and batoids with respect to fishing gear.
Sl. 2: Število primerkov določenih morskih psov in skatov glede na ribolovno orodje.

Since online communities and website administrators may react negatively to the utilisation of their online content by researchers, all internet content scraping activity was performed responsibly, following the ethical code proposed by Monkman *et al.* (2017) and avoiding compromising any personal data or image. Species identification follows Barone *et al.* (2022), while taxonomic nomenclature follows Froese and Pauly (2023). The IUCN Red List status of the identified elasmobranchs in the Mediterranean Sea follows Otero *et al.* (2019). Wherever possible, the following information was collected for each identified species through contacts with the owner of the post: total length (TL), total weight (TW), depth of capture (image of the echosounder

screen), and locality of deployment. To determine the sex of the specimens, the best image depicting the genital opening (for females) or claspers (for males) was captured, and the time code of the image was recorded. Neonates are defined as post-hatching or post-birth free-swimming young bearing fresh, unhealed, or healing umbilical scars in the case of placental species, or those at or near-birth size in the case of aplacental or ovoviparous species (Castro, 1993). Juveniles include all post-neonatal individuals prior to sexual maturation (Castro, 1993). The downloaded source videos as well as the captured frames of the identified species are preserved in the author's archives and available upon request for further examination.




Fig. 3: Numerical occurrence of identified sharks and batoids with respect to depth of capture.
Sl. 3: Število primerkov določenih morskih psov in skatov glede na globino ulova.

Data analysis

Differences in the number of identified elasmobranch species and the respective numbers of specimens were analysed with regards to type of fishing gear, depth of capture, and season of capture using Welch ANOVA, Kruskal-Wallis, and Dunn's post hoc tests. The latter was performed using Bonferroni-adjusted p -values (Parab & Bhalerao, 2010; Özeltindiş *et al.*, 2021). The chosen p -value was set at 0.05. Statistical analyses were performed employing PAST-Palaeontological statistics, version 4.03 (Hammer *et al.* 2001).

RESULTS

Total elasmobranchs

A total of 67 shark and batoid specimens representing 8 orders, 15 families, and 21 species appeared in the examined video footages. In addition to the identified species, there were several specimens identified at the generic level (a total of 6 genera). The identified species are presented in Tab. 1 in taxonomic order. Collated data regarding the date of capture, depth of capture, type of fishing gear used, species observed, and individual

Tab. 1: Taxonomic order of elasmobranch species identified in the video footages, along with the respective number of specimens (n=67) and the percentages of occurrence in the total number of specimens. DD: Data Deficient; LC: Least Concern; NT: Near-Threatened; VU: Vulnerable; EN: Endangered; CR: Critically Endangered. The asterisk (*) denotes species protected by law in Turkish seas.

Tab. 1: Taksonomski pregled prepoznanih vrst hrustančnic v videoposnetkih, število primerkov (n=67) in delež pojavljanja glede na celotno število primerkov. DD: Pomanjkljivi podatki; LC: Najmanj ogrožene; NT: potencialno ogrožene; VU: ranljive; EN: ogrožene; CR: skrajno ogrožene. Zvezdica (*) označuje z zakonom zaščitene vrste v turških morjih.

TAXONOMICAL ORDER	N	% of Total N	IUCN Red List Status for Mediterranean
HEXANCHIFORMES			
Hexanchidae			
<i>Hexanchus griseus</i> (Bonnaterre, 1788)	2	1.34	LC
LAMNIFORMES			
Lamnidae			
<i>Isurus oxyrinchus</i> Rafinesque, 1810*	1	0.67	CR
Alopiidae			
<i>Alopias vulpinus</i> (Bonnaterre, 1788)*	1	0.67	EN
CARCHARHINIFORMES			
Scyliorhinidae			
<i>Scyliorhinus canicula</i> (Linnaeus, 1758)	1	0.67	LC
<i>Scyliorhinus stellaris</i> (Linnaeus, 1758)	3	2.01	NT
<i>Scyliorhinus</i> sp.	3	2.01	
Triakidae			
<i>Mustelus asterias</i> Cloquet, 1819	1	0.67	VU
<i>Mustelus mustelus</i> (Linnaeus, 1758)	3	2.01	VU
Carcharhinidae			
<i>Prionace glauca</i> (Linnaeus, 1758)*	2	1.34	CR
SQUALIFORMES			
Dalatiidae			
<i>Dalatias licha</i> (Bonnaterre, 1788)	2	1.34	VU
Oxynotidae			
<i>Oxynotus centrina</i> (Linnaeus, 1758)*	2	1.34	CR
Squalidae			

<i>Squalus blainville</i> (Risso, 1827)*	9	6.03	DD
SQUATINIFORMES			
Squatinae			
<i>Squatina oculata</i> Bonaparte, 1840*	1	0.67	CR
<i>Squatina squatina</i> (Linnaeus, 1758)*	11	7.37	CR
TORPEDINIFORMES			
Torpedinidae			
<i>Torpedo marmorata</i> Risso, 1810	2	1.34	LC
<i>Torpedo</i> sp.	1	0.67	
RAJIFORMES			
Rajidae			
<i>Dipturus</i> sp.	1	0.67	
<i>Leucoraja naevus</i> (Müller & Henle, 1841)	1	0.67	NT
<i>Leucoraja</i> sp.	1	0.67	
<i>Raja clavata</i> Linnaeus, 1758*	4	2.68	NT
<i>Raja miraletus</i> Linnaeus, 1758	1	0.67	LC
<i>Raja</i> sp.	1	0.67	
<i>Rostroja alba</i> (Lacepède, 1803)	1	0.67	EN
MYLIOBATIFORMES			
Dasyatidae			
<i>Dasyatis pastinaca</i> (Linnaeus, 1758)	1	0.67	VU
<i>Dasyatis</i> sp.	2	1.34	
Aetobatidae			
<i>Aetomylaeus bovinus</i> (Geoffroy St. Hilaire, 1817)	2	1.34	CR
Myliobatidae			
<i>Myliobatis aquila</i> (Linnaeus, 1758)	7	4.69	VU




Fig. 4: Numerical occurrence of identified sharks and batoids with respect to season of capture.
Sl. 4: Število primerkov določenih morskih psov in skatov glede na sezono ulova.

remarks are presented in Appendix 1. The majority of the identified elasmobranchs consisted of sharks ($n=42$; 62.69%), whereas batoids comprised 37.31% ($n=25$) of the identified species. The most frequently observed species was *Squatina squatina* ($n=11$; 7.37%), followed by *Squalus blainville* ($n=9$; 6.03%), and *Myliobatis aquila* ($n=7$; 4.69%) (Tab. 1). As seen in Table 1, the percentage of occurrences for the remaining elasmobranchs was less than 6 percent of the total sample. Demersal elasmobranchs comprised the majority of the identified elasmobranchs ($n=63$; 94.03%), with only 5.97% of the total sample ($n=4$) represented by pelagic sharks (*Isurus oxyrinchus*, *Alopias vulpinus*, and *Prionace glauca*) (Tab. 1).

Elasmobranch species vs. fishing gears

Incidental captures of identified elasmobranchs mostly occurred in association with demersal fishing gears, with 50.75% ($n=34$) of the total sample

captured in demersal long-line fishery, followed by trammel netting ($n=27$; 40.30%). Pelagic long-lining ($n=3$) and recreational angling ($n=3$) individually comprised 4.48 percent of the total captures (Fig. 2). When analysing the captures based on the type of fishing gear used, there was no statistically significant difference (Kruskal-Wallis test, $p>0.05$, $p=0.24$). Since most of the captures (91.04%) occurred in demersal long-lining and trammel netting, a second analysis was conducted, focussing solely on catches in these two fisheries. However, no statistically significant difference was observed between demersal long-lining and trammel netting (Welch ANOVA, $p>0.05$, $p=0.68$; Kruskal-Wallis test, $p>0.05$, $p=0.32$).

Elasmobranch species vs. depth of capture

Depth of capture information was recorded for 51 specimens (76.12% of total elasmobranchs). The deepest depth of capture ranged from 500 to 600

m, with recreational anglers deploying a purpose-made shark tackle for game fishing targeting blunt-nose sixgill shark (*Hexanchus griseus*); however, the majority of captures occurred at depths between 51 and 100 m ($n=23$; 45.10%), followed by the 0-50 m, 101-200 m ($n=11$; 21.57% for both strata), and >200 m depth ranges ($n=6$; 11.76%) (Fig. 3). No statistically significant difference was found between the numerical distribution of captured specimens and depth of capture (Kruskal-Wallis test, $p>0.05$, $p=0.38$).

Elasmobranch species vs. season of capture

The seasonal distribution of elasmobranch captures is depicted in Figure 4, showing occurrences throughout the year. Captures were most abundant in August ($n=16$; 23.88%), followed by April ($n=13$; 19.40%), July ($n=10$; 14.93%), and January ($n=9$; 13.43%). The percentage of elasmobranch species present in commercial catches during the remaining months was less than 10 percent (ranging from 8.96% to 1.49%) (Fig. 4), with a complete absence of any elasmobranch species in the video footages of commercial catches in May and October. Nevertheless, no statistically significant difference was found in the seasonal distribution of elasmobranch captures (Kruskal-Wallis test, $p>0.05$, $p=0.17$; Dunn's post hoc test, $p>0.05$).

Notes on reproductive biology of several rare and endangered elasmobranchs observed in the studied videos

A close examination of the video footage ($n=31$; total duration 9:10:05 hours) in this study has revealed remarkable data suggesting that the northeastern Aegean Sea may serve as a potential nursery and/or breeding ground for several rare and endangered elasmobranchs.

On 30 July 2023, a young-of-the-year blunt-nose sixgill shark (*H. griseus*; TL ca. 150 cm) was incidentally captured in trammel net fishery at a 46-60 m depth range (Fig. 5a, b). The shark was completely motionless during the retention period onboard, and its post-release survival is uncertain.

On 22 June 2023, a newborn blue shark (*P. glauca*; TL ca. 50 cm) with a healing umbilical scar visible on the ventral surface between the pectoral fins, was incidentally captured in demersal long-lines deployed at a depth of 109 m (Fig. 5c, d). The newborn blue shark was released alive.

On 14 April 2022, a pregnant female common angel shark (*S. squatina*) became entangled in trammel nets deployed at a depth range of 19 to 25 m off the southwestern coast of Gökçeada island. The specimen aborted an unknown number

of near-term embryos onboard. On 3 March 2023, another pregnant female of *S. squatina* (TL ca. 170 cm) was captured by recreational anglers at a depth of 33 m, and aborted 5 near-term embryos upon being pulled onboard (Fig. 6a, b). In both instances, the pregnant females and pups were all released alive; however, their post-release survival status is not known. Available data suggest that mature and pregnant females, and juveniles (TL ca. 30–40 cm) frequently aggregate in the shallow areas (≤ 25 m depth) off the southwestern coast of Gökçeada island, while adult males occur more frequently in deeper areas (~60 m depth) between the island and the Gallipoli Peninsula.

Two mature females of *Mustelus mustelus* (TL ca. 110 cm and ca. 140 cm) and one year-of-the-young specimen (TL ca. 40 cm) were incidentally captured. Both females were retained and landed.

A juvenile shortfin mako shark (*Isurus oxyrinchus*; TL ca. 100 cm) was captured by a recreational angler off the northern coast of Gökçeada on 5 August 2023 (Fig. 6c).

Two female kite fin sharks (*Dalatias licha*; both TL ≥ 100 cm) were captured upon entanglement in trammel nets at a depth range of 171 to 300 m (Fig. 6d). Their swollen bellies suggested possible pregnancy, as preying on other specimens after becoming ensnared in trammel net, unlike when they get tangled in cod-ends of bottom trawls, may not be possible. Similarly, two mature female angular rough sharks (*Oxynotus centrina*; both TL ca. 60 cm) were captured individually in demersal long-lining and trammel netting in the northern littoral waters of Gökçeada. Their swollen bellies suggested potential pregnancy.

No data were obtained on the reproductive biology of other elasmobranchs identified from the video footages.

DISCUSSION AND CONCLUSIONS

For the last 50 years, the chondrichthyan fauna of the northern Aegean Sea has been investigated by several researchers through general ichthyological inventory studies, megafauna observations, or elasmobranch-specific studies. In a general ichthyological inventory study, Papaconstantinou and Tortonesi (1980) reported 15 species of demersal elasmobranchs from the Gulf of Thermaikos (northwestern Aegean Sea). Ulutürk (1987) identified 13 species of demersal and pelagic chondrichthyans in the waters surrounding the Gökçeada island (northeastern Aegean Sea). In a more recent general ichthyological inventory study, Eryılmaz (2003) documented 15 species of demersal sharks and batoids in the waters of the island of Bozcaada (also northeastern Aegean Sea).



Fig. 5: (a, b) Juvenile specimen of *Hexanchus griseus* (TL ca. 150 cm); (c, d) newborn specimen of *P. glauca* (TL ca. 50 cm), arrow indicating the unhealed umbilical scar.

Sl. 5: (a, b) Mladostni primerek vrste *H. griseus* (dolžina približno 150 cm); (c, d) novostoteni primerek vrste *P. glauca* (dolžina približno 50 cm), puščica označuje nezaceljeno poporodno brazgotino.

In one of the few studies specifically focussed on sharks, Kabasakal and Kabasakal (2004) recorded 20 species in the region, including megafauna members such as *Carcharodon carcharias* and *Cetorhinus maximus*, and rare squaliform sharks such as *D. licha* and *O. centrina*. According to Cengiz *et al.* (2011), the fish fauna of the Gulf of Saroz comprises 28 chondrichthyan species, including the rare lamniform shark, *Carcharias taurus*. However, since the authors have no information on where the examined specimen of the latter was deposited, this record is considered questionable. In an extensive study on the biodiversity of the northern Aegean Sea, Altuğ *et al.* (2011) reported the presence of 12 species of demersal sharks and batoids in the study area. In a recent survey on deep-sea fishes of the northern Aegean Sea, Gönülal (2016) recorded 13 species of chondrichthyans occurring in depths ranging between 500 and 1000 m.

The common feature of the studies mentioned so far is that the sampling method has always been extractive sampling, carried out using fishing gears such as bottom trawl, beam trawl, trammel net, gill net, or demersal longline. Also, for many years, fishery-dependent surveys have been the primary, if not the only, source of scientific information on most chondrichthyan species in the northern Aegean Sea and in the Mediterranean Sea (e.g., Kabasakal & Kabasakal, 2004; Damalas & Megalofonou, 2012; Sperone *et al.*, 2012; Kabasakal *et al.*, 2017). In recent years, however, the contribution of local ecological knowledge (LEK) and citizen science has steadily increased. And although social media-based data mining as a research method cannot substitute traditional stratified/random *in situ* surveys in the studies on chondrichthyans in the northern Aegean Sea, it can provide complementary data to fill the knowledge gaps arising from financial constraints or a shortage of researchers.

A recent study carried out by Taklis *et al.* (2020) underscored the value and importance of social media in collecting baseline information, while identifying and/or addressing important conservation issues related to sharks in Greece, with a specific mention of chondrichthyan records from the northern Aegean Sea. Kabasakal and Bilecenoğlu (2020) compiled an inventory of rare and large shark species in Turkish waters solely from internet-based news, which also included current northern Aegean records. Marine top predators found in the waters around Gökçeada have been investigated in a study that used LEK as a supporting tool for data collection (Kesici *et al.*, 2021) and reported rare occurrences of *C. carcharias* (n=2) and *I. oxyrinchus*

(n=4) around the island. Following data mining digital archives and gathering supportive data from citizen science and social media, Moutopoulos *et al.* (2022) examined historical records of shark presence during the early developmental phases of the Greek fishery, highlighting historical records of *C. carcharias* and *C. maximus* in the northern Aegean Sea. As corroborated by the cited current studies (Kabasakal & Bilecenoğlu, 2020; Taklis *et al.*, 2020; Kesici *et al.*, 2021; Moutopoulos *et al.*, 2022), social media have proven to be an increasingly efficient tool in the efforts for the conservation of and raising awareness on large predatory sharks, which can notably supplement the monitoring of the cartilaginous fish status in a certain region and time interval (Boldrocchi & Storai, 2021; Casola *et al.*, 2022).

Fishery-dependent extractive surveys have revealed that chondrichthyan species are being incidentally captured in multi-modal fisheries operating across an extensive area of the northern Aegean Sea, ranging from the shallows of the continental shelf to bathyal grounds (≤ 1000 m depth) (Papaconstantinou & Tortonese, 1980; Ulutürk, 1987; Eryılmaz, 2003; Kabasakal & Kabasakal, 2004; Gönülal, 2016). The majority (>90%) of shark and batoid species identified in the present study were captured by small-scale fishermen (trammel-netters and demersal longliners). This is consistent with the findings of Giovos *et al.* (2021), who emphasised that the underreporting of catches by small-scale fisheries poses a threat to elasmobranchs. According to Giovos *et al.* (2021), commercial gill-netting and demersal long-lining each contribute to over 40 percent of the bycatch of sharks and batoids in the north Aegean Sea. With the implementation of laws aimed at protecting cartilaginous fish, fishers tend to not report such catches, as highlighted by Giovos *et al.* (2021). Therefore, an ethical examination of the videos uploaded by fishermen on social media, driven as much by the desire to showcase their (legal) catches as by self-promotion (Monkman *et al.*, 2017; Shiffman, 2018), can be an effective method of recording previously unreported cartilaginous fish catches. The present preliminary study relies on the examination of 31 video footages, but a more comprehensive understanding of the bycatch impact would require the implementation of a questionnaire and onboard monitoring involving fisheries observers, and social media-based investigations complementing the data collected through large-scale fisheries surveys.

The present data set provides useful observations on the potential reproductive areas for



Fig. 6: (a) Pregnant female of *Squatina squatina* (TL ca. 170 cm) aborting near-term embryos while retained on board, with the arrow indicating the aborted specimen; (b) a near-term embryo of *S. squatina* before being released alive; (c) juvenile specimen of *Isurus oxyrinchus* (TL ca. 100 cm); (d) the arrow indicates a female specimen of *Dalatias licha* with two tub gurnards (*Chelidonichthys lucerna*) on the right.

Sl. 6: (a) Breja samica vrste *Squatina squatina* (dolžina približno 170 cm), ki je na krovu splavila skoraj razvite zarodke, s puščico, ki označuje splavljen primerek; (b) skoraj popolno razvit zarodek vrste *S. squatina*, preden so ga živega izpustili; (c) mladostni primerek vrste *Isurus oxyrinchus* (dolžina približno 100 cm); (d) puščica označuje samico vrste *Dalatias licha* z dvema velikima krulcema (*Chelidonichthys lucerna*) na desni.

several elasmobranch species in the area encompassing the islands of Gökçeada and Semadirek, the Gulf of Saroz, and the Gallipoli Peninsula. Previous occurrences of newborns, juveniles, and pregnant females in this region can be summarised as follows: Newborn specimens of *H. griseus* (TL 660 mm) and *D. licha* (TL range

338–372.5 mm), observed bearing unhealed umbilical scars, have been reported from the upper slope waters off the northern coast of the island of Gökçeada (Kabasakal & Kabasakal, 2002, 2004). Rare occurrences of newborn specimens of *O. centrina* observed in bottom-trawling have been reported from the Gulf of Thermaikos (TL

248 mm; Papaconstantinou & Tortonese, 1980) and off the southwestern coast of the island of Bozcaada (TL 225 mm; Eryilmaz, 2003). On 22 September 2008, a mature female of *O. centrina* carrying 12 developing ova (TL 651 mm) was captured in a scientific bottom-trawl survey in the Gulf of Saros (Yiğın et al., 2016). Taklis et al. (2020) reported captures of juvenile specimens of *P. glauca* (TW range 2000–5000 g) in commercial fisheries in various locations across the northern Aegean Sea. The capture of a newborn blue shark in demersal long-lining (which was subsequently released alive), documented in the present study, supports the suggestion that the northern Aegean Sea may serve as a parturition and nursery ground for *P. glauca*. Recent records of newborn specimens of *I. oxyrinchus* and *C. carcharias* from several locations in the northern Aegean Sea have been mentioned in Kabasakal (2015), Tunçer and Kabasakal (2016), Taklis et al. (2020), and Kabasakal et al. (2022).

While not included in the present study, juvenile specimens of *Etmopterus spinax* (TL range 120–227 mm), incidentally captured in commercial bottom-trawling at depths of 300 to 400 m, have also been reported (Kabasakal & Kabasakal, 2004). A juvenile specimen of *Squatina oculata* (TL 300 mm) and a subadult specimen of *S. squatina* (TL 750 mm) have been reported off the island of Gökçeada (Kabasakal & Kabasakal, 2004). İşmen et al. (2009) reported the observation of a newborn *S. squatina* (TL 232 mm) in the Gulf of Saros. In the present study, observations of newborn specimens and pregnant females of *S. squatina* confirmed the presumed areas of reproduction for the common angel shark in the northern Aegean Sea. Last but not least, pregnant females of *Mustelus asterias* (TL 1530 mm; bearing 21 near-term embryos) and of *S. blainville* (TL 870 mm; bearing 7 near-term embryos) have been incidentally captured in the Gulf of Saros (Kabasakal & Kabasakal, 2004). Therefore, past and recent occurrences of newborns, juveniles, and pregnant females in the studied region support the possibility of multiple elasmobranch nurseries across the shelf and bathyal grounds of the northern Aegean Sea. This has led to the designation of an Important Shark and Ray Area (ISRA) and an Area of Interest (Aoi) (Jabado et al., 2023).

The geographical area in which the study was conducted falls within the boundaries of the proposed buffer zone of the recently designated Thracian Sea Shelf ISRA on account of the presence of threatened species (e.g., *A. bovinus*), range-restricted species (*Raja radula*), reproductive areas (e.g., *S. canicula*), and undefined ag-

gregations (*D. pastinaca*) (Jabado et al., 2023). Furthermore, the geographical boundaries of the Truva Shelf Aoi, where potential reproduction areas for *Squatina* species are assumed to exist, overlap with the southern part of the area investigated in the present study (Jabado et al., 2023). As a result, the findings of this study contribute supportive new data that reinforce the establishment of the Thracian Sea Shelf ISRA and the Truva Shelf Aoi.

Eight of the species identified in this study are protected in Turkish seas. In addition, the majority of these species (n=16; 76.19%) fall within the susceptible categories of Near Threatened, Vulnerable, Endangered, or Critically Endangered of the IUCN Red List (Otero et al., 2019). The majority of captured elasmobranchs (n=63; 94.02%) were released alive without prolonged retention on deck and harsh handling, which is promising for the survival of bycaught elasmobranchs in the region. The fishermen's statements in the examined footages demonstrate their awareness of conservation laws as well as recognition of protected (e.g., *Squatina* spp., *O. centrina*) or vulnerable species (e.g., *D. licha*), as does their release of the captured specimens back to the sea. This corroborates the suggestion by Boldrocchi and Storai (2020) that social media can play a crucial role in raising fishermen's awareness about cartilaginous fish and bridging knowledge gaps.

To conclude, this study presents a regional example of a good practice in using social media films to gather additional and complementary information about the life histories and current status of elasmobranch species in a specific region. In recent years, the utilisation of footage published on social media films has emerged as an increasingly promising data collection method and an approach supported by the findings of various researchers (Boldrocchi & Storai, 2020; Mancusi et al., 2020; Taklis et al., 2020; Kesici et al., 2021; Bargnesi et al., 2020, 2022; Gallo et al., 2022). Compared to traditional systematic scientific sampling, it is clear that this study approach still presents some uncertainties (e.g., TL information dependent on the statements of the fishermen) and weaknesses (e.g., the intervals at which the fishermen upload footage may not accurately reflect the true periodicity of fishing days). However, considering the *K-selected* life history characteristics of elasmobranchs (e.g., slow growth, late maturation, low fecundity, etc.) and the current extinction crisis that threatens their survival (Fowler et al., 2005; Dulvy et al., 2021), this non-invasive and non-extractive visual sampling method, if applied according to

the methodology described herein, can complement traditional systematic sampling by addressing the knowledge gaps that arise from limited research staff or expansion of the study area. Last but not least, the present study evidently suffers from a very small sample size. Therefore, more effort is needed before drawing any conclusions about the impact of bycatch on the populations of sharks and batoids in the northern Aegean Sea.

ACKNOWLEDGMENTS

The author thanks the fishermen who shared the footages of their commercial captures on social media or personally with the author. Special thanks go to two anonymous reviewers for their critical revisions, which improved the content and quality of the article.

Appendix 1: Elasmobranch species (n=67) captured in commercial fisheries in the northern Aegean Sea and appearing in social media videos, along with fishing data, remarks about the observed specimens, and sources of video footages. Specimen time code indicates the time interval during which an elasmobranch specimen is visible in the respective video.

Priloga 1: Hrustančnice (n=67 vrst), ujete v komercialnem ribolovu v severnem Egejskem morju, ki se pojavljajo v videoposnetkih v socialnih medijih, skupaj s podatki o ulovu, opombami v zvezi z opazovanimi primerki in viri videoposnetkov. Časovna koda primerka označuje časovni interval, v katerem je bil primerek opažen v danem videoposnetku.

Date	Locality	Depth (m)	Fishing gear	Species	Nr	Specimen time codes	Remarks	Source
14 Sep 2021	Gökçeada, off Yıldız Bay	85-92	Demersal long line	<i>Mustelus mustelus</i>	1	9:49-12:26	Female, TL ca. 110 cm; taken on board	https://www.youtube.com/watch?v=LeNLmXCkkm0
14 Sep 2021	Gökçeada, off Yıldız Bay	85-92	Demersal long line	<i>Raja clavata</i>	1	12:58-13:38	Female, DW ca. 40 cm; released alive	https://www.youtube.com/watch?v=LeNLmXCkkm0
14 Sep 2021	Gökçeada, off Yıldız Bay	85-92	Demersal long line	<i>Raja clavata</i>	1	13:48-14:14	Female, DW ca. 50 cm; released alive	https://www.youtube.com/watch?v=LeNLmXCkkm0
14 Sep 2021	Gökçeada, off Yıldız Bay	85-92	Demersal long line	<i>Raja clavata</i>	1	15:25-15:50	Male, adult; claspers very longer than pelvic fins and visible at the time code (15:29) of the source video; DW ca. 70 cm; released alive	https://www.youtube.com/watch?v=LeNLmXCkkm0
14 Sep 2021	Gökçeada, off Yıldız Bay	85-92	Demersal long line	<i>Mustelus mustelus</i>	1	16:40-19:20	Female, genital opening is visible at the time code (19:01) of the source video, TL ca. 140 cm; taken on board	https://www.youtube.com/watch?v=LeNLmXCkkm0
20 Nov 2021	Between Gökçeada and Semadirek islands, off SW of Semadirek island, between Enez and Gökçeada	102-107	Demersal long line	<i>Squalus blainville</i>	1	22:17-22:36	Female, genital opening is visible at the time code (22:30) of the source video, TL ca. 50 cm; released alive; depth information given at the time codes (18:32 and 33:55)	https://www.youtube.com/watch?v=PNCcAy7f7g
20 Nov 2021	Between Gökçeada and Semadirek islands, off SW of Semadirek island, between Enez and Gökçeada	102-107	Demersal long line	<i>Myliobatis aquila</i>	1	22:44-23:08	Female, DW ca. 100 cm; dorsal fin is behind pelvic fins (22:48); released alive; depth information given at the time codes (18:32 and 33:55)	https://www.youtube.com/watch?v=PNCcAy7f7g
15 Dec 2021	Between Gökçeada and Semadirek islands, around Zürafa rocks point	90-320	Demersal long line	<i>Scyliorhinus stellaris</i>	1	21:15-21:53	Female, TL ca. 100 cm; released alive; depth information is given at the time codes (3:18-3:21) of the source video	https://www.youtube.com/watch?v=rTr1_WgVA84
15 Dec 2021	Between Gökçeada and Semadirek islands, around Zürafa rocks point	90-320	Demersal long line	<i>Myliobatis aquila</i>	1	29:57-30:35	Female, DW ca. 100 cm; no claspers (30:31); dorsal fin is behind pelvic fins (30:35); released alive; depth information is given at the time codes (3:18-3:21) of the source video	https://www.youtube.com/watch?v=rTr1_WgVA84
15 Jan 2022	Between Gökçeada and Semadirek islands, around Zürafa rocks point	?	Demersal long line	<i>Myliobatis aquila</i>	1	8:45-8:56	DW ca. 60 cm; released alive	https://www.youtube.com/watch?v=IHfYuOwNolk
15 Jan 2022	Between Gökçeada and Semadirek islands, around Zürafa rocks point	?	Demersal long line	<i>Squalus blainville</i>	1	16:04-16:26	TL ca. 40 cm; released alive	https://www.youtube.com/watch?v=IHfYuOwNolk

15 Jan 2022	Between Gökçeada and Semadirek islands, around Zürafa rocks point	?	Demersal long line	<i>Mustelus mustelus</i>	1	21:49-22:04	TL ca. 40 cm; released alive	https://www.youtube.com/watch?v=IHfYuOwNolk
1 Apr 2022	Gökçeada-Büyükkemikli	500-600	Rod and reel	<i>Hexanchus griseus</i>	1	14:40-19:45	Captured with a purpose-made shark rig for big game fishery, baited with bonito, <i>Sarda sarda</i> , released alive, TL ca. 450 cm, female	https://www.youtube.com/watch?v=k6T-LjX8kN0
7 Apr 2022	Tuzla	14-16	Trammel net	<i>Squatina squatina</i>	1	1:43-2:27	All female and released alive	Video provided by the fisherman and available on request
14 Apr 2022	Between Kaşkaval cape and Kabatepe on mainland	70	Demersal long line	<i>Aetomylaeus bovinus</i>	1	6:10-6:30	Claspers are visible between the time codes (6:10-6:30) of the source video, released alive	https://www.youtube.com/watch?v=bF8lwek96sc
14 Apr 2022	Between Kaşkaval cape and Kabatepe on mainland	70	Demersal long line	<i>Aetomylaeus bovinus</i>	1	14:19-14:54	Claspers are visible at the time code (14:44) of the source video, released alive	https://www.youtube.com/watch?v=bF8lwek96sc
14 Apr 2022	Between Kaşkaval cape and Kabatepe on mainland	70	Demersal long line	<i>Mustelus asterias</i>	1	9:15-9:50	Landed for auction	https://www.youtube.com/watch?v=bF8lwek96sc
14 Apr 2022	Between Kaşkaval cape and Kabatepe on mainland	70	Demersal long line	<i>Oxynotus centrina</i>	1	10:28-12:46	Female, TL ca. 60 cm, genital opening is visible at the time code (11:56) of the source video, released alive	https://www.youtube.com/watch?v=bF8lwek96sc
14 Apr 2022	Between Kaşkaval cape and Kabatepe on mainland	70	Demersal long line	<i>Dipturus</i> sp.	1	13:10-13:32	Released alive	https://www.youtube.com/watch?v=bF8lwek96sc
14 Apr 2022	Pirgos	19-25	Trammel net	<i>Squatina squatina</i>	1	1:15-1:30	Pregnant female, aborted nearterm embryos on board, all released alive but their post-release survival status uncertain	Video provided by the fisherman and available on request
6 Jun 2022	Tepeköy coast	10-15 m	Trammel net	<i>Torpedo</i> sp.	1	12:51-13:00	Released alive	https://www.youtube.com/watch?v=INbVVQoQs-g
10 Jun 2022	Kefaloz	20	Trammel net	<i>Squatina squatina</i>	1	0:00-1:07	4 adult females (TL 80-90 cm and TW 10-20 kg) 4 juveniles (TL 30-40 cm), sexes not reported	Video provided by the fisherman and available on request
11 Aug 2022	Between Kaşkaval cape and Kabatepe on mainland	60	Trammel net	<i>Torpedo marmorata</i>	1	5:43-5:46 13:37-13:50	Spiracle tentacles and dorsal texture are visible between the time codes (13:37-13:39) of the source video, released alive	https://www.youtube.com/watch?v=Wv3ck9wCsqU
11 Aug 2022	Between Kaşkaval cape and Kabatepe on mainland	60	Trammel net	<i>Squatina squatina</i>	1	7:31-7:50	Male, claspers are visible at the time code (7:40) of the source video, released alive	https://www.youtube.com/watch?v=Wv3ck9wCsqU
11 Aug 2022	Between Kaşkaval cape and Kabatepe on mainland	60	Trammel net	<i>Squatina squatina</i>	1	7:59-8:01	Specimen is visible at the time codes (7:59-8:01) of the source video, released alive	https://www.youtube.com/watch?v=Wv3ck9wCsqU
11 Aug 2022	Between Kaşkaval cape and Kabatepe on mainland	60	Trammel net	<i>Squatina squatina</i>	1	9:52-10:12	Specimen is visible at the time codes (9:52-10:12) of the source video, juvenile male with minute claspers shorter than pelvic fins (10:09), released alive	https://www.youtube.com/watch?v=Wv3ck9wCsqU
17 Aug 2022	North coast of Gökçeada	?	Demersal long line	<i>Myliobatis aquila</i>	1	10:18-10:24	Released alive	https://www.youtube.com/watch?v=fgXlkx2dUTo

17 Aug 2022	North coast of Gökçeada	?	Demersal long line	<i>Scyliorhinus canicula</i>	1	23:20-23:29	Released alive	https://www.youtube.com/watch?v=fgXlkx2dUTo
2 Sep 2022	Between Gökçeada and Semadirek islands, around Zürafa rocks point	95-100	Demersal long line	<i>Rostroraja alba</i>	1	13:39-14:03	Immature male, short clasper is visible at the time code (13:49) of the source video; snout very long and pointed (13:51); tip of pectoral fins angular (13:52); released alive	https://www.youtube.com/watch?v=rgYVaHoLPI
1 Jan 2023	North coast of Gökçeada	117-171	Demersal long line	<i>Squalus blainville</i>	1	13:51-14:10	Female, genital opening is visible at the time code (14:08) of the source video; released alive	https://www.youtube.com/watch?v=3b0gHFvSQyE
14 Jan 2023	Between Kefaloz point and Kabatepe on mainland	27	Trammel net	<i>Leucoraja naevus</i>	1	12:39-12:50	Male, claspers are visible at the time code (12:43) of the source video, released alive	https://www.youtube.com/watch?v=LBkx3zFGE-Y
14 Jan 2023	Between Kefaloz point and Kabatepe on mainland	27	Trammel net	<i>Squatina squatina</i>	1	15:33-16:00	Subadult male, claspers are visible at the time code (15:52) of the source video, released alive	https://www.youtube.com/watch?v=LBkx3zFGE-Y
14 Jan 2023	Between Kefaloz point and Kabatepe on mainland	27	Trammel net	<i>Squatina squatina</i>	1	16:32-16:52	Subadult male, claspers are visible at the time code (16:33) of the source video, released alive	https://www.youtube.com/watch?v=LBkx3zFGE-Y
14 Jan 2023	Between Kefaloz point and Kabatepe on mainland	27	Trammel net	<i>Squatina squatina</i>	1	17:00-17:11	Released alive	https://www.youtube.com/watch?v=LBkx3zFGE-Y
14 Jan 2023	Between Kefaloz point and Kabatepe on mainland	27	Trammel net	<i>Leucoraja</i> sp.	1	17:46-17:55	Released alive	https://www.youtube.com/watch?v=LBkx3zFGE-Y
14 Jan 2023	Between Kefaloz point and Kabatepe on mainland	27	Trammel net	<i>Squatina squatina</i>	1	19:34-19:48	Both released alive	https://www.youtube.com/watch?v=LBkx3zFGE-Y
3 Feb 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	231	Demersal long line	<i>Myliobatis aquila</i>	1	11:31-11:38	Released alive	https://www.youtube.com/watch?v=nXo9pPNUcvk
3 Feb 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	231	Demersal long line	<i>Squalus blainville</i>	1	12:03-12:28	Female, genital opening is visible at the time code (12:26) of the source video; no anal fin; TL ca. 50 cm; released alive	https://www.youtube.com/watch?v=nXo9pPNUcvk
3 Feb 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	231	Demersal long line	<i>Squalus blainville</i>	1	15:14-15:27	Female, genital opening is visible at the time code (15:16) of the source video; no anal fin; TL ca. 50 cm; released alive	https://www.youtube.com/watch?v=nXo9pPNUcvk
3 Feb 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	231	Demersal long line	<i>Myliobatis aquila</i>	1	16:20-16:23	DW ca. 100 cm, released alive	https://www.youtube.com/watch?v=nXo9pPNUcvk
3 Mar 2023	Çanakkale Strait, southern entrance	?	Recreational angling	<i>Squatina squatina</i>	1	0:00-1:20	1 pregnant female, TL ca. 170 cm, giving birth on board; 5 newborns; all released alive, but post-release survival is unknown	https://www.instagram.com/reel/CpUZYM0ghmz?igshid=YmMyMTA2M2Y=

22 Apr 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	86	Demersal long line	<i>Squalus blainville</i>	1	7:20-7:42	Female, genital opening is visible at (7:40); time codes of the descriptive characters in the source video are as follows: free rear tips of pectoral fins, and continuous posterior edge of caudal fin (7:40), spurs in front of dorsal fins (7:41), released alive	https://www.youtube.com/watch?v=TbSdp2-F0Tg
22 Apr 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	86	Trammel net	<i>Scyliorhinus stellaris</i>	1	11:59-13:02	Female, genital opening is visible at the time code (12:54) of the source video, TL ca. 100 cm; descriptive characters are visible at the following time codes: widely separated nasal flaps (12:35), dorsally positioned long eyes (12:35), released alive	https://www.youtube.com/watch?v=TbSdp2-F0Tg
12 Jun 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	103	Trammel net	<i>Squalus blainville</i>	1	9:04-10:22	All released alive; time codes of the descriptive characters in the source video are as follows: white coloration of the posterior edge of first dorsal fin (9:10), white coloration and continuous contour of the ventral edge of upper caudal lobe (9:12), free rear tip of pectoral (9:13) and spur (11:53)	https://www.youtube.com/watch?v=PegisdVPFMg
12 Jun 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	103	Trammel net	<i>Myliobatis aquila</i>	1	15:21-15:35	Female, genital opening is visible at the time code (15:29) of the source video; some other descriptive characters are visible at the following time codes of the same video; whip-shaped tail (15:30), short and obtuse snout (15:32) released alive	https://www.youtube.com/watch?v=PegisdVPFMg
19 Jun 2023	Between Gökçeada and Semadirek islands	300	Trammel net	<i>Dalatias licha</i>	1	0:00-0:11	Not alive, the swollen belly of the specimen suggests that it was a pregnant female, TL > 100 cm	https://www.instagram.com/reel/Ctqe3lUASpc/?igshid=MzRlODBiNWFlZA==
8 Jly 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	125	Demersal long line	<i>Dasyatis pastinaca</i>	1	16:13-16:24	Released alive	https://www.youtube.com/watch?v=HXFrVvayR8
15 Jly 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	171	Trammel net	<i>Dalatias licha</i>	1	10:35-10:45	Not alive, the swollen belly of the specimen suggests that it was a pregnant female, TL > 100 cm	https://www.youtube.com/watch?v=y8yU1bgrfJM
15 Jly 2023	Between Gökçeada and Semadirek islands	?	Pelagic long line for sword fish	<i>Alopias vulpinus</i>	1	1:30-1:52	Female, TL ca. 400 cm	Video provided by the fisherman and available on request
15 Jly 2023	Between Gökçeada and Semadirek islands	?	Pelagic long line for sword fish	<i>Prionace glauca</i>	1	0:00-1:00	Female, TL 250-300 cm	Video provided by the fisherman and available on request
22 Jly 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	109	Demersal long line	<i>Prionace glauca</i>	1	14:14-15:23	New-born female; TL ca. 50 cm, healing umbilical scar on ventral surface is visible at the time code (14:52) of the source video, released alive	https://www.youtube.com/watch?v=BtS00_DN8Q
30 Jly 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	46-60	Trammel net	<i>Scyliorhinus</i> sp.	1	10:29-10:31	Released alive	https://www.youtube.com/watch?v=ZX6VgINwdcM

30 Jly 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	46-60	Trammel net	<i>Oxynotus centrina</i>	1	16:48-17:43	Female, TL ca. 60 cm, genital opening is visible at the time code (17:35) of the source video; retained in a recirculating water container to recovery, then released alive	https://www.youtube.com/watch?v=ZX6VgINwdcM
30 Jly 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	46-60	Trammel net	<i>Hexanchus griseus</i>	1	19:28-21:34	TL ca. 150 cm, juvenile, the following descriptive characters are visible at the time codes of the source video: 6 gill slits (21:05), single dorsal fin (21:06) and dentition (21:15); the specimen swallowed a piece of net and spiral valve is extended out from the cloaca (19:30-19:35); fishermen released the specimen, but it's post-release survival is not certain	https://www.youtube.com/watch?v=ZX6VgINwdcM
30 Jly 2023	Between Gökçeada and Semadirek islands, around Zürafa rocks point	46-60	Demersal long line	<i>Dasyatis</i> sp.	1	38:23-38:37	Broke the line and escaped	https://www.youtube.com/watch?v=ZX6VgINwdcM
31 Jly 2023	Saroz Bay	33	Recreational angling	<i>Torpedo marmorata</i>	1	0:00-0:23	In the footage it is visible that the head and pectoral fins are fused, head-pectoral fin complex moving with continuous undulations, and the origin of the second dorsal fin is at the level of pelvic fin tips; released alive	https://www.instagram.com/reel/CvULGBnN4wR/?igshid=MTc4MmM1Yml2Ng==
4 Aug 2023	Entrance of Saroz Bay, Büyükkemikli	?	Demersal long line	<i>Squalus blainville</i>	1	5:16-5:28	Released alive	https://www.youtube.com/watch?v=jAlyQD-U2as
4 Aug 2023	Entrance of Saroz Bay, Büyükkemikli	?	Demersal long line	<i>Raja</i> sp.	1	5:31-5:46	Released alive	https://www.youtube.com/watch?v=jAlyQD-U2as
4 Aug 2023	Entrance of Saroz Bay, Büyükkemikli	?	Demersal long line	<i>Dasyatis</i> sp.	1	5:48-6:03	Released alive	https://www.youtube.com/watch?v=jAlyQD-U2as
4 Aug 2023	Entrance of Saroz Bay, Büyükkemikli	?	Trammel net	<i>Scyliorhinus</i> sp.	1	14:16-14:23	Released alive	https://www.youtube.com/watch?v=jAlyQD-U2as
4 Aug 2023	Entrance of Saroz Bay, Büyükkemikli	?	Trammel net	<i>Scyliorhinus</i> sp.	1	22:32-22:50	TL ca. 100 cm; released alive	https://www.youtube.com/watch?v=jAlyQD-U2as
4 Aug 2023	Entrance of Saroz Bay, Büyükkemikli	?	Trammel net	<i>Squalus blainville</i>	1	24:23-24:45	TL ca. 50 cm; released alive	https://www.youtube.com/watch?v=jAlyQD-U2as
5 Aug 2023	Gökçeada	?	Pelagic long line for sword fish	<i>Isurus oxyrinchus</i>	1	0:00-0:38	TL ca. 100 cm; no umbilical scar is visible between pectoral fins on ventral surface; released alive	https://www.tiktok.com/@agasonfishing/video/7263391305470872837?_t=8egjCQnMnZd&_i=1
8 Aug 2023	Gökçeada	224-300	Trammel net	<i>Raja miraletus</i>	1	5:46-6:00	DW ca. 40 cm; retained onboard	https://www.youtube.com/watch?v=u2pLcdiKCDs
8 Aug 2023	Gökçeada	224-300	Trammel net	<i>Squatina oculata</i>	1	11:42-12:20	Female, TL ca. 120 cm; released alive	https://www.youtube.com/watch?v=u2pLcdiKCDs
8 Aug 2023	Gökçeada	160	Demersal long line	<i>Scyliorhinus stellaris</i>	1	21:20-21:25	Female, TL ca. 100 cm; depth information is given at the time code (16:44-16:46) of the source video; released alive	https://www.youtube.com/watch?v=u2pLcdiKCDs
8 Aug 2023	Gökçeada	160	Demersal long line	<i>Raja clavata</i>	1	22:01-22:12	Female, DW ca. 50 cm; released alive	https://www.youtube.com/watch?v=u2pLcdiKCDs

PRELIMINARNA RAZISKAVA O MORSKIH PSIH IN SKATIH, UJETIH V KOMERCIALNEM RIBIŠTVU SEVERNEGA EGEJSKEGA MORJA NA OSNOVI PODATKOV IZ SOCIALNIH MEDIJEV

Hakan KABASAKAL

Istanbul University, Institute of Science, Fisheries Technologies and Management Program, Süleymaniye, Esnaf Hastanesi 4. Kat, 34116
Fatih, İstanbul, Türkiye
WWF Türkiye
e-mail: kabasakal.hakan@gmail.com

POVZETEK

*V zadnjih letih so filmi v socialnih medijih vedno bolj uporabni kot obetavna metoda za pridobivanje podatkov o hrustančnicah. Pričujoča raziskava predstavlja primer dobre prakse pri uporabi filmov v socialnih medijih za zbiranje dodatnih in dopolnilnih informacij o morskih psih in skatih v severnem Egejskem morju. V pregledanih videoposnetkih je avtor prepoznal 67 primerkov morskih psov in skatov, ki so pripadali 8 redovom, 15 družinam in 21 vrstam. Večina je bilo morskih psov. Naključno ujete hrustančnice so bile ujete s pridnim ribolovnim orodjem, pri čemer so dobro polovico primerkov ujeli s parangalom. Naključni ulovi novoskotenih primerkov sinjega morskega psa (*Prionace glauca*) in navadnega sklata (*Squatina squatina*) podpirajo domnevo, da lahko predstavlja severno Egejsko morje jaslice za obe vrsti morskih psov.*

Ključne besede: hrustančnice, ulovi, ohranjanje, socialni mediji, oportunistično vzorčenje

REFERENCES

- Altuğ, G., Y. Aktan, M. Oral, B. Topaloğlu, A. Dede, Ç. Keskin, M. İşinibilir, M. Çardak, & P.S. Çiftçi (2011): Biodiversity of the northern Aegean Sea and southern part of the Sea of Marmara. *Mar. Biodivers. Rec.*, 4, e65. doi:10.1017/S1755267211000662.
- Bargnesi, F., S. Lucrezi, & F. Ferretti (2020): Opportunities from citizen science for shark conservation, with a focus on the Mediterranean Sea. *Eur. Zool. J.*, 87, 20-34. doi: 10.1080/24750263.2019.1709574.
- Bargnesi, F., S. Moro, A. Leone, I. Giovos, & F. Ferretti (2022): New technologies can support data collection on endangered shark species in the Mediterranean Sea. *Mar. Ecol. Prog. Ser.*, 689, 57-76. https://doi.org/10.3354/meps14030.
- Barone, M., C. Mazzoldi & F. Serena (2022): Sharks, Rays and Chimaeras in Mediterranean and Black Sea – Key to Identification. FAO, Rome, http://dx.doi.org/10.4060/cc0830en.
- Barrull, J., I. Mate & M. Bueno (1999): Observaciones de tiburones (Chondrichthyes Euselachii) en aguas de Cataluña (Mediterráneo NO), con algunos aspectos generales de su ecología. *Sci. gerund.*, 24, 127-151.
- Başusta, N., Ü. Erdem & C. Çevik (1998): An investigation on chondrichthyes in Iskenderun Bay. *CBUJOS Series (Biology)*, 1, 63-69. (in Turkish).
- Boldrocchi, G. & T. Storai (2020): Data-mining social media platforms highlights conservation action for the Mediterranean Critically Endangered blue shark *Prionace glauca*. *Aquatic Conserv. Mar. Freshw. Ecosyst.*, 31, 3087-3099. doi: 10.1002/aqc.3690.
- Bonanomi, S., J. Pulcinella, C.M. Fortuna, F. Moro & A. Sala (2018): Elasmobranch bycatch in the Italian Adriatic pelagic trawl fishery. *PLoS one*, 13(1), e0191647.
- Bonanomi, S., F. Moro, A. Colombelli, J. Pulcinella & C.M. Fortuna (2022): A 14-year time series of marine megafauna bycatch in the Italian midwater pair trawl fishery. *Sci. Data*, 9, 51. doi: 10.1038/s41597-022-01155-2.
- Cengiz, Ö., A. İşmen, U. Özekinci & A. Öztekin (2011): An investigation on fish fauna of Saros Bay (Northern Aegean Sea). *AKU-J. Sci.*, 11, 31-37.
- Capapé, C. (1989): Les Sélaciens des côtes méditerranéennes: aspects généraux de leur écologie et exemples de peuplements. *Océanis*, 15, 309-331.
- Capapé, C., O. Guélorget., Y. Vergne, A. Marquès & J.-P. Quignard (2006): Skates and rays (Chondrichthyes) from waters off the Languedocian coast (southern France, northern Mediterranean). *Annales, Ser. Hist. Nat.*, 16, 166-178.
- Casola, W.R., J.M. Beall, M.N. Peterson, L.R. Larson & C.S. Price (2022): Influence of social media on fear of sharks, perceptions of intentionality associated with shark bites, and shark management preferences. *Front. Commun.*, 7. doi: 10.3389/fcomm.2022.1033347.
- Castro, J.I. (1993): The shark nursery of Bulls Bay, South Carolina, with a review of the shark nurseries of the southeastern coast of the United States. In: Demski, L.S., Wourms, J.P. (Eds.), *The reproduction and development of sharks, skates, rays and ratfishes*. *Environ. Biol. Fishes*, 38, 37-48.
- Damalas, D. & P. Megalofonou (2012): Occurrences of large sharks in the open waters of the southeastern Mediterranean Sea. *J. Nat. Hist.*, 46, 2701-2723. http://dx.doi.org/10.1080/00222933.2012.716864.
- De Maddalena, A. & L. Pisticelli (2001): Analisi preliminare dei selaci registrati presso il mercato ittico di Milano (aprile-settembre 2000). *Boll. Mus. Civ. Stor. Nat. Venezia*, 52, 129-145.
- Dulvy, N.K., N. Pacoureau, C.L. Rigby, R.A. Pollom, R.W. Jabado, D.A. Ebert, B. Finucci, C.M. Pollock, J. Cheok, D.H. Derrick, K.B. Herman, C.S. Sherman, W.J. Vander Wight, J.M. Lawson, R.H.L. Walls, J.K. Carlson, P. Charvet, K.K. Bineesh, D. Fernando, G.M. Ralph, J.H. Matsushiba, C. Hilton-Taylor, S.V. Fordham & C.A. Simpfendorfer (2021): Overfishing drives over onethird of all sharks and rays toward a global extinction crisis. *Curr. Biol.*, 31, 1-15. https://doi.org/10.1016/j.cub.2021.08.062.
- Ennajjar, S., B. Saïdi & M.N. Bradaï (2022): Elasmobranchs in Tunisia: Status, Ecology, and Biology. In book: *Sharks - Past, Present and Future [Working Title]*. doi: http://dx.doi.org/10.5772/intechopen.108629.
- Eryilmaz, L. (2003): A study on the fishes of Bozcaada island (north Aegean Sea). *Turk. J. Mar. Sci.*, 9, 121-137.
- Ferretti, F., R.A. Myers, F. Serena & H.K. Lotze (2008): Loss of large predatory sharks from the Mediterranean Sea. *Conserv. Biol.*, 22, 952-964.
- Fowler, S.L., R.D. Cavanagh, M. Camhi, G.H. Burgess, G.M. Cailliet, S.V. Fordham, C.A. Simpfendorfer & J.A. Musick (2005): *Sharks, Rays and Chimaeras: The Status of the Chondrichthyan Fishes*. Status Survey. IUCN/SSC Shark Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK. x + 461 pp.
- Froese, R. & D. Pauly (eds.) (2023): *FishBase*. World Wide Web electronic publication. www.fishbase.org (last accession: 31 October 2023).
- Gallo, S., C. Mancusi, A. Mohammed, B. Rigers, A. Barash, M. Barone, M. Bottaro, P. Carbonara, C. Roberto, I. Četković, S. Clo, E. de Sabata, S. Enajjar, E. Shakman, F. Garibaldi, G. Giglio, I. Giovos, H. Kabasakal, L. Lanteri, S. Lelli, L. Lipej, C. Mazzoldi, P. Micarelli, G. Morey, S. Moro, M.N. Bradai, G.N. di Sciara, C. Romano, B. Saïdi, A. Soldo, E. Sperone, F. Tiralongo, B. Zava & F. Serena (2022): Sightings of large elasmobranchs from the Mediterranean: new data from MEDLEM database in the last five years (2017–2022). 2022 IEEE International Workshop on Metrology for the Sea; Learning to Measure Sea Health Parameters (MetroSea), pp. 293-297, doi: 10.1109/MetroSea55331.2022.9950984.

- Giovas, I., R.N. Aga Spyridopoulou, N. Doumpas, K. Glaus, P. Kleitou, Z. Kazlari, D. Katsada, D. Loukovitis, I. Mantzouni, M. Papapetrou, Y.P. Papastamatiou & D. K. Moutopoulos (2021): Approaching the “real” state of elasmobranch fisheries and trade: A case study from the Mediterranean. *Ocean Coast. Manag.*, 211. <https://doi.org/10.1016/j.ocecoaman.2021.105743>.
- Gönülal, O. (2016): A preliminary study on the bathymetric distribution of the deep sea fishes from Northern Aegean Sea. *Bio. Di. Con.*, 9, 144-148.
- Hammer, Ø., D.A.T. Harper & D.R. Paul (2001): Past: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontol. Electron.*, 4: Available at: http://palaeo-electronica.org/2001_1/past/issue1_01.htm.
- İşmen, A., C.Ç. Yiğın, U. Altınağaç & A. Ayaz (2009): Length-weight relationships for ten shark species from Sargos Bay (North Aegean Sea). *J. Appl. Ichthyol.*, 25(Suppl. 1), 109-112. doi: 10.1111/j.1439-0426.2009.01263.x.
- Jabado, R.W., E. García-Rodríguez, P.M. Kyne, R. Charles, A.H. Armstrong, J. Bortoluzzi, T.L. Mouton, A. Gonzalez Pestana, A. Battle-Morera, C. Rohner & G. Notarbartolo di Sciarra (2023): Mediterranean and Black Seas: A regional compendium of Important Shark and Ray Areas. Dubai: IUCN SSC Shark Specialist Group.
- Jessup, D.A. (2003): Opportunistic research and sampling combined with fish and wildlife management actions or crisis response. *ILAR Journal*, 44, 277-285.
- Kabasakal, H. (2015): Occurrence of shortfin mako shark, *Isurus oxyrinchus* Rafinesque, 1810, off Turkey's coast. *Mar. Biodivers. Rec.*, 8. doi:10.1017/S1755267215001104.
- Kabasakal, H., E. Bayrı & G. Alkan (2022): Distribution and status of the great white shark, *Carcharodon carcharias*, in Turkish waters: a review and new records. *Annales, Ser. Hist. Nat.*, 32, 325-342. doi:10.19233/ASHN.2022.34
- Kabasakal, H. & M. Bilecenoglu (2020): Shark infested internet: an analysis of internet-based media reports on rare and large sharks of Turkey. *FishTaxa*, 16, 8-18.
- Kabasakal, H. & E. Kabasakal (2002): Morphometrics of young kitefin sharks, *Dalatias licha* (Bonnaterre, 1788), from northeastern Aegean Sea, with notes in its biology. *Annales, Ser. Hist. Nat.*, 12, 161-166.
- Kabasakal, H. & E. Kabasakal (2004): Sharks captured by commercial fishing vessels off the coast of Turkey in the northern Aegean Sea. *Annales, Ser. Hist. Nat.*, 14, 171-180.
- Kabasakal, H., S.Ü. Karhan & S. Sakınan (2017): Review of the distribution of large sharks in the seas of Turkey (eastern Mediterranean). *Cah. Biol. Mar.*, 58, 219-228. doi:10.21411/CBM.A.96D9F948.
- Kesici, N.B., C. Dalyan, O. Gönülal, A. Akkaya, P. Lyne, S. Tüzün & B. Yıldırım (2021): A preliminary study on marine top predators inhabiting Gökçeada Island, the North Aegean Sea. *J. Black Sea/Medit. Environ.*, 27, 34-48.
- Kim, Y., J. Huang & S. Emery (2016): Garbage in, garbage Out: data collection, quality assessment and reporting standards for social media data use in health research, infodemiology and digital disease detection. *J. Med. Internet Res.*, 18. doi: 10.2196/jmir.4738.
- Mancusi, C., R. Baino, C. Fortuna, L. de Sola, G. Morey, M. Bradai, A. Kallianotis, A. Soldo, F. Hemida, A. Saad, M. Dimech, P. Peristeraki, M. Bariche, S. Clo, E. de Sabata, L. Castellano, F. Garibaldi, L. Lanteri, F. Tinti, A. Pais, E. Sperone, P. Micarelli, F. Poisson, L. Sion, R. Carlucci, D. Cebrian-Menchero, B. Séret, F. Ferretti, A. El-Far, İ. Saygu, E. Shakman, A. Bartoli, J. Gualart, D. Damalas, P. Megalofonou, M. Vacchi, M. Bottaro, G. Notarbartolo di Sciarra, M. Follesa, R. Cannas, H. Kabasakal, B. Zava, G. Cavlan, A. Jung, M. Abudaya, J. Kolutari, A. Barash, A. Joksimovic, B. Marceta, L. G. Vilas, F. Tiralongo, I. Giovos, F. Bargnesi, S. Lelli, M. Barone, S. Moro, C. Mazzoldi, C. Charis, A. Abella & F. Serena (2020): MEDLEM database, a data collection on large Elasmobranchs in the Mediterranean and Black seas. *Mediterr. Mar. Sci.*, 21, 276-288.
- Monkman, G.G., M. Kaiser & K. Hyder (2017): The Ethics of Using Social Media in Fisheries Research, *Rev. Fish. Sci., Aquac.* <http://dx.doi.org/10.1080/23308249.2017.1389854>.
- Moutopoulos, D.K., E. Lazari, G. Katselis & I. Giovos (2022): From extermination to conservation: historical records of shark presence during the early and development phase of the Greek fishery. *Animals*, 12. <https://doi.org/10.3390/ani12243575>.
- Otero, M., F. Serena, V. Gerovasileiou, M. Barone, M. Bo, J.M. Arcos, A. Vulcano & J. Xavier (2019): Identification guide of vulnerable species incidentally caught in Mediterranean fisheries. IUCN, Malaga, Spain, 204 pp.
- Özaltındış, T., A.M. Köse & E. Ö. Özdamar (2021): ANOM Test as an Alternative Approach to Group Comparisons. *SDU J. Nat. Appl., Sci.* 25, 89-98. doi:10.19113/sdufenbed.820031.
- Papaconstantinou, C. & E. Tortonese (1980): On a collection of fishes from Thermaikos Gulf (NE Greece). *Thalassographica*, 3, 15-42.
- Parab, S. & S. Bhaleao (2010): Choosing statistical test. *Int. J. Ayurveda Res.*, 1, 187-191.
- Rafrafi-Nouira, S., O. El Kamel-Moutalibi, C. Reynaud, M. Boumaïza & C. Capapé (2015): Additional and unusual captures of elasmobranch species from the northern coast of Tunisia (central Mediterranean). *J. Ichthyol.*, 55, 337-345.
- Ragonese, S., S. Vitale, M. Dimech & S. Mazzola (2013): Abundances of demersal sharks and chimaera from 1994-2009 scientific surveys in the central Mediterranean Sea. *PLoS One*, 8. doi:10.1371/journal.pone.0074865.

Serena, F., A.J. Abella, F. Bargnesi, M. Barone, F. Colloca, F. Ferretti, F. Fiorentino, J. Jenrette & S. Moro (2020): Species diversity, taxonomy and distribution of chondrichthyes in the Mediterranean and Black Sea. *Eur. Zool. J.*, 87, 497-536. <https://doi.org/10.1080/24750263.2020.1805518>.

Shiffman, D.S. (2018): Social media for fisheries science and management professionals: how to use it and why you should? *Fisheries*, 43, 123-129. doi:10.1002/fsh.10031.

Sperone, E., G. Parise, A. Leone, C. Milazzo, V. Circosta, G. Santoro, G. Paolillo, P. Micarelli & S. Tripepi (2012): Spatiotemporal patterns of distribution of large predatory sharks in Calabria (central Mediterranean, southern Italy). *Acta Adriat.*, 53, 13-24..

Storai, T. (2004): Check-list degli elasmobranchi delle acque Toscane (Mar Ligure meridionale, Mar Tirreno settentrionale). Parte I: Squali. *Atti Soc. tosc. Sci. nat. Mem., Serie B.* 111. 7-11.

Storai, T., B. Cristo, M. Zuffa, L. Zinzula, A. Floris & A.T. Campanile (2006): The Sardinian large elasmobranch database. *Cybium*, 30, 141-144.

Taklis, C., I. Giovos & A.A. Karamanlidis (2020): Social media: a valuable tool to inform shark conservation in Greece. *Mediterr. Mar. Sci.*, 21, 493-498. doi:<http://dx.doi.org/10.12681/mms.22165>.

Tunçer, S. & H. Kabasakal (2016): Capture of a juvenile shortfin mako shark, *Isurus oxyrinchus* Rafinesque, 1810 (Chondrichthyes: Lamnidae) in the Bay of Edremit, northern Aegean Sea (Turkey). *Annales, Ser. Hist. Nat.*, 26, 31-36. doi:10.19233/ASHN.2016.5.

Ulutürk, T. (1987): Fish fauna, background radioactivity of the Gökçeada marine environment. *Aquat. Sci. Eng.*, 1, 95-119. (formerly Journal of Fisheries & Aquatic Sciences of Istanbul University).

Yığın, C.Ç., A. İşmen & U. Önal (2016): Occurrence of a rare shark, *Oxynotus centrina* (Chondrichthyes: Oxynotidae), from Saros Bay, North Aegean Sea, Turkey. *J. Black Sea/Medit. Environ.*, 22, 103-109.