

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 33, 2023, 2





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 33, 2023, 2

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 33, leto 2023, številka 2

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / *Società storica del Litorale - Capodistria*[®]Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / *Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente*[®]**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / *Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran*
SI-6330 Piran / *Pirano*, Fornace/*Fornace* 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 23. 12. 2023.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS) in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / *La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in:* BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).To delo je objavljeno pod licenco / *Quest'opera è distribuita con Licenza / This work is licensed under a* Creative Commons BY-NC 4.0.Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/> / *The submission guidelines and all scientific journals and articles are available free of charge on the website https://zdjp.si/en/p/annalesshn/* / *Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito https://zdjp.si/en/p/annalesshn/*

VSEBINA / *INDICE GENERALE* / CONTENTSSREDOZEMSKÉ HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
*MEDITERRANEAN SHARKS AND RAYS***Christian CAPAPÉ, Christian REYNAUD & Farid HEMIDA**

The First Substantiated Records of Smoothback Angelshark *Squatina oculata* (Squatinae) from the Algerian Coast (Southwestern Mediterranean Sea) 143
Prvi utemeljeni zapis o pojavljanju pegastega sklata Squatina oculata (Squatinae) iz alžirske obale (jugozahodno Sredozemsko morje)

Tanguy CARPAYE-TAILAMEE & Mattéo MAUREL

Perspective on Great White Sharks (*Carcharodon carcharias*) in the Northwestern Mediterranean and Recommendations for Further Field Research 151
Pogled na velikega belega morskega volka (Carcharodon carcharias) v severozahodnem Sredozemlju in priporočila za nadaljnje terenske raziskave

Hakan KABASAKAL

A Preliminary Social Media Survey of Sharks and Batoids Captured in North Aegean Sea Commercial Fisheries 165
Preliminarna raziskava o morskih psih in skatih, ujetih v komercialnem ribištvu severnega Egejskega morja na osnovi podatkov iz socialnih medijev

Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ

On The Occurrence of Norwegian Skate, *Dipturus nidarosiensis* (Rajidae) on the Algerian Coast (Southwestern Mediterranean Sea) 187
O pojavljanju norveške raže, Dipturus nidarosiensis (Rajidae), ob alžirski obali (jugozahodno Sredozemsko morje)

Alen SOLDÓ

The First Record of Complete Albinism in Common Stingray *Dasyatis pastinaca* (Linnaeus, 1758) 193
Prvi zapis o najdbi popolnega albinističnega primerka navadnega morskega biča, Dasyatis pastinaca (Linnaeus, 1758)

Christian CAPAPÉ, Christian REYNAUD & Farid HEMIDA

Capture of a Giant Round Fantail Stingray *Taeniurops grabatus* (Dasyatidae) from the Algerian Coast (Southwestern Mediterranean Sea) 199
Ulov okroglega morskega biča (Taeniurops grabatus) (Dasyatidae) iz alžirske obale (jugozahodno Sredozemsko morje)

IHTIOFAUNA**ITTIOFAUNA****ICHTHYOFAUNA****Nicola BETTOSO & Diego BORME**

Recent Record of the Atlantic Pomfret *Brama brama* (Bonnaterre, 1788) (Scombriformes: Bramidae) in the Gulf of Trieste (Northern Adriatic Sea) 207
Recentni zapis o pojavljanju kostanjevke Brama brama (Bonnaterre, 1788) (Scombriformes: Bramidae) v Tržaškem zalivu (severno Jadransko morje)

Alan DEIDUN, Bruno ZAVA, Alessio MARRONE, Johann GALTIES, Arnold SCIBERRAS & Maria CORSINI-FOKA

The Confirmed Occurrence of *Schedophilus medusophagus* (Cocco, 1839) and *Petromyzon marinus* Linnaeus, 1758 in Maltese Waters, Central Mediterranean 213
Potrjeno pojavljanje meduzojedca, Schedophilus medusophagus (Cocco, 1839), in morskega piškurja, Petromyzon marinus Linnaeus, 1758, v malteških vodah, osrednje Sredozemsko morje

Gianni INSACCO, Gildo GAVANELLI, Bruno ZAVA & Maria CORSINI-FOKA

An Overlooked Finding of *Mola alexandrini* (Ranzani, 1839) in the Adriatic Sea 221
Spregledana najdba vrste Mola alexandrini (Ranzani, 1839) v Jadranskem morju

Borut MAVRIČ, Lovrenc LIPEJ, Jelena BELAMARIĆ, Dule BULAJA, Matea ŠPIK & Petar KRUŽIĆ

Additional Data on the Bump-Head Sunfish, *Mola alexandrini* (Ranzani, 1839) in the Adriatic Sea 229
Dodatni podatki o pojavljanju grbastega morskega meseca, Mola alexandrini (Ranzani, 1839) v Jadranskem morju

Lana KHREMA, Amina ALNESSER, Adib SAAD & Christian CAPAPÉ

First Substantiated Record of Painted Eel
Echelus myrus (Ophichthidae) from the Syrian
 Marine Waters (Eastern Mediterranean Sea) 235
Prvi utemeljeni zapis o pojavljanju pisane
jegulje, Echelus myrus (Ophichthidae),
iz morskih voda Sirije (vzhodno
Sredozemsko morje)

BIOTSKA GLOBALIZACIJA
 GLOBALIZZAZIONE BIOTICA
 BIOTIC GLOBALIZATION

Deniz ERGUDEN, Deniz AYAS & Zafer KUŞATAN

The Presence of *Hippocampus fuscus* Rüppell,
 1838, in the Northeastern Mediterranean Sea 243
Pojavljanje morskega konjička vrste
Hippocampus fuscus Rüppell, 1838,
v severovzhodnem Sredozemskem morju

Christian CAPAPÉ & Adib SAAD

Confirmed Occurrence of Pharaoh
 Cardinal Fish *Apogonichthyoides*
pharaonis (Osteichthyes: Apogonidae)
 from the Syrian Coast
 (Eastern Mediterranean Sea) 249
Potrjeno pojavljanje faraonskega
kraljička Apogonichthyoides
pharaonis (Osteichthyes: Apogonidae)
iz sirske obale (vzhodno
Sredozemsko morje)

Deniz ERGUDEN, Deniz AYAS & Cemal TURAN

First Record of *Epinephelus areolatus*
 (Epinephelidae) from the South-Eastern
 Mediterranean, Turkey 255
Prvi zapis o pojavljanju rdečepikčaste
kirnje Epinephelus areolatus
(Epinephelidae) v jugovzhodnem
Sredozemskem morju (Turčija)

FAVNA
 FAUNA
 FAUNA

Andrea LOMBARDO & Giuliana MARLETTA

First Record of the Marine Heterobranch
Spinoaglaja wildpretii (Ortea, Bacallado &
 Moro, 2003) (Cephalaspidea: Aglajidae)
 in Sicily (Ionian Sea) with Notes on
 Its Biology and Ecology 263
Prvi zapis o pojavljanju morskega zaškrgarja
vrste Spinoaglaja wildpretii (Ortea,
Bacallado & Moro, 2003) (Cephalaspidea:
Aglajidae) na Siciliji (Jonsko morje) z
zapiski o njeni biologiji in ekologiji

Marco FANTIN, Saul CIRIACO, Lisa FARESI, Chiara SCRIGNER, Juri VECCHI, Domen TRKOV & Lovrenc LIPEJ

First Evidence of the Presence of *Okenia*
picoensis Paz-Sedano, Ortigosa & Pola,
 2017 (Gastropoda: Nudibranchia)
 in the Adriatic Sea 271
Prvi zapis o pojavljanju vrste gološkrgarja
Okenia picoensis Paz-Sedano, Ortigosa &
Pola, 2017 (Gastropoda: Nudibranchia)
iz Jadranskega morja

FLORA
 FLORA
 FLORA

Amelio PEZZETTA & MARCO PAOLUCCI

La flora di Lama dei Peligni (Abruzzo,
 Italia): aggiornamento sistematico e
 nuove segnalazioni 279
Flora občine Lama dei Peligni
(Abruci, Italija): sistematična
posodobitev in nove najdbe

MISCELLANEA**Al VREZEC**

Bird (Aves) Descriptions of Joannes
 Antonius Scopoli (1723-1788):
 General Overview 327
Opisi ptic (Aves) Joannesa Antoniusa Scopolija
(1723-1788): osnovni pregled

OCENE IN POROČILA
 RECENSIONI E RELAZIONI
 REVIEWS AND REPORTS

Mitja KALIGARIČ

Recenzija knjige:
 Podobe iz modrine 365

IN MEMORIAM**Elide CATALFAMO**

Ricordo del professor Guido Bressan 369

Kazalo k slikam na ovitku 371

Index to images on the cover 371

THE CONFIRMED OCCURRENCE OF *SCHEDOPHILUS MEDUSOPHAGUS* (COCCO, 1839) AND *PETROMYZON MARINUS* LINNAEUS, 1758 IN MALTESE WATERS, CENTRAL MEDITERRANEAN SEA

Alan DEIDUN

Department of Geosciences, University of Malta, Msida MSD 2080, Malta
e-mail: alan.deidun@um.edu.mt

Bruno ZAVA

Wilderness studi ambientali, via Cruillas 27, 90146 Palermo, Italy
Museo Civico di Storia Naturale di Comiso, Via degli Studi 9, Comiso (RG), Italy
e-mail: wildernessbz@hotmail.com

Alessio MARRONE

Department of Geosciences, University of Malta, Msida MSD 2080, Malta
e-mail: alessio.marrone@um.edu.mt

Johann GALDIES

Department of Geosciences, University of Malta, Msida MSD 2080, Malta
e-mail: johann.galdies@um.edu.mt

Arnold SCIBERRAS

The Exterminator Ltd. 5, The Service Hub, Triq San Gorg, Naxxar NXR 2541 Malta
e-mail: bioislets2@gmail.com

Maria CORSINI-FOKA

Institute of Oceanography, Hellenic Centre for Marine Research. Hydrobiological Station of Rhodes, Cos Street, 85100 Rhodes, Greece
e-mail: mcorsini@hcmr.gr

ABSTRACT

Specimens of two uncommon fish species, Schedophilus medusophagus and Petromyzon marinus, were recently collected in nearshore waters off the island of Malta in the central Mediterranean. Several morphological and biological parameters for the specimens are described. The findings ascertain the occurrence of the two species in the area and corroborate furthermore the substantial contribution to wildlife sightings made through social media and by citizen scientists, thus further increasing knowledge on Mediterranean biodiversity.

Keywords: Centrolophidae, Petromyzontidae, Malta, Mediterranean Sea, citizen science

PRESENZA CONFERMATA DI *SCHEDOPHILUS MEDUSOPHAGUS* (COCCO, 1839) E *PETROMYZON MARINUS* LINNAEUS, 1758 NELLE ACQUE MALTESI, MEDITERRANEO CENTRALE

SINTESI

Esemplari di due specie di pesci non comuni, Schedophilus medusophagus e Petromyzon marinus, sono stati recentemente raccolti nelle acque dell'isola di Malta nel Mediterraneo centrale. Alcuni parametri morfologici e biologici degli esemplari sono descritti. I ritrovamenti accertano la presenza di queste due specie nell'area, corroborando inoltre l'importante contributo che i social media e i citizen scientists danno per quanto concerne le osservazioni in natura e quindi per un ulteriore incremento della conoscenza della biodiversità del Mediterraneo.

Parole chiave: Centrolophidae, Petromyzontidae, Malta, Mediterraneo, citizen science

INTRODUCTION

In the Mediterranean Sea, four species of the family Centrolophidae are known: *Hyperoglyphe periformis* (Mitchill, 1818), *Centrolophus niger* (Gmelin, 1789), *Schedophilus ovalis* (Cuvier, 1833) and *Schedophilus medusophagus* (Cocco, 1839) (Kovačić et al., 2021). According to Borg et al. (2023), in the waters around the Maltese archipelago, the occurrence of *C. niger* and *S. ovalis* is well documented, while the presence of *S. medusophagus* requires confirmation.

In the same basin, two species of Petromyzontidae occur, *Lampetra fluviatilis* (Linnaeus, 1758) and *Petromyzon marinus* Linnaeus, 1758, of which only the latter species has been previously reported from the Maltese Islands (Borg et al., 2023).

Through the cooperation with citizen scientists, individuals of *S. medusophagus* and *P. marinus* were collected from Maltese waters. The main morphological features and meristic characters of the individuals were described. The present note confirms the occurrence of both species in Maltese waters and adds knowledge on the characteristic fish diversity in this area of the central Mediterranean Sea.

MATERIAL AND METHODS

On 10 February 2023, a slowly-swimming individual of *S. medusophagus* was scooped by a fisher from the surface at Pieta yacht marina, island of Malta (coordinates 35°53'38.21"N, 14°29'52.49"E), by means of a handnet.

On 23 March 2023, a dead individual of *P. marinus* was observed stranded on the beach, after a severe storm, at Fomm ir-Rih, along the north-west coast of the island of Malta (35°54'24.17"N; 14°20'27.90"E); the individual was collected by an angler.

Both fishes were unusual to those who collected them, such that they immediately contacted one of the authors (A.D.) through the "Spot the Alien Fish" citizen science platform on Facebook (<https://www.facebook.com/aliensmalta>), a campaign implemented since 2017 within the Department of Geosciences at the University of Malta.

The samples of *S. medusophagus* and *P. marinus* are currently preserved in the collection of the Department of Geosciences at the University of Malta under the catalogue numbers OMRG/GS/01/2023 and OMRG/GS/02/2023, respectively.

Abbreviations used: Total length, TL; Standard length, SL; Head length, HL.



Fig. 1: The de-frozen specimen of *Schedophilus medusophagus* collected in Malta in 2023 (scale bar = 10 cm). [Detail: freshly caught specimen].

Sl. 1: Odmrznjen primerek meduzojedca (*Schedophilus medusophagus*), ujetega na Malti v letu 2023 (merilo = 10 cm). [Detajl: sveže ujeti primerek].

RESULTS

Schedophilus medusophagus (Cocco, 1839)

The specimen was a juvenile with TL 145 mm and weight 39.3 g. Brief description (Fig. 1): body compressed and high; a single long dorsal fin, its origin before pectoral fin origin; snout short, slightly less than eye diameter; pectoral and pelvic fins inserted at the same level; caudal fin bilobate; lateral line relatively curved anteriorly, over the operculum and the pectoral fin. Dorsal fin rays III+46; Anal fin spines plus soft rays 32; Pectoral fin rays 19; Ventral fin rays 5; Caudal fin rays 22; gill rakers 11+1+6 on first gill arch; 15 spines on preoperculum. Body depth 47, head length 28.3, predorsal 25.4, preventral 31, preanal 54.6, all as % of SL; caudal peduncle length 52.2, eye diameter 21.6, preorbital distance 18.8, postorbital 59.1, all as % of HL (Tab. 1). Colour of fresh and de-frozen specimen were similar: background of body light blue; irregular darker grey patches and wavy horizontal stripes on sides, some patches extending at the base of dorsal and anal fins; pectoral and pelvic fins dark; a continuous dark stripe at the middle of the dorsal and anal fins (Fig. 1).

Petromyzon marinus (Linnaeus, 1758)

The lamprey was an adult with TL 505 mm and weight 360 g. Main morphological characters: body anguilliform, two dorsal fins in the posterior half of the body (Fig. 2A), the base of the first 65 mm long, the base of the second 142 mm; eye length 5.9 mm. Six branchial openings on the left side (Fig. 2B), seven on the right; branchial length 48 mm; interbranchial opening distance 10.3 mm. Oral disc (Fig. 2C): length 38.5 mm; teeth on concentric series; one bicuspid supraoral tooth, 4 endolateral bicuspid teeth on each side, bilobed lingual teeth, infraoral lamina with 8 unicuspid teeth. Colour: body mottled black dorsally and laterally, in a marbled pattern and uniformly pale ventrally.

DISCUSSION

The morphological and meristic characters as well as the colour of the Centrolophidae specimen under study were in full agreement with the description of *S. medusophagus* given by Tortonese (1959), Heidrich (1986), Fahay (2007) and Milana et al. (2011). Our specimen differed from its close relative *S. ovalis* by virtue of its soft and limp body (rigid and firm in *S. ovalis*) and based on differences in the dorsal fin count: *S. ovalis* is generally characterized by 30–32 rays in its dorsal fins, whilst our specimen displayed 46 dorsal fin rays (Heidrich, 1986; Rafrafi-Nouira et al., 2015a).

Tab. 1: Morphometric measurements (mm) of the *Schedophilus medusophagus* specimen from Malta.

Tab. 1: Morfometrične meritve (v mm) primerka meduzojedca (*Schedophilus medusophagus*) iz Malte.

Measurements	mm
Total length (TL)	145
Standard length (SL)	113
Maximum body depth	53
Minimum body depth	10.8
Caudal peduncle length	16.7
Head length (HL)	32
Eye diameter	6.9
Preorbital distance	6
Postorbital distance	18.9
Interorbital distance	12.6
Dorsal fin base length	88.1
Anal fin base length	42
Pectoral fin length	24
Ventral fin length	23.8
Predorsal length	28.7
Preventral length	35
Preanal length	61.7
Maximum height of dorsal fin	24.4
Maximum height of anal fin	14.8

The Cornish blackfish *S. medusophagus* is a mesopelagic fish encountered in the temperate waters of the north Atlantic and the Mediterranean, prevalently within the western half of the basin, but also in the Adriatic Sea and in the central Mediterranean, such as in Tunisia and within the Strait of Sicily (Bauchot, 1987; Dulčić, 1998; Dulčić & Lipej, 2002; Bradai et al., 2004; Bañón et al., 2012; Battaglia et al., 2014; Rafrafi-Nouira et al., 2015a; Hattour & Koched, 2017; Kovačić et al., 2020). Up to date, no records of the species are documented in the eastern side of the basin (Golani et al., 2006; Bilecenoğlu et al., 2014; Akel & Karachle, 2017; Ali, 2018; Bariche & Fricke, 2020; Golani, 2021) and its presence in Hellenic waters is considered



Fig. 2: The whole de-frozen specimen of *Petromyzon marinus* collected in Malta in 2023 (A), the abnormal six gill openings in the left side (B) (scale bars in A and B = 10 cm) and the oral disc, 38.5 mm in length (C).

Sl. 2: V celoti odmrznjen primerek morskega piškurja (*Petromyzon marinus*), ujetega na Malti v 2023 (A), nenavadnih šest škržnih rež na levi strani (B) (merilo za A in B = 10 cm) in oralni disk, 38,5 mm v dolžino (C).

questionable (Papaconstantinou, 2014). It reaches a maximum TL of 50 cm; adult specimens are mainly found at 300–900 m of depth. Juveniles are usually detected at the surface, as in the present case, frequently associated with jellyfish (Haedrich, 1986; Bauchot, 1987), especially *Pelagia noctiluca* (family Pelagiidae), that constitute the main food source for adults and juveniles of this species (Garibaldi et al., 2010; Battaglia et al., 2014).

In March and May 2022, during a bloom of *Pelagia noctiluca*, unknown centrolophids, possibly *S. medusophagus*, were repeatedly observed in the surface waters around the Maltese Islands, and photos were submitted to the above-mentioned Maltese platform (cf. Deidun et al., 2022), but samples were not retained. The Cornish blackfish *S. medusophagus* and the Imperial blackfish *S. ovalis* may be sometimes confused with each other, especially at juvenile stage, or may be confused with species of other families (see Nour et al., 2022; Borg et al., 2023). In the case reported in the present study, the availability of a sample allowed the authors to ascertain the identity of the same as belonging to *S. medusophagus* and to confirm the

occurrence of the species in Maltese waters. Therefore, *S. medusophagus* can be added to the list of 349 bony fish species that are to date confirmed as occurring in the same waters (Borg et al., 2023).

The morphological characters, dentition, and other associated structures of the oral disc, as well as the colour of the lamprey under study fully agreed with the description of *L. marinus* given by Bauchot (1987) and Renaud (2011). The latter cited literature and the work by Maitland (1972) allowed us to discard *L. fluviatilis* when identifying the specimen, by virtue of the marbled dark pattern on its back and along the sides (uniform colour pattern in *L. fluviatilis*), the occurrence of closely-packed teeth in radiating rows on the oral disc (teeth are widely-spaced in *L. fluviatilis*) as well as the occurrence of two large teeth in the supra-oral dental plate (one small tooth at most in *L. fluviatilis*). The number of branchial openings, six instead of seven on the left side of the body, was probably a morphological aberration. Six branchial openings on one side, generally on the left, as well as multiple tails, have been observed in rare cases in lampreys (Renaud, 2011;

Hume *et al.*, 2014). It is to be noted that in the *P. marinus* specimen from Syria reported in Saad *et al.* (2021), the first dorsal fin appears positioned in the anterior half of the body, while in the Petromyzontidae both dorsal fins are usually positioned in the posterior half. This could perhaps be another type of morphological abnormality in lampreys, although, to our knowledge, this is as yet unreported in the literature.

The sea lamprey *P. marinus* occurs at depths ranging between 1 m and 4000 m, in the marine, freshwater and brackish waters of the north Atlantic Ocean, the western and central Mediterranean Sea and in the Adriatic (Catalano *et al.*, 1997; Renaud, 2011; Milana *et al.*, 2011; Karachle & Machias, 2014; Rafrafi-Nouira *et al.*, 2015b; Tutman *et al.*, 2020; Giglio & Sperone, 2021; Froese & Pauly, 2023; Antognazza *et al.*, 2023). Scattered records of the species have also been reported from the eastern Ionian Sea (Karachle & Machias, 2014), the north Aegean (Economidis *et al.*, 1999; Papaconstantinou, 2014) and the southeastern Aegean Sea (Filiz *et al.*, 2012), as well as from the Levantine Sea waters of Turkey and Syria (Çevik *et al.*, 2010; Saad *et al.*, 2021).

Sea lampreys are parasitic on sharks, large bony fishes and marine mammals. The sample studied here probably has lost its host during the mentioned storm, and after death it was washed up on the beach.

The sea lamprey has been historically reported in coastal waters of the Maltese archipelago, but

there are no recent records (Borg *et al.*, 2023), suggesting that this species is uncommon. In the Tunisian coasts, close to the Maltese Islands, the species is recorded as an occasional visitor, and it is not considered to have established viable populations (Rafrafi-Nouira *et al.*, 2015b). The finding described in the present study indicates that *P. marinus* occasionally occurs in the wider area around the Maltese islands. Nevertheless, it is difficult to ascertain if there is a viable population of this anadromous species established within Maltese waters given the non-retrieval of live individuals but only of a stranded one; in addition, along the Maltese coastline, there is a lack of freshwater systems suitable for its reproduction in the area.

The GBIF database was consulted on the 24th of August 2023 and, although several Mediterranean Atlantic findings exist within the database for both *S. medusophagus* and for *P. marinus*, no specimens of these species are listed for Maltese waters.

ACKNOWLEDGEMENTS

Authors thank Mr Eric Zarb and Mr Kevin Gauci, the citizen scientists who kindly provided to the authors the *Schedophilus medusophagus* and *Petromyzon marinus* specimens respectively and shared information on their collection. They also are grateful to the anonymous reviewers for their useful comments on the manuscript.

POTRJEHO POJAVLJANJE MEDUZOJEDCA, *SCHEDOPHILUS MEDUSOPHAGUS* (COCCO, 1839), IN MORSKEGA PIŠKURJA, *PETROMYZON MARINUS* LINNAEUS, 1758, V MALTEŠKIH VODAH, OSREDNJE SREDOZEMSKO MORJE

Alan DEIDUN

Department of Geosciences, University of Malta, Msida MSD 2080, Malta
e-mail: alan.deidun@um.edu.mt

Bruno ZAVA

Wilderness studi ambientali, via Cruillas 27, 90146 Palermo, Italy
Museo Civico di Storia Naturale di Comiso, Via degli Studi 9, Comiso (Rg), Italy
e-mail: wildernessbz@hotmail.com

Alessio MARRONE

Department of Geosciences, University of Malta, Msida MSD 2080, Malta
e-mail: alessio.marrone@um.edu.mt

Johann GALDIES

Department of Geosciences, University of Malta, Msida MSD 2080, Malta
e-mail: johann.galdies@um.edu.mt

Arnold SCIBERRAS

The Exterminator Ltd. 5, The Service Hub, Triq San Gorg, Naxxar NXR 2541 Malta
e-mail: bioislets2@gmail.com

Maria CORSINI-FOKA

Institute of Oceanography, Hellenic Centre for Marine Research. Hydrobiological Station of Rhodes, Cos Street, 85100 Rhodes, Greece
e-mail: mcorsini@hcmr.gr

POVZETEK

V malteških vodah osrednjega Sredozemskega morja so avtorji pred kratkim ulovili primerke dveh manj pogostih vrst, meduzojedca (*Schedophilus medusophagus*) in morskega piškurja (*Petromyzon marinus*). Opisujejo številne morfološke in biološke parametre primerkov. Ugotovitve dokazujejo pojavljanje obeh vrst na obravnavanem območju in potrjujejo pomemben prispevek socialnih medijev in ljubiteljske znanosti, s čimer se dodatno povečuje poznavanje o biotski raznovrstnosti Sredozemskega morja.

Ključne besede: Centrolophidae, Petromyzontidae, Malta, Sredozemsko morje, ljubiteljska znanost

REFERENCES

- Akel, E.H.Kh. & P.K. Karachle (2017):** The Marine Ichthyofauna of Egypt. Egypt. J. Aquat. Biol. Fish., 21(3), 81-116.
- Ali, M.F. (2018):** An updated checklist of the marine fishes from Syria with emphasis on alien species. Mediterr. Mar. Sci., 19(2), 388-393. <https://doi.org/10.12681/mms.15850>.
- Antognazza, C.M., A. Gentile, G. Crosa & S. Zaccara (2023):** The presence of lampreys in the Tyrrhenian rivers of the Campania region (Southern Italy): A new record of the Sea Lamprey *Petromyzon marinus* (Linnaeus 1758). Environments, 10, 125. <https://doi.org/10.3390/environments10070125>.
- Bañón, R., A. Alonso-Fernández, D.T.G. Quigley, A. Miranda & J.C. Arronte (2012):** Unusual shallow inshore records of Cornish blackfish *Schedophilus medusophagus* (Stromateoidei: Centrolophidae) from Galician waters (NW Spain). Cah. Biol. Mar., 53, 271-277. <https://doi.org/10.21411/CBM.A.73E1719B>.
- Bariche, M. & R. Fricke (2020):** The marine ichthyofauna of Lebanon: an annotated checklist, history, biogeography, and conservation status. Zootaxa, 4775(1), 1-157. <https://doi.org/10.11646/zootaxa.4775.1.1>.
- Battaglia, P., S. Musolino, V. Esposito, G. Amendolia, P. Consoli, F. Andaloro & T. Romeo (2014):** Feeding habits of juvenile fishes belonging to three medusivorous species (Centrolophidae and Nomeidae) from the Strait of Messina (central Mediterranean Sea). Mar. Biol. Res., 10(9), 927-933. <http://dx.doi.org/10.1080/17451000.2013.863357>.
- Bauchot, M.L. (1987):** Lamproies et Myxines. In: Fischer, W., M.L. Bauchot & M. Schneider (eds.): Fiches FAO d'identification des espèces pour les besoins de la pêche. (Révision 1). Méditerranée et mer Noire. Zone de pêche 37. Volume 2. Vertébrés. FAO, Rome, pp. 762-765.
- Bilecenoğlu, M., M. Kaya, B. Cihangir & E. Çiçek (2014):** An updated checklist of the marine fishes of Turkey. Turk. J. Zool., 38, 901-929. <https://doi.org/10.3906/zoo-1405-60>.
- Borg, J.A., D. Dandria, J. Evans, L. Knittweis & P. J. Schembri (2023):** A critical checklist of the marine fishes of Malta and surrounding waters. Diversity, 15, 225. <https://doi.org/10.3390/d15020225>.
- Bradai, M.N., J.P. Quignard, A. Bouain, O. Jarboui, A. Ouannes-Ghorbel, L. Ben Abdallah, J. Zaouali & S. Ben Salem (2004):** Ichtyofaune autochtone et exotique des côtes tunisiennes: recensement et biogéographie. Cybium, 28(4), 315-328. <https://doi.org/10.26028/cybium/2004-284-003>.
- Catalano, E., P. Gianguzza & B. Zava (1997):** Sulla presenza di *Petromyzon marinus* Linnaeus, 1758 (Cyclostoata, Petromyzonidae) in acque siciliane. Biol. Mar. Medit., 4(1), 513-514.
- Çevik, C., D. Ergüden & N. Tekelioğlu (2010):** Confirmation of the presence of the sea lamprey, *Petromyzon marinus* Linnaeus, 1758 in the Levantine Sea (Petromyzoniformes: Petromyzonidae). Zool. Middle East, 49, 107-108. <https://doi.org/10.1080/09397140.2010.10638398>.
- Deidun, A., B. Zava & M. Corsini-Foka (2022):** Distribution extension of *Lutjanus argentimaculatus* (Forsskal, 1775) (Lutjanidae) and *Psenes pellucidus* Lütken, 1880 (Nomeidae) to the waters of Malta, central Mediterranean Sea. Annales, Ser. Hist. Nat., 32(1), 49-58. <https://doi.org/10.19233/ASHN.2022.06>.
- Dulčić, J. (1998):** First record of the cornich blackfish *Schedophilus medusophagus* (Pisces: Centrolophidae) larvae from the Adriatic Sea. J. Mar. Biol. Assoc. U. K., 78(3), 1035-1038. <https://doi.org/10.1017/S002531540004501X>.
- Dulčić, J. & L. Lipej (2002):** Rare and little-known fishes in the Eastern Adriatic during last two decades (1980-2001). Periodicum Biologorum, 104(2), 185-194.
- Economidis, P.S., A. Kallianiotis & H. Psaltopoulou (1999):** Two records of sea lamprey from the north Aegean Sea. J. Fish Biol., 55, 1114-1118. <https://doi.org/10.1111/j.1095-8649.1999.tb00745.x>.
- Fahay, M.P. (2007):** Early Stages of Fishes in the Western North Atlantic Ocean. Volume Two. Scorpaeniformes through Tetraodontiformes. Northwest Atlantic Fisheries Organization, 2007. Available at: <https://www.nafo.int/Library/Fahay-Early-Stages-of-Fishes/Early-Stages-of-Fishes-PDFs> (accessed on 20-4-2023).
- Filiz, H., G. Bilge & M. Erdem (2012):** New occurrence record of the lamprey in the southern Aegean Sea. pp. 319-320. In: Thessalou-Legaki, M., O. Aydogan, P. Bekas, G. Bilge, Y.O. Boyaci, E. Brunelli, V. Circosta, F. Crocetta, F. Durucan, M. Erdem, A. Ergolavou, H. Filiz, F. Fois, E. Gouva, K. Kapisir, S. Katsanevakis, Z. Kljajić, E. Konstantinidis, G. Konstantinou, D. Koutsogiannopoulos, S. Lamon, V. Mačić, R. Mazzette, D. Meloni, A. Mureddu, I. Paschos, C. Perdikaris, F. Piras, D. Poursanidis, A.A. Ramos-Esplà, A. Rosso, P. Sordino, E. Sperone, A. Sterioti, E. Taşkin, F. Toscano, S. Tripepi, L. Tsiakiros & A. Zenetos: New Mediterranean biodiversity records (December 2012). Mediterr. Mar. Sci., 13, 312-327. <https://doi.org/10.12681/mms.313>.
- Froese, R. & D. Pauly (2023):** FishBase. World Wide Web electronic publication. www.fishbase.org (Accessed on 26-4-2023).
- Garibaldi, F., L. Orsi Relini & G. Relini (2010):** Medusivorous fishes of the Ligurian Sea 2. The specialist, *Schedophilus medusophagus*, Cocco 1839. Rapp. Comm. int. Mer Médit., 39, 525.

- Giglio, G. & E. Sperone (2021):** First documented record of *Petromyzon marinus* Linnaeus, 1758 in Calabria (Central Mediterranean, Southern Italy). p. 208. In: Santín, A., R. Aguilar, O. Akyol, C.R. Begburs, L. Benoit, G. Chimienti, F. Crocetta, C. Dalyan, A. De La Linde Rubio, B. Dragičević, J. Dulčić, G. Giglio, O. Gönülal, T. Kebapcioglu, N.B. Kesici, S. Kiparissis, V. Kousteni, E. Mancini, F. Mastrototaro, T. Menut, F. Montesanto, P. Peristeraki, D. Poursanidis, J. Renoult, L. Sánchez-Tocino, E. Sperone & F. Tiralongo: New records of rare species in the Mediterranean Sea (March 2021). *Mediterr. Mar. Sci.*, 22(1), 199-217. <https://doi.org/10.12681/mms.25295>.
- Golani, D. (2021):** An updated Checklist of the Mediterranean fishes of Israel, with illustrations of recently recorded species and delineation of Lessepsian migrants. *Zootaxa*, 4956(1), 1-108. <https://doi.org/10.11646/zootaxa.4956.1.1>.
- Golani, D., B. Oztürk & N. Başusta (2006):** The Fishes of the Eastern Mediterranean. Turkish Marine Research Foundation, Istanbul, Turkey, 259 pp.
- Hattour, A. & W. Koched (2017):** Nouvelles occurrence et signalisation d'espèces de la famille des Centrolophidae (Actinopterygiens, Perciformes) dans les eaux Tunisiennes. *Bulletin de l'Institut National des Sciences et Technologies de la Mer de Salammbô*, 44, 195-202.
- Heidrich, R.L. (1986):** Centrolophidae. In: Whitehead, P. J. P., M. L. Bauchot, J. C. Hureau, J. Nielsen & E. Tortonese (eds.): Check-list of the fishes of the north-eastern Atlantic and of the Mediterranean (CLOFNAME), Volume 3. UNESCO, Paris, pp. 1177-1182.
- Hume, J.B., C.W. Bean & C.E. Adams (2014):** Morphological abnormalities in a population of *Lampetra planeri*, with a short review of petromyzontid teratologies. *J. Fish Biol.*, 84, 1614-1619. <https://doi.org/10.1111/jfb.12369>.
- Karachle, P.K. & A. Machias (2014):** First confirmed record of *Petromyzon marinus* Linnaeus, 1758 from west Greece. *Cah. Biol. Mar.*, 55(3), 367-370. <https://doi.org/10.21411/CBM.A.969D4E94>.
- Kovačić, M., L. Lipej & J. Dulčić (2020):** Evidence approach to checklists: critical revision of the checklist of the Adriatic Sea fishes. *Zootaxa*, 4767(1), 1-55. <https://doi.org/10.11646/zootaxa.4767.1.1>.
- Kovačić, M., L. Lipej, J. Dulčić, S.P. Iglesias & M. Goren (2021):** Evidence-based checklist of the Mediterranean Sea fishes. *Zootaxa*, 4998(1), 1-115. <https://doi.org/10.11646/zootaxa.4998.1.1>.
- Maitland, P.S. (1972):** A key to the freshwater fishes of the British Isles with notes on their distribution and ecology. Freshwater Biological Association Scientific Publication No. 27, 139 pp.
- Milana, V., A. Fusari, A. R. Rossi & L. Sola (2011):** Molecular and morphological identification of an uncommon centrolophid fish. *Cent. Eur. J. Biol.*, 6(3), 440-445. <https://doi.org/10.2478/s11535-011-0016-x>.
- Nour, O.M., S.A.A. Al Mabruk, B. Zava, P. Gianguzzza, M. Corsini-Foka & A. Deidun (2022):** First record of *Naso annulatus* (Quoy & Gaimard, 1825) and further records of *Scatophagus argus* (Linnaeus, 1766) and *Charybdis (Charybdis) natator* (Herbst, 1794) in the Mediterranean Sea. *Biol. Invasions Rec.*, 11(3), 785-795. <https://doi.org/10.3391/bir.2022.11.3.20>.
- Papaconstantinou, C. (2014):** Fauna Graeciae. An updated checklist of the fishes in the Hellenic Seas. Monographs on Marine Sciences, 7. Hellenic Centre for Marine Research (HCMR), Athens, 340 pp.
- Rafrafi-Nouira, S., D. Golani, O. El Kamel-Moutalibi, M. Boumaiza, C. Reynaud & C. Capapé (2015a):** First record of imperial blackfish, *Schedophilus ovalis* (Actinopterygii: Perciformes: Centrolophidae), from the Tunisian coast, central Mediterranean. *Acta Ichthyol. Piscat.*, 45(2), 203-206. <https://doi.org/10.3750/AIP2015.45.2.11>.
- Rafrafi-Nouira, S., O. El Kamel-Moutalibi, M. Boumaiza, C. Reynaud & C. Capapé (2015b):** On the occurrence of sea lamprey *Petromyzon marinus* (Agnatha: Petromyzonidae) in Tunisian waters (Central Mediterranean Sea). *Cah. Biol. Mar.*, 56, 73-76. <https://doi.org/10.21411/CBM.A.42E4EC16>.
- Renaud, C.B. (2011):** Lampreys of the world. An annotated and illustrated catalogue of lamprey species known to date. FAO Species Catalogue for Fishery Purposes. No. 5. FAO, Rome, 109 pp.
- Saad, A., I. Barakat, M. Masri, W. Sabour & C. Capapé (2021):** First substantiated record of sea lamprey *Petromyzon marinus* (Agnatha: Petromyzontidae) from the Syrian coast (Eastern Mediterranean Sea). *FishTaxa*, 20, 21-24.
- Tortonese, E. (1959):** Revisione dei Centrolophidae (Pisces Perciformes) del Mare Ligure. I–Annali Mus. civ. St. nat., Genova, 71, 57-82.
- Tutman, P., I. Buj, M. Čaleta, Z. Marčić, A. Hamzić & A. Adrović (2020):** Review of the lampreys (Petromyzontidae) in Bosnia and Herzegovina: a current status and geographic distribution. *J. Vertebr. Biol.*, 69(1), 19046. <https://doi.org/10.25225/jvb.19046>.