

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 32, 2022, 1





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 32, 2022, 1

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 32, leto 2022, številka 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačič (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®

Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®

**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran / Pirano, Fornace/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 30. 06. 2022.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS) in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation:

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY-NC 4.0.



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>



VSEBINA / INDICE GENERALE / CONTENTS 2022(1)

SREDOZEMSKI MORSKI PSI
SQUALI MEDITERRANEI
*MEDITERRANEAN SHARKS***Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ**

Observations on Thresher Shark, *Alopias vulpinus* (Chondrichthyes: Alopiidae) from the Coast of Algeria (Southwestern Mediterranean Sea) 1
Opazovanja morskih lisic, Alopias vulpinus (Chondrichthyes: Alopiidae) ob alžirski obali (jugozahodno Sredozemsko morje)

Elif ÖZGÜR ÖZBEK & Hakan KABASAKAL

Notes on Smoothback Angel Shark, *Squatina oculata* (Squatiniformes: Squatinidae) caught in the Gulf of Antalya 9
Zapis o pegastih sklatih, Squatina oculata (Squatiniformes: Squatinidae), ujetih v Antalijskem zalivu

Alessandro PAGANO & Alessandro DE MADDALENA

Underwater Observations of the Rare Angular Roughshark *Oxynotus centrina* (Chondrichthyes: Squalidae) in the Waters of Santa Tecla (Sicily, Italy) 17
Podvodna opazovanja redkega morskega prašiča, Oxynotus centrina (Chondrichthyes: Squalidae) v vodah Sante Tecele (Sicilija, Italija)

Deniz ERGÜDEN, Deniz AYAS & Hakan KABASAKAL

Morphometric Measurements of Three Young Carcharhinid Species from Northeastern Levant (Mediterranean Sea) 25
Morfometrične meritve mladičev treh vrst morskih psov iz družine Carcharhinidae iz severnovzhodnega Levanta (Sredozemsko morje)

Hakan KABASAKAL

Projections on the Future of Deep-Sea Sharks in the Sea of Marmara, Where Deep Zones Are Threatened by Deoxygenation: a Review 35
Napovedi o prihodnosti globomorskih morskih psov v Marmarskem morju, ogroženem zaradi pomanjkanja kisika: pregled

BIOINVAZIJA
BIOINVASIONE
*BIOINVASION***Alan DEIDUN, Bruno ZAVA & Maria CORSINI-FOKA**

Distribution Extension of *Lutjanus argentimaculatus* (Lutjanidae) and *Psenes pellucidus* (Nomeidae) to the Waters of Malta, Central Mediterranean Sea 49
Širjenje areala vrst Lutjanus argentimaculatus (Lutjanidae) in Psenes pellucidus (Nomeidae) v malteške vode (osrednje Sredozemsko morje)

Sami M. IBRAHIM, Abdulrazziq A. ABDULRAZIQ, Abdulghani ABDULGHANI, Sara A.A. AL MABRUK, David SALVATORI, Bruno ZAVA, Maria CORSINI-FOKA & Alan DEIDUN

First Record of *Enchelycore anatina* (Muraenidae) from Libyan Waters and an Additional Record from Southern Italy (Western Ionian Sea) 59
Prvi zapis o pojavljanju kavljazobe murene Enchelycore anatina (Muraenidae) iz libijskih voda in dodatni zapis za južno Italijo (zahodno Jonsko morje)

Rasha Ali HENEISH & Samir Ibrahim RIZKALLA

Morphometric and Meristic Characteristics of a New Record of Bluespot Mullet *Crenimugil seheli* (Pisces: Mugilidae) in Egyptian Mediterranean waters 67
Novi zapis o pojavljanju vrste Crenimugil seheli (Pisces: Mugilidae) v egiptovskih sredozemskih vodah in njene morfolometrične in meristične značilnosti

Yana SOLIMAN, Adib SAAD, Vienna HAMMOUD & Christian CAPAPÉ

Heavy Metal Concentrations in Tissues of *Siganus rivulatus* (Siganidae) from the Syrian Coast (Eastern Mediterranean Sea) 75
Vsebnost težkih kovin v tkivih marmoriranega morskega kunca Siganus rivulatus (Siganidae) iz sirske obale (vzhodno Sredozemsko morje)

IHTIOLOGIJA
 ITTIOLOGIA
 ICHTHYOLOGY

Jihade ALAHYENE, Brahim CHIAHOU, Hammou EL HABOUZ & Abdelbasset BEN-BANI

Length Based Growth Estimation of the Blue Shark *Prionace glauca* from the Moroccan Central Atlantic Coast 85
Dolžinsko-masni odnos in ocena rasti pri sinjem morskem psu (Prionace glauca) iz osrednje atlantske obale Maroka

Okan AKYOL, Altan LÖK & Funda ERDEM

Occurrence of *Cubiceps gracilis* (Nomeidae) in the Eastern Mediterranean Sea 101
Pojavljanje klateža, Cubiceps gracilis (Nomeidae), v vzhodnem Sredozemskem morju

Farid HEMIDA, Boualem BRAHMI, Christian REYNAUD & Christian CAPAPÉ

Occurrence of the Rare Driftfish *Cubiceps gracilis* (Nomeidae) from the Algerian Coast (Southwestern Mediterranean Sea) 107
Pojavljanje redkega klazeža Cubiceps gracilis (Nomeidae) z alžirske obale (jugozahodno Sredozemsko morje)

Deniz ERGÜDEN & Cemal TURAN

A Rare Occurrence of *Carapus acus* (Carapidae) in the Eastern Mediterranean, Turkey 113
Redko pojavljanje strmorinca Carapus acus (Carapidae) v vzhodnem Sredozemskem morju (Turčija)

Laith JAWAD, Murat ŞIRIN, Miloslav PETRTÝL, Ahmet ÖKTENER, Murat ÇELİK & Audai QASIM

Skeletal Abnormalities in Four Fish Species Collected from the Sea of Marmara, Turkey 119
Skeletne anomalije pri štirih vrstah rib iz Marmarskega morja (Turčija)

RAZMNOŽEVALNA EKOLOGIJA
 ECOLOGIA RIPRODUTTIVA
 REPRODUCTIVE ECOLOGY

Amaria Latefa BOUZIANI, Khaled RAHMANI, Samira AIT DARNA, Alae Eddine BELMAHI, Sihem ABID KACHOUR & Mohamed BOUDERBALA

Gonadal Histology in *Diplodus vulgaris* from the West Algerian Coast 137
Histologija gonad pri navadnem šparu (Diplodus vulgaris) iz zahodne alžirske obale

Cheikhna Yero GANDEGA, Nassima EL OMRANI, Rezan O. RASHEED, Mohammed RAMDANI & Roger FLOWER

The Growth and Reproduction of Two Sparidae, *Pagrus caeruleostictus* and *Pagellus bellottii* in Northern Mauritanian Waters (Eastern Tropical Atlantic) 143
Rast in razmnoževanje dveh vrst pagrov, Pagrus caeruleostictus in Pagellus bellottii v severnih mavretanskih vodah (vzhodni tropski Atlantik)

Nassima EL OMRANI, Hammou EL HABOUZ, Abdellah BOUHAIMI, Jaouad ABOU OUALID, Abdellatif MOUKRIM, Jamila GOUZOU, Mohammed RAMDANI, Roger FLOWER & Abdelbasset BEN-BANI

The Reproductive Biology of the Pouting *Trisopterus luscus* from the Atlantic Coast of Morocco 155
Reproduktivna biologija francoskega moliča (Trisopterus luscus) iz atlantske obale Maroka

Mourad CHÉRIF, Rimel BENMESSAOUD & Christian CAPAPÉ

Growth Patterns and Age Structure of *Mullus surmuletus* (Mullidae) from the Northern Coast of Tunisia (Central Mediterranean Sea) 173
Rastni parametri in starostna struktura progastih bradačev Mullus surmuletus (Mullidae) iz severne tunizijske obale (osrednje Sredozemsko morje)

FLORA

FLORA

FLORA

Martina ORLANDO-BONACA, Erik LIPEJ, Romina BONACA & Leon Lojze ZAMUDA

Improvement of the Ecological Status of the *Cymodocea nodosa* Meadow near the Port of Koper 185
Izboljšanje ekološkega stanja morskega travnika kolenčaste cimodoceje (Cymodocea nodosa) v bližini kopskega pristanišča

FAVNA

FAVNA

FAVNA

Manja ROGELJA, Martin VODOPIVEC & Alenka MALEJ

Cestum veneris Lesueur, 1813 (Ctenophora) – a Rare Guest in the Northern Adriatic Sea 197
Cestum veneris Lesueur, 1813 (Ctenophora) – redek gost v severnem Jadranu

Adla KAHRIĆ, Dalila DELIĆ & Dejan KULIJER

Notospermus annulatus (Nemertea: Lineidae), a New Record for Bosnia and Herzegovina 205
Notospermus annulatus (Nemertea: Lineidae), prvi zapis o pojavljanju za Bosno in Hercegovino

Andrea LOMBARDO & Giuliana MARLETTA

Report of an Interesting *Trapania* (Gastropoda: Nudibranchia: Goniodorididae) Specimen from Central Eastern Sicily 211
Zapis o zanimivem primerku iz rodu Trapania (Gastropoda: Nudibranchia: Goniodorididae) iz osrednje vzhodne Sicilije

Abdelkarim DERBALI & Othman JARBOUI

Stock Assessment, Cartography and Sexuality of the Wedge Clam *Donax trunculus* in the Gulf of Gabes (Tunisia) 217
Ocena staleža, kartografija in spolnost klinaste školjke Donax trunculus v gabeškem zalivu (Tunizija)

Abdelkarim DERBALI, Aymen HADJ TAIEB & Othman JARBOUI

Length-Weight Relationships and Density of Bivalve Species in the Shellfish Production Area of Zarzis (Tunisia, Central Mediterranean Sea) 229
Dolžinsko-masni odnos in gostota školjk na gojišču školjk v predelu Zarsisa (Tunizija, osrednje Sredozemsko morje)

Toni KOREN

The Diversity of Moths (Lepidoptera: Heterocera) of Significant Landscape Donji Kamenjak and Medulin Archipelago, Istria, Croatia 237
Raznolikost nočnih metuljev (Lepidoptera: Heterocera) Pomembne pokrajine Donji Kamenjak in Medulinski arhipelag, Istra, Hrvaška

OCENE IN POROČILA

RECENSIONI E RELAZIONI

REVIEWS AND REPORTS

Ines Mandić Mulec & Nives Ogrinc

Recenzija knjige: Mikrobn biogeokemija vod 263

Kazalo k slikam na ovitku 265

Index to images on the cover 265

received: 2022-02-17

DOI 10.19233/ASHN.2022.03

UNDERWATER OBSERVATIONS OF THE RARE ANGULAR ROUGHSHARK *OXYNOTUS CENTRINA* (CHONDRICHTHYES: SQUALIDAE) IN THE WATERS OF SANTA TECLA (SICILY, ITALY)

Alessandro PAGANO

Via Duca degli Abruzzi 40, 95127 Catania, Italy
e-mail: alessandro.pagano@tim.it

Alessandro DE MADDALENA

Shark Museum, 26 Forest Hill Road, Simon's Town, 7975 Cape Town, South Africa
e-mail: alessandrodemaddalena@gmail.com

ABSTRACT

*Three encounters with the uncommon and elusive angular roughshark, *Oxynotus centrina*, are reported from the waters off Santa Tecla, Catania, Italy, in the Western Ionian Sea. The three observations occurred on 11 June 2019, 26 February 2020 and 26 January 2022, between 74 and 84 m depth. The sharks encountered were three different female individuals. Photographs of the sharks were taken on all the encounters.*

Key words: angular roughshark, *Oxynotus centrina*, Sicily, Ionian Sea, Mediterranean Sea

OSSERVAZIONI SUBACQUEE DEL RARO PESCE PORCO *OXYNOTUS CENTRINA* (CHONDRICHTHYES: SQUALIDAE) NELLE ACQUE DI SANTA TECLA (SICILIA, ITALIA)

SINTESI

*Vengono riportati tre incontri con il poco comune ed elusivo pesce porco, *Oxynotus centrina*, nelle acque di Santa Tecla, a Catania, in Italia, nel Mar Ionio Occidentale. Le tre osservazioni hanno avuto luogo l'11 giugno 2019, il 26 febbraio 2020 e il 26 gennaio 2022, tra i 74 e gli 84 m di profondità. Gli squali incontrati erano tre diversi individui, di lunghezza stimata di 75, 60 e 50 cm rispettivamente, tutti di sesso femminile. La documentazione fotografica è stata raccolta durante tutti gli incontri.*

Parole chiave: pesce porco, *Oxynotus centrina*, Sicilia, Mar Ionio, Mare Mediterraneo

INTRODUCTION

The angular roughshark, *Oxynotus centrina* (Linnaeus, 1758), belongs to the Order Squaliformes and the family Oxynotidae. The morphology of this species includes: no anal fin, broad dorsal fins with first dorsal fin origin over pectoral fin origin, fin spines on both dorsal fins, short pectoral fins, moderately long caudal fin upper lobe and short lower lobe, no caudal fin posterior notch, stout body, ventral ridges, large dermal denticles, short snout, small mouth with enlarged labial parts, large nostrils, large eyes, large spiracles, and five pairs of short gill slits. The coloration of the dorsal surfaces is grey-brown or blackish with a lighter pattern on the lateral part of the head, sides, caudal peduncle and caudal fin; the ventral surfaces are partially lighter. Its size at birth is 21–24 cm and the maximum size is 150 cm. The embryonic development of this species is aplacental viviparous, with unknown gestation and litter size of 7–23 young. It is a benthic species that lives on continental shelves

and upper slope, at depths ranging from 50 m to at least 725 m, with nocturnal habits, feeding on bony fishes, small-spotted catshark eggs, crustaceans, and polychaetes (Bigelow & Schroeder, 1948; Bass *et al.*, 1976; Compagno, 1984; De Maddalena *et al.*, 2015).

It is found in the Eastern Atlantic Ocean and its presence throughout the Mediterranean Sea is well documented, despite being an uncommon species in the entire area, and very rare in most sites (Tortonese, 1956; Capapé, 1977; Cadenat & Blache, 1981; Bauchot, 1987; Vanni, 1992; Moreno, 1995; Mizzan, 1994; Barrull *et al.*, 1999; Barrull & Mate, 2001; Barrull & Mate, 2002; Dulvy *et al.*, 2003; Lipej *et al.*, 2004; Kabasakal, 2010; De Maddalena *et al.*, 2015; Kousteni & Megalofonou, 2016; Koehler, 2018; Capapé *et al.*, 2021; Gajić *et al.*, 2021).

Keeping in mind the paucity of observation of live individuals of this species (Kabasakal, 2009), it was considered important to report on recent underwater observations that occurred in the waters of the Western Ionian Sea, off the Eastern Sicilian coast.



Fig. 1: The estimated 75 cm TL angular roughshark, *Oxynotus centrina*, observed off Santa Tecla, Catania, on the Eastern Coast of Sicily, Italy, on 11 June 2019 at 10:16, at 74 m depth (photo: A. Pagano).

Sl. 1: Približno 75 cm dolga samica morskega prašiča, *Oxynotus centrina*, opažena 11. junija 2019 ob 10:16, na globini 74 m ob Santa Tecli, Catania, na vzhodni obali Sicilije v Italiji (foto: A. Pagano).

Tab. 1: Main details of the three encounters with angular roughsharks, *Oxynotus centrina*, reported in this article.**Tab. 1: Glavne podrobnosti o treh srečanjih z morskimi prašiči, *Oxynotus centrina*, o katerih avtorja poročata v prispevku.**

Date	Location	Depth (m)	Sex	Estimated TL
11 June 2019	Santa Tecla, Catania, Italy	74	F	75 cm
26 February 2020	Santa Tecla, Catania, Italy	84	F	60 cm
26 January 2022	Santa Tecla, Catania, Italy	84	F	50 cm

MATERIAL AND METHODS

The first author experienced three encounters with angular roughsharks while scuba diving with another diver, Eugenio Longo, by Closed Circuit Re-breather in the waters of the Western Ionian Sea, off Santa Tecla, Catania, on the Eastern Coast of Sicily, Italy (latitude: 37.639682105936735 N, longitude: 15.183723565445055 E). The weather conditions

were good, with calm sea. The dive site is located 150 m from the shore, where the bottom depth is between 80 and 110 m. The site is a rocky landslide that ends on a sandy bottom. The average temperature of the water was 14°C on all three encounters. The first author dives about 50 times per year on the observation site, with an average of one dive per week.

Sharks were photographed using a Nikon D800 reflex camera with a Tokina 10-17 mm lens for subse-



Fig. 2: The estimated 60 cm TL angular roughshark observed on the same site on 26 February 2020 at 10:26, at 84 m depth (photo: A. Pagano).

Sl. 2: Približno 60 cm dolga samica morskega prašiča, *Oxynotus centrina*, opažena na isti lokaliteti 26. februarja 2020 ob 10:26, na globini 84 m (foto: A. Pagano).

quent analyses of their morphology and photoidentification of the individuals occurring in the area.

RESULTS AND DISCUSSION

On each encounter an individual female angular roughshark was observed (Tab. 1). The first encounter occurred on the morning of 11 June 2019, at 10:16, when the divers were at 74 m depth and the angular roughshark observed was estimated at 75 cm total length (TL). The second encounter occurred on the morning of 26 February 2020, at 10:26, when the divers were at 84 m depth, only 30 m from the site of the first encounter. This second shark was estimated at 60 cm total length. The third encounter occurred in the morning of 26 January 2022, at 11:26, once again when the divers were at 84 m depth, exactly in the same site of the second encounter. This third individual was estimated at 50 cm total length. Each encounter lasted 10 to 20 minutes.

Other species observed on the three encounters included the swallowtail seaperch, *Anthias anthias*

(Linnaeus, 1758), the serpent eel, *Ophisurus serpens* (Linnaeus, 1758), the yellow gorgonian, *Eunicella cavolinii* (Koch, 1887), the small polyped gorgonian, *Paramuricea clavata* (Risso, 1826), and the red coral, *Corallium rubrum* (Linnaeus, 1758).

Based on the shape of the dorsal fins, the colour pattern and the size, it is evident that the angular roughsharks encountered were three different individuals. The total length of the sharks was estimated by comparison with the size of the underwater housing of the camera. The sharks initially kept a distance from the divers, apparently disturbed by the light, but then let the divers approach them at very close range and to take close-up pictures of the subjects. Therefore, photographic documentation of the shark was taken on all three encounters.

Eco-tourism, including the activities of shark diving are crucial to draw public attention to the importance of protecting sharks (De Maddalena & Galli, 2017). Currently the Mediterranean Sea offers very few spots where it is possible to have the chance to dive with some species of sharks. Among those documented spots are



Fig. 3: The estimated 50 cm TL angular roughshark observed on the same site on 26 January 2022, at 11:26, at 84 m depth (photo: A. Pagano).

Sl. 3: Približno 50 cm dolga samica morskega prašiča, *Oxynotus centrina*, opažena na isti lokaliteti 26. januarja 2022 ob 11:26, na globini 84 m (foto: A. Pagano).

the Messina Strait, Italy, for the bluntnose sixgill shark, *Hexanchus griseus* (Bonnaterre, 1788) (Celona *et al.*, 2005), the offshore waters of Beirut, Lebanon, for the smalltooth sand tiger, *Odontaspis ferox* (Risso, 1810) (Barrull & Mate, 2002; De Maddalena *et al.*, 2015), Boncuk Bay, Turkey, Lampione, Italy, Hadera and Ashkelon, Israel, for the sandbar shark, *Carcharhinus plumbeus* (Nardo, 1827), and Hadera and Ashkelon, Israel, for the dusky shark, *Carcharhinus obscurus* (Le Sueur, 1818) (Barash *et al.*, 2018; Zemah Shamir *et al.*, 2019; Kabasakal, 2020; Cattano *et al.*, 2021). The fact that there are so few spots where it is possible to shark dive in the Mediterranean area, makes the site where the three observations of angular roughsark are described in this article even more important. The fact that the species is so uncommon in the entire Mediterranean makes the observation site unique and its preservation critical.

Since female angular roughsharks attain sexual maturity between 66 and 75 cm TL (De Maddalena *et al.*, 2015), we can assume that the female sharks observed off Santa Tecla were sexually mature or subadults, and speculate that perhaps their presence in the area

may be related to mating or parturition. Unfortunately, the observation area, despite being also a site for the reproduction of catsharks, *Scyliorhinus* sp., (as clearly shown by egg cases observed by the first author), is not protected in any way. Fishing with gillnets, longlines and pots is still common practice and unregulated. The species is endangered according to the assessment of the IUCN red list (IUCN, 2022) and it's apparently extinct from some locations including the Adriatic Sea and the Gulf of Lion. We strongly recommend the angular roughshark be listed as a protected species, and the site be properly preserved and closely monitored to ensure viable habitat remains.

ACKNOWLEDGEMENTS

The authors wish to thank Eric Glenn Haenni for taking the time to edit the manuscript, and the anonymous referees for their useful comments on the manuscript. Alessandro Pagano thanks Massimo Ardizzoni and Eugenio Longo. Alessandro De Maddalena thanks Alessandra, Antonio and Phoebe for their support and love.

PODVODNA OPAZOVANJA REDKEGA MORSKEGA PRAŠIČA, *OXYNOTUS CENTRINA*
(CHONDRICHTHYES: SQUALIDAE) V VODAH SANTE TECLE (SICILIJA, ITALIJA)

Alessandro PAGANO

Via Duca degli Abruzzi 40, 95127 Catania, Italy
e-mail: alessandro.pagano@tim.it

Alessandro DE MADDALENA

Shark Museum, 26 Forest Hill Road, Simon's Town, 7975 Cape Town, South Africa
e-mail: alessandrodemaddalena@gmail.com

POVZETEK

Avtorja poročata o treh srečanjih z nenavadnim in izmuzljivim morskim prašičem, *Oxynotus centrina*, v bližini Sante Tecla (Catania, Italija), v zahodnem Jonskem morju. Morskega prašiča so opazovali 11. junija 2019, 26. februarja 2020 in 26. januarja 2022, med 74 in 84 m globine. Bile so tri različne samice, vse tudi fotografirane.

Ključne besede: morski prašič, *Oxynotus centrina*, Sicilija, Jonsko morje, Sredozemsko morje

REFERENCES

- Barash, A., R. Pickholtz, E. Pickholtz, L. Blaustein & G. Rilov (2018):** Seasonal aggregations of sharks near coastal power plants in Israel: An emerging phenomenon. *Mar. Ecol. Prog. Ser.*, 590, 145-154. DOI: 10.3354/meps12478.
- Barrull, J. & I. Mate (2001):** First confirmed record of Angular Roughshark *Oxynotus centrina* (Linnaeus, 1758) predation on shark egg case of small-spotted catshark *Scyliorhinus canicula* (Linnaeus, 1758) in Mediterranean waters. *Annales, Ser. Hist. Nat.*, 11(1), 23-28.
- Barrull, J. & I. Mate (2002):** Tiburones del Mediterráneo. Librería El Set-ciències, Arenys de Mar, 292 pp.
- Barrull, J., I. Mate & M. Bueno (1999):** Observaciones de tiburones (Chondrichthyes, Euselachii) en aguas de Cataluña (Mediterráneo NO), con algunos aspectos generales de su ecología. *Sci. gerund.*, 24, 127-151.
- Bass, A.J., J.D. D'Aubrey & N. Kistnasamy (1976):** Sharks of the east coast of Southern Africa. 6. The families Oxynotidae, Squalidae, Dalatiidae and Echinorhinidae. *Invest. Rep. Oceanogr. Res. Inst.*, Durban, 45, 103 pp.
- Bauchot, M.L. (1987):** Requins. In: Fischer W., M. Schneider & M.-L. Bauchot (eds.): Fiches FAO d'identification des espèces pour les besoins de la pêche. (Révision I). Méditerranée et Mer Noire. Zone de pêche 37. Vol. 2. Vertébrés. CEE, FAO, Rome, pp. 767-843.
- Bigelow, H.B. & W.C. Schroeder (1948):** Sharks. In: Fishes of the Western North Atlantic. Part one: Lancelets, Ciclostomes, Sharks. Memoir Sears Foundation for Marine Research, Yale University, pp. 53-576.
- Cadenat, J. & J. Blache (1981):** Requins de Méditerranée et d'Atlantique (plus particulièrement de la Côte Occidentale d'Afrique). *Faune Tropicale*, ORSTOM, Paris, 21, 1-330.
- Capapé, C. (1977):** Liste commentée des sélachiens de la région de Toulon (de La Ciotat à Saint-Tropez). *Bull. Mus. Hist. Nat. Marseille*, 37, 5-9.
- Capapé, C., S. Rafrari-Nouira, M.M. Ben Amor, K. Ounifi-Ben Amor & C. Reynaud (2021):** *Oxynotus centrina* from the Tunisian coast (Mediterranean Sea). *Thalassia Sal.*, 43, 35-42. DOI: 10.1285/i15910725v43p35.
- Cattano, C., G. Turco, M. Di Lorenzo, M. Gristina, G. Visconti & M. Milazzo (2021):** Sandbar shark aggregation in the central Mediterranean Sea and potential effects of tourism. *Aquat. Conserv.*, 31(6), 1420-1428.
- Celona, A., A. De Maddalena, & T. Romeo (2005):** Bluntnose sixgill shark, *Hexanchus griseus* (Bonnaterre, 1788), in the eastern north Sicilian waters. *Boll. Mus. stor. nat. Venezia*, 56, 137-151.
- Compagno, L.J.V. (1984):** FAO Species Catalogue. Vol.4. Sharks of the World. An annotated and illustrated catalogue of sharks species known to date. Parts 1 and 2. FAO Fish. Synop., 125, 1-655.
- De Maddalena, A., H. Baensch & W. Heim (2015):** Sharks Of The Mediterranean. An Illustrated Study of All Species. McFarland & Co., Jefferson, 204 pp.
- De Maddalena, A. & P. Galli (2017):** Sharks of the Maldives. Editoriale Magenes, Milano, 216 pp.
- Dulvy, N.K., Y. Sadovy & J.D. Reynolds (2003):** Extinction vulnerability in marine populations. *Fish Fish.*, 4, 25-64. DOI: 10.1046/j.1467-2979.2003.00105.x.
- Gajić, A.A., S. Lelo, A. Joksimović, A. Pešić, J. Tomanić, H. Beširović & B. Dragičević (2021):** Contemporary records of the rare and critically endangered angular rough shark, *Oxynotus centrina* (Linnaeus, 1758), from the eastern Adriatic Sea. *J. Fish Biol.*, 100(1), 329-334.
- IUCN (2022):** The IUCN Red List of Threatened Species. Version 2021-3: <https://www.iucnredlist.org>.
- Kabasakal, H. (2009):** Observations on a rare shark, *Oxynotus centrina* (Chondrichthyes: Oxynotidae), in the sea of Marmara (North-Western Turkey). *Pan-Am. J. Aquat. Sci.*, 4, 609-612.
- Kabasakal, H. (2010):** Historical and contemporary records of the angular rough shark *Oxynotus centrina* (Chondrichthyes; Oxynotidae) in Turkish waters. *Mediterr. Mar. Sci.*, 11(2), 361-367.
- Kabasakal, H. (2020):** A Field Guide to the Sharks of Turkish Waters. Turkish Marine Research Foundation (TUDAV) Publication No. 55, Istanbul, 133 pp.
- Koehler, L. (2018):** New records of angular rough sharks *Oxynotus centrina* in the coastal waters of Malta, with observations on post-capture resilience and release behaviour. *J. Fish Biol.*, 92(6), 2039-2044. DOI:10.1111/jfb.13461.
- Kousteni V. & P. Megalofonou (2016):** Observations on the biological traits of the rare shark *Oxynotus centrina* (Chondrichthyes: Oxynotidae) in the Hellenic Seas. *J. Fish Biol.*, 89(3), 1880-1888. DOI: 10.1111/jfb.13077.
- Lipej, L., A. De Maddalena & A. Soldo (2004):** Sharks of the Adriatic Sea. Knjižnica Annales Major, Koper, 254 pp.
- Mizzan, L. (1994):** I Leptocardi, Ciclostomi e Selaci delle collezioni del Museo Civico di Storia Naturale di Venezia - 1) Leptocardia, Agnatha, Gnathostomata - Chondrichthyes (esclusi Rajiformes). *Boll. Mus. stor. nat. Venezia*, 45, 123-137.

Moreno, J.A. (1995): Guía de los tiburones del Atlántico Nororiental y Mediterráneo. Ed. Pirámide, Madrid, 310 pp.

Tortonese, E. (1956): Fauna d'Italia vol. II. Leptocardia, Ciclostomata, Selachii. Calderini, Bologna, 334 pp.

Vanni, S. (1992): Cataloghi del Museo di Storia Naturale dell'Università di Firenze, Sezione di Zoologia "La Specola". XI. Chondrichthyes. Atti Soc. Toscana Sci. Nat. Mem. B, 99, 85-114.

Zemah Shamir, Z., S. Zemah Shamir, N. Becker, A. Scheinin & D. Tchernov (2019): Evidence of the impacts of emerging shark tourism in the Mediterranean. Ocean Coast. Manag., 178. DOI: 10.1016/j.ocecoaman.2019.104847.