

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 32, 2022, 1





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 32, 2022, 1

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 32, leto 2022, številka 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capapé (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Mitja Kaligarič, Marcelo Kovačič (HR), Andrej Kranjc, Lovrenc Lipej, Vesna Mačič (ME), Alenka Malej, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®

Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®

**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, **internet:** www.zdjp.si

Redakcija te številke je bila zaključena 30. 06. 2022.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za raziskovalno dejavnost Republike Slovenije (ARRS) in Mestna občina Koper

Annales - Series Historia Naturalis izhaja dvakrat letno.

Naklada/Tiratura/Circulation:

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY-NC 4.0.



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>



VSEBINA / INDICE GENERALE / CONTENTS 2022(1)

SREDOZEMSKI MORSKI PSI
SQUALI MEDITERRANEI
 MEDITERRANEAN SHARKS

Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ

Observations on Thresher Shark, *Alopias vulpinus* (Chondrichthyes: Alopiidae) from the Coast of Algeria (Southwestern Mediterranean Sea) 1
Opazovanja morskih lisic, Alopias vulpinus (Chondrichthyes: Alopiidae) ob alžirski obali (jugozahodno Sredozemsko morje)

Elif ÖZGÜR ÖZBEK & Hakan KABASAKAL

Notes on Smoothback Angel Shark, *Squatina oculata* (Squatiniformes: Squatinidae) caught in the Gulf of Antalya 9
Zapis o pegastih sklatih, Squatina oculata (Squatiniformes: Squatinidae), ujetih v Antalijskem zalivu

Alessandro PAGANO & Alessandro DE MADDALENA

Underwater Observations of the Rare Angular Roughshark *Oxynotus centrina* (Chondrichthyes: Squalidae) in the Waters of Santa Tecla (Sicily, Italy) 17
Podvodna opazovanja redkega morskega prašiča, Oxynotus centrina (Chondrichthyes: Squalidae) v vodah Sante Tecele (Sicilija, Italija)

Deniz ERGÜDEN, Deniz AYAS & Hakan KABASAKAL

Morphometric Measurements of Three Young Carcharhinid Species from Northeastern Levant (Mediterranean Sea) 25
Morfometrične meritve mladičev treh vrst morskih psov iz družine Carcharhinidae iz severnovzhodnega Levanta (Sredozemsko morje)

Hakan KABASAKAL

Projections on the Future of Deep-Sea Sharks in the Sea of Marmara, Where Deep Zones Are Threatened by Deoxygenation: a Review 35
Napovedi o prihodnosti globomorskih morskih psov v Marmarskem morju, ogroženem zaradi pomanjkanja kisika: pregled

BIOINVAZIJA
 BIOINVASIONE
 BIOINVASION

Alan DEIDUN, Bruno ZAVA & Maria CORSINI-FOKA

Distribution Extension of *Lutjanus argentimaculatus* (Lutjanidae) and *Psenes pellucidus* (Nomeidae) to the Waters of Malta, Central Mediterranean Sea 49
Širjenje areala vrst Lutjanus argentimaculatus (Lutjanidae) in Psenes pellucidus (Nomeidae) v malteške vode (osrednje Sredozemsko morje)

Sami M. IBRAHIM, Abdulrazziq A. ABDULRAZIQ, Abdulghani ABDULGHANI, Sara A.A. AL MABRUK, David SALVATORI, Bruno ZAVA, Maria CORSINI-FOKA & Alan DEIDUN

First Record of *Enchelycore anatina* (Muraenidae) from Libyan Waters and an Additional Record from Southern Italy (Western Ionian Sea) 59
Prvi zapis o pojavljanju kavljazobe murene Enchelycore anatina (Muraenidae) iz libijskih voda in dodatni zapis za južno Italijo (zahodno Jonsko morje)

Rasha Ali HENEISH & Samir Ibrahim RIZKALLA

Morphometric and Meristic Characteristics of a New Record of Bluespot Mullet *Crenimugil seheli* (Pisces: Mugilidae) in Egyptian Mediterranean waters 67
Novi zapis o pojavljanju vrste Crenimugil seheli (Pisces: Mugilidae) v egiptovskih sredozemskih vodah in njene morfolometrične in meristične značilnosti

Yana SOLIMAN, Adib SAAD, Vienna HAMMOUD & Christian CAPAPÉ

Heavy Metal Concentrations in Tissues of *Siganus rivulatus* (Siganidae) from the Syrian Coast (Eastern Mediterranean Sea) 75
Vsebnost težkih kovin v tkivih marmoriranega morskega kunca Siganus rivulatus (Siganidae) iz sirske obale (vzhodno Sredozemsko morje)

IHTIOLOGIJA
 ITTIOLOGIA
 ICHTHYOLOGY

Jihade ALAHYENE, Brahim CHIAHOU, Hammou EL HABOUZ & Abdelbasset BEN-BANI

Length Based Growth Estimation of the Blue Shark *Prionace glauca* from the Moroccan Central Atlantic Coast 85
Dolžinsko-masni odnos in ocena rasti pri sinjem morskem psu (Prionace glauca) iz osrednje atlantske obale Maroka

Okan AKYOL, Altan LÖK & Funda ERDEM

Occurrence of *Cubiceps gracilis* (Nomeidae) in the Eastern Mediterranean Sea 101
Pojavljanje klateža, Cubiceps gracilis (Nomeidae), v vzhodnem Sredozemskem morju

Farid HEMIDA, Boualem BRAHMI, Christian REYNAUD & Christian CAPAPÉ

Occurrence of the Rare Driftfish *Cubiceps gracilis* (Nomeidae) from the Algerian Coast (Southwestern Mediterranean Sea) 107
Pojavljanje redkega klazeža Cubiceps gracilis (Nomeidae) z alžirske obale (jugozahodno Sredozemsko morje)

Deniz ERGÜDEN & Cemal TURAN

A Rare Occurrence of *Carapus acus* (Carapidae) in the Eastern Mediterranean, Turkey 113
Redko pojavljanje strmorinca Carapus acus (Carapidae) v vzhodnem Sredozemskem morju (Turčija)

Laith JAWAD, Murat ŞIRIN, Miloslav PETRTÝL, Ahmet ÖKTENER, Murat ÇELİK & Audai QASIM

Skeletal Abnormalities in Four Fish Species Collected from the Sea of Marmara, Turkey 119
Skeletne anomalije pri štirih vrstah rib iz Marmarskega morja (Turčija)

RAZMNOŽEVALNA EKOLOGIJA
 ECOLOGIA RIPRODUTTIVA
 REPRODUCTIVE ECOLOGY

Amaria Latefa BOUZIANI, Khaled RAHMANI, Samira AIT DARNA, Alae Eddine BELMAHI, Sihem ABID KACHOUR & Mohamed BOUDERBALA

Gonadal Histology in *Diplodus vulgaris* from the West Algerian Coast 137
Histologija gonad pri navadnem šparu (Diplodus vulgaris) iz zahodne alžirske obale

Cheikhna Yero GANDEGA, Nassima EL OMRANI, Rezan O. RASHEED, Mohammed RAMDANI & Roger FLOWER

The Growth and Reproduction of Two Sparidae, *Pagrus caeruleostictus* and *Pagellus bellottii* in Northern Mauritanian Waters (Eastern Tropical Atlantic) 143
Rast in razmnoževanje dveh vrst pagrov, Pagrus caeruleostictus in Pagellus bellottii v severnih mavretanskih vodah (vzhodni tropski Atlantik)

Nassima EL OMRANI, Hammou EL HABOUZ, Abdellah BOUHAIMI, Jaouad ABOU OUALID, Abdellatif MOUKRIM, Jamila GOUZOU, Mohammed RAMDANI, Roger FLOWER & Abdelbasset BEN-BANI

The Reproductive Biology of the Pouting *Trisopterus luscus* from the Atlantic Coast of Morocco 155
Reproduktivna biologija francoskega moliča (Trisopterus luscus) iz atlantske obale Maroka

Mourad CHÉRIF, Rimel BENMESSAOUD & Christian CAPAPÉ

Growth Patterns and Age Structure of *Mullus surmuletus* (Mullidae) from the Northern Coast of Tunisia (Central Mediterranean Sea) 173
Rastni parametri in starostna struktura progastih bradačev Mullus surmuletus (Mullidae) iz severne tunizijske obale (osrednje Sredozemsko morje)

FLORA

FLORA

FLORA

Martina ORLANDO-BONACA, Erik LIPEJ, Romina BONACA & Leon Lojze ZAMUDA

Improvement of the Ecological Status of the *Cymodocea nodosa* Meadow near the Port of Koper 185
Izboljšanje ekološkega stanja morskega travnika kolenčaste cimodoceje (Cymodocea nodosa) v bližini kopskega pristanišča

FAVNA

FAVNA

FAVNA

Manja ROGELJA, Martin VODOPIVEC & Alenka MALEJ

Cestum veneris Lesueur, 1813 (Ctenophora) – a Rare Guest in the Northern Adriatic Sea 197
Cestum veneris Lesueur, 1813 (Ctenophora) – redek gost v severnem Jadranu

Adla KAHRIĆ, Dalila DELIĆ & Dejan KULIJER

Notospermus annulatus (Nemertea: Lineidae), a New Record for Bosnia and Herzegovina 205
Notospermus annulatus (Nemertea: Lineidae), prvi zapis o pojavljanju za Bosno in Hercegovino

Andrea LOMBARDO & Giuliana MARLETTA

Report of an Interesting *Trapania* (Gastropoda: Nudibranchia: Goniodorididae) Specimen from Central Eastern Sicily 211
Zapis o zanimivem primerku iz rodu Trapania (Gastropoda: Nudibranchia: Goniodorididae) iz osrednje vzhodne Sicilije

Abdelkarim DERBALI & Othman JARBOUI

Stock Assessment, Cartography and Sexuality of the Wedge Clam *Donax trunculus* in the Gulf of Gabes (Tunisia) 217
Ocena staleža, kartografija in spolnost klinaste školjke Donax trunculus v gabeškem zalivu (Tunizija)

Abdelkarim DERBALI, Aymen HADJ TAIEB & Othman JARBOUI

Length-Weight Relationships and Density of Bivalve Species in the Shellfish Production Area of Zarzis (Tunisia, Central Mediterranean Sea) 229
Dolžinsko-masni odnos in gostota školjk na gojišču školjk v predelu Zarsisa (Tunizija, osrednje Sredozemsko morje)

Toni KOREN

The Diversity of Moths (Lepidoptera: Heterocera) of Significant Landscape Donji Kamenjak and Medulin Archipelago, Istria, Croatia 237
Raznolikost nočnih metuljev (Lepidoptera: Heterocera) Pomembne pokrajine Donji Kamenjak in Medulinski arhipelag, Istra, Hrvaška

OCENE IN POROČILA

RECENSIONI E RELAZIONI

REVIEWS AND REPORTS

Ines Mandić Mulec & Nives Ogrinc

Recenzija knjige: Mikrobn biogeokemija vod 263

Kazalo k slikam na ovitku 265

Index to images on the cover 265

received: 2022-04-11

DOI 10.19233/ASHN.2022.20

CESTUM VENERIS LESUEUR, 1813 (CTENOPHORA) – A RARE GUEST IN THE NORTHERN ADRIATIC SEA

Manja ROGELJA

University of Primorska, Aquarium Piran, Kidričevo nabrežje 4, 6330 Piran, Slovenia
e-mail: manja.rogelja@upr.si

Martin VODOPIVEC & Alenka MALEJ

Marine Biology Station Piran, National Institute of Biology, SI-6330 Piran, Slovenia
e-mail: alenka.malej@nib.si

ABSTRACT

We report the occurrence of the ctenophore *Cestum veneris* Lesueur, 1813 in the Gulf of Trieste (northern Adriatic Sea). Three individuals were found in March 2022, representing the first documented record in the scientific literature in more than 130 years. They were observed and photographed in the sea, and two specimens were brought to the Piran Aquarium, where their behaviour was observed. Historical and recent literature on the occurrence of this species in the Adriatic Sea was reviewed. The occurrence in the Gulf of Trieste is likely related to the transport of *C. veneris* with the currents, and therefore we present simulation results for the period January-February. Despite the relatively rare detections near the coast, this species is probably widespread in the offshore waters of the central and southern Adriatic.

Key words: winter gelatinous plankton, Ctenophora, distribution, Gulf of Trieste

CESTUM VENERIS LESUEUR, 1813 (CTENOPHORA) – OSPITE RARO NELL'ADRIATICO SETTENTRIONALE

SINTESI

Gli autori segnalano la presenza dello ctenoforo *Cestum veneris* Lesueur, 1813 nel Golfo di Trieste (mare Adriatico settentrionale). A marzo 2022 sono stati avvistati tre esemplari, che rappresentano il primo ritrovamento documentato dopo oltre 130 anni. Osservati e fotografati prima in mare, due esemplari sono stati portati all'Aquario di Pirano, dove è stato seguito il loro comportamento. È stata esaminata la letteratura storica e recente sulla presenza di questa specie nel mare Adriatico. Gli avvistamenti nel Golfo di Trieste sono probabilmente legati al trasporto di *C. veneris* con le correnti, e nell'articolo vengono presentati i risultati della simulazione per il periodo gennaio - febbraio. Nonostante i rilevamenti relativamente rari in prossimità della costa, questa specie è probabilmente diffusa nel mare aperto dell'Adriatico centro-meridionale.

Parole chiave: plancton gelatinoso invernale, Ctenophora, distribuzione, Golfo di Trieste

INTRODUCTION

Plankton in the northern Adriatic, including the Gulf of Trieste, is affected in winter and early spring not only by the well-mixed and cold water, but also by the greater influx of water from the south (Zavatarelli & Pinardi, 2003), which brings some otherwise rare species to this northernmost part of the Adriatic (Fonda Umani et al., 1992). These dynamics also affect the structure of the gelatinous plankton community. According to Miloš & Malej (2005), the cold-water group of gelatinous mesozooplankton was more diverse and less abundant, while some organisms originating from the deep southern waters of the Adriatic, such as the siphonophore *Vogtia pentacantha*, were occasionally observed. The most common winter-spring macrojellyfish in the northern Adriatic included the syphozoans *Aurelia solida*, *Rhizostoma pulmo*, the hydrozoan *Aequorea forskalea*, and the ctenophore *Leucothea multicornis*, while *Cestum veneris* was observed in the central and southern Adriatic and only in a few years in the northern Adriatic (Pestorić et al., 2021).

C. veneris is a tentaculate ctenophore species that lives circumglobally in temperate, subtropical and tropical seas (Mayer, 1912; GBIF, 2019). It is considered epipelagic (Harbison et al., 1978), but has also been observed in the mesopelagic zone (Lindsay et al., 2015; Hidaka et al., 2021).

Péron and Lesueur, who found this gelatinous species in coastal waters near Nice, France, on 12 May, 1809, named it after the magical girdle of Aphrodite - Ceste de Venus - and assigned this beautiful new ctenophore *C. veneris* with a distinctly elongated and transversely compressed body to the beroids (Lesueur, 1813). However, they had only one specimen available for examination. Lesueur reported in the same note (1813) that this organism, unknown to him and his collaborator Péron, was occasionally observed in large numbers in the harbour of Villefranche, where it

was called by fishermen 'sabres de mer' (a sea saber, a kind of curved sword used especially by pirates). Later Gegenbaur (1856) created the order Cestida and the family Cestidae, which today includes two monotypic genera: *Cestum veneris* Lesueur 1813 and *Velamen parallelum* Krumbach 1925.

In our contribution, we reviewed the historical and current literature on the occurrence of *C. veneris* in the Adriatic Sea. The recent discovery of this organism in the Gulf of Trieste, the first documented after 138 years (Graeffe, 1884), prompted us to describe its distribution and discuss some ecological characteristics of this beautiful ctenophore. We also present modelling results of the winter circulation, which likely support the entry of planktonic organisms of more southern origin into the northern Adriatic and the Gulf of Trieste.

MATERIAL AND METHODS

In total, three specimens of *C. veneris* were observed, two on 2 March and one on 3 March 2022 near the coast of Piran, Gulf of Trieste. The first observed specimen was not visible from the surface, but was accidentally felt by hand while taking underwater photos. During the capture, a part of the animal broke off, so the ctenophore was immediately released after inspection. At the same time, another specimen was observed 20 m away and collected undamaged. The next day, a third specimen was found at the same location and was also collected. The sea temperature was 8°C, salinity 38.6 and the sea surface near the shore was calm with almost no waves.

Two captured specimens were brought to the Piran Aquarium, but the specimen collected on March 2 began to disintegrate very quickly. The specimen caught on March 3 was immediately transferred to the 325 litres kreisel tank with a water temperature of 13°C and salinity of 38. It was fed freshly hatched *Artemia* nauplii. Photos and videos were taken, but the quality




Fig. 1: Photo of *Cestum veneris* in the sea (left) and in the aquarium (right).
Sl. 1: Fotografija *Cestum veneris* v morju (levo) in v akvariju (desno).




Fig. 2: Characteristics of the central part of *Cestum veneris*.

Sl. 2: Značilnosti osrednjega dela *Cestum veneris*.

was not very good due to the great transparency and rapid movements of the *C. veneris*.

We reviewed the available published sources on ctenophores in the Adriatic Sea in the last 150 years since the first description of *C. veneris*. With regard to the ecology of this species, we also consulted other published sources. The global distribution of *C. veneris* shown in Fig. 2 is from the Global Biodiversity Information Facility database (GBIF, 2019).

The CROCO ocean model (Coastal and Regional Ocean COMMunity model, formerly known as ROMS_AGRIF; www.croco-ocean.org) was used to simulate the Adriatic circulation during the 2000-2018 period. The model configuration was similar to the one used in Vodopivec *et al.* (2017), but with horizontal resolution reduced to 4 km and new atmospheric forcing provided by ERA5 (Hersbach *et al.*, 2020). Barotropic currents and free surface were averaged over the entire simulation period for January and February to provide an overview of the general circulation during the months when the *Cestum veneris* was likely transported from the central to the northern Adriatic.

RESULTS AND DISCUSSION

In the current classification, *C. veneris* is listed as a single species of the genus, belonging to the family Cestidae along with another monotypic genus *Velamen* Krumbach 1925 (Mills, 1998 - present):

Phylum Ctenophora
Class Tentaculata
Order Cestida Gegenbaur 1856
Family Cestidae Gegenbaur 1856
Genus *Cestum* Gegenbaur 1856
Species *Cestum veneris* Lesueur 1813

C. veneris is characterised by a transparent, ribbon-shaped body, flattened in the tentacular axis while the lobular axis is elongated (Fig. 1). Adults can reach a height of about 8 cm and a length up to nearly 1,5 m but usually less than 80 cm. In contrast to those in g. *Velamen*, where subtentacular meridional canals arise directly without bending, in *Cestum* they arise from the stomodaeum then bend outward and run along the midline of the body. Substomodeal comb rows are greatly elongated, while four subtentacular ones are rudimentary on aboral edge close to the sense organ (Fig. 2). On both sides of the mouth, grooves with tentilla extend along the entire oral margin of the body. Gonads run in continuous line under substomodaeal meridional canals.

All Ctenophora are carnivorous and their feeding mode is related to their morphology: they forage with tentacles, lobes, or feed by engulfing the prey (Haddock, 2007). *C. veneris* search for prey hovering or gliding horizontally with the oral edge leading (Matsumoto & Harbison, 1993) and tentilla stream aborally like curtains. After a few meters, *Cestum*




Fig. 3: Global distribution of *Cestum veneris* (<https://www.gbif.org/species/2501198>).

Sl. 3: Globalna razširjenost rebrače *Cestum veneris* (<https://www.gbif.org/species/2501198>).

move vertically and reverse swimming direction. Tentillae with sticky colloblasts attach to prey such as small copepods and then contract. Prey items are transferred to the oral groove and moved toward the mouth using cilia (Matsumoto & Harbison, 1993). Besides comb-row hovering or gliding, *Cestum* can also swim by muscular body undulations (Ceccaldi, 1972).

***Cestum veneris* distribution**

Cestum veneris is probably one of the most fragile ctenophores. Therefore, it is not surprising that we cannot catch them with conventional nets, which have been most commonly used in plankton surveys in the past. This characteristic, therefore, has a major impact on assessing *C. veneris* distribution and abundance. Nevertheless, researchers in the past have described the distribution of this species based on observations either from the coast or during research cruises. Their distribution was studied in particular during the ‘first golden age of gelata research’, from the late 19th to the early 20th century (Haddock, 2004), when many new species of Ctenophora were described, (Chun, 1878; Bigelow, 1904; Mayer, 1912). The development of autonomous diving, remote sensing, and new *in situ* observation techniques improved the collection of data on delicate gelatinous organisms. In addition, citizen science has enabled the collection of large amounts of data over broad areas and long periods of time. The availability of photographs and video clips facilitated the recognition of observed organisms by experts. Today, platforms such as Mer et littoral - European Marine Life or Global Biodiversity Information Facility provide information on the distribution of many marine species, including *C. veneris* (Fig. 3).

There are relatively few published data on ctenophores in the Adriatic Sea (Pestorić *et al.*, 2021). The rare records of ctenophores are usually not the result of systematic studies of these gelatinous organisms but are rather based on incidental observations such as ours. Some evidence come from published regular plankton observations for those species that are better preserved and later from the establishment of citizen science programs. The non-native *Mnemiopsis leidyi* is more systematically studied and is therefore a rare exception due to its significant impact on the ecosystem (Malej *et al.*, 2017; Budiša *et al.*, 2021).

The first record of *C. veneris* from the Adriatic Sea dates back to the 19th century when Graeffe (1884) reported rare, occasional observations of single specimens in the Gulf of Trieste. Krumbach (1911) found four specimens near Rovinj in mid-December 1910; two were about 10 cm and two about 25 cm long. In the southern Adriatic, Babić (1913) reported one small specimen observed near the island of Mljet in March 1912 and three larger specimens (about 50 cm) near Komiža (island of Vis). These few observations suggest that *C. veneris* was rare in the Adriatic, but Benović & Lučić (2001) described massive strandings and swarms near the island of Korčula and Dubrovnik after strong southerly winds. For more information on temporal and spatial distribution of *C. veneris* in the Adriatic, see Pestorić *et al.* (2021) and Viočić *et al.* (2022). It was described as common in the central and southern Adriatic in winter and formed blooms in some years (1999, 2013). In the northern Adriatic, only few individuals were recorded in 2015–2017 period, while none were observed in the Gulf of Trieste (Pestorić *et al.*, 2021). Also, in the checklist of Ctenophora in the Italian seas (Mills, 2008), *C. veneris* was listed only for the central

and southern Adriatic. In the Mediterranean (Gulf of Naples), blooms of *C. veneris* were reported by Chun (1878) between December and May. Harbison et al. (1978), presenting the distribution and natural history of oceanic ctenophores of the North Atlantic, reported that *C. veneris* was the most abundant ctenophore species, and its density exceeded 1 per m³ at some stations.

Being an oceanic species, it is not surprising that it is so rare in the shallow northern Adriatic and Gulf of Trieste. Our observation of *C. veneris* in March 2022 is the first from the Gulf of Trieste (Fig. 2) since the report of Graeffe (1884). All three observed specimens were in shallow water, had a rectilinear posture, and swam just below the sea surface. They were swimming by bending, which resembled a wave moving from one side of their body to the other. The undulation started on the side of their body where the movement was directed and ran over to the other side like a wave. They could swim in either direction by changing sides at the beginning of the bend. Similar behaviour of individuals up to 80 cm in length was observed in diving by Ceccaldi (1972), who also noticed curling (rolling upon itself) in captivity and attributed this behaviour to stress.

Our observations in the aquarium agree with those of Ceccaldi (1972). Immediately after release into the kreisel tank, the *Cestum* curled up and remained in this shape for some time. Then it slowly stretched and moved in the same manner as it had previously been observed in the sea. In the following hours it was mostly fully stretched and moved with the circular flow of the water. A continuous movement of the cilia was observed. *Cestum* also ate newly hatched *Artemia* nauplii offered as food, as individual nauplii were observed in the stomach in the following hours (Fig. 2). During this time, it moved effortlessly and remained undamaged. The length was roughly estimated at 40 to 50 cm, measured by the size of the container in which they were held.

The rare sporadic records in the northern Adriatic and the Gulf of Trieste (Graeffe, 1884; Krumbach, 1911; Pestorić et al., 2021; this article) during winter are likely due to the transport of *C. veneris* with the currents. This assumption is supported by the modeled barotropic currents for January and February, shown in Figure 4. The northward current along the eastern Adriatic coast is clearly visible. It passes the Kvarner Bay (KV) and splits into two branches at the southern tip of the Istrian peninsula (IS). The stronger branch veers westward and crosses the basin, while the weaker branch travels further north towards the Gulf of Trieste (GT).

The presence of *C. veneris* in the Gulf of Trieste in March 2022 could be caused by a bora (strong NE wind) event lasting several days, which occurred




Fig. 4: Average January and February barotropic currents and free surface for the 2000-2018 period (model results). The arrow top-right represents current speed of 0.1 m/s, the colour scale is in meters. Arrows are plotted for every fourth grid cell. GT – Gulf of Trieste, IS – Istrian peninsula, KV – Kvarner Bay.

Sl. 4: Povprečni barotropni tokovi in prosta gladina v januarju in februarju za obdobje 2000-2018 (modelski rezultati). Puščica v desnem zgornjem kotu slike predstavlja tok s hitrostjo 0.1 m/s. Barvna skala je v metrih. Puščice so narisane za vsako četrto točko modelske mreže. GT – Tržaški zaliv, IS – Istra, KV – Kvarner.

shortly before its discovery. A study of topographic control of wind-driven circulation has shown that the bora induces a compensating current on the south-eastern side of the Gulf of Trieste (Malačič et al., 2012 - Figure 7). The leeward current (flowing into the gulf) is present almost throughout the entire water column, except for a few meters below the surface, where the current is windward. The inflowing north-eastward current could be responsible for the transport of the *C. veneris* individuals into the Gulf of Trieste.

In summary, despite the relatively rare finds near the Adriatic coasts, especially in the shallow northernmost area, this species is probably widespread in the offshore waters of the central and southern Adriatic.

ACKNOWLEDGMENTS

The authors acknowledge the financial support from the Slovenian Research Agency (research core funding No. P1-0237 “Coastal Sea Research” and project Z7-1884).

CESTUM VENERIS LESUEUR, 1813 (CTENOPHORA) – REDEK GOST V SEVERNEM JADRANU

Manja ROGELJA

Univerza na Primorskem, Akvarij Piran, Kidričevo nabrežje 4, 6330 Piran, Slovenia
e-mail: manja.rogelja@upr.si

Martin VODOPIVEC & Alenka MALEJ

Morska biološka postaja Piran, Nacionalni inštitut za biologijo, Fornace 41, SI-6330 Piran, Slovenia
e-mail: alenka.malej@nib.si

POVZETEK

Avtorji poročajo o redkem pojavu rebrače venerin pas (*Cestum veneris* Leseuer, 1813) v Tržaškem zalivu (severni Jadran). Trije osebki venerinega pasu so bili opaženi marca 2022, kar je prva v znanstveni literaturi dokumentirana najdba po več kot 130 letih. Osebke smo opazovali in fotografirali v morju, dva pa prenesli tudi v Akvarij Piran, kjer smo opazovali obnašanje. Pregledali smo historično in novejšo literaturo o pojavljanju te vrste rebrač v Jadranskem morju. Pojav venerinega pasu v Tržaškem zalivu povezujemo z vnosom oz. transportom vodnih mas iz južnejših delov. Na osnovi nekaterih objav za Jadransko in Sredozemsko morje ter opazovanj v severnem Atlantiku sklepamo, da je ta vrsta, kljub ne ravno pogostim opažanjem v priobalnih vodah, dokaj običajna v odprtih vodah srednjega in južnega Jadrana.

Ključne besede: zimski želatinozni plankton, Ctenophora, razširjenost, Tržaški zaliv

REFERENCES

- Babić, K. (1913):** Planktonički celenterati iz Jadranskog mora. Rad JAZU Zagreb, 55, 186-202.
- Benović, A. & D. Lučić (2001):** Jellyfish outbreaks: natural cycles or stress response effects. CIESM Workshop Series, 14, 59-62.
- Bigelow (1904):** Medusae from the Maldive Islands. Bull. Museum Comp. Zool., 39, 245-269
- Budiša, A., P. Paliaga, T. Juretić, D. Lučić, N. Supić, Z. Pasarić, T. Djakovac, M. Mladinić, V. Dadić & V. Tičina (2021):** Distribution, diet and relationships of the invasive ctenophore *Mnemiopsis leidyi* with anchovies and zooplankton, in the northeastern Adriatic Sea. Mediterranean Marine Science, 22(4), 827-842.
- Ceccaldi H.J. (1972):** Observations biologiques de *Cestus veneris*. Tethys, 4, 707-710.
- Chun, C. (1878):** Die im Golf von Neapel erscheinenden Rippenquallen. Mittheilungen. a.d. Zoologischen Station zu Neapel, 1, 180-217.
- Fonda Umani, S., P. Franco, E. Ghirardelli & A. Malej (1992):** Outline of oceanography and the plankton of the Adriatic Sea. In: Marine Eutrophication and Population Dynamics (Eds. Colombo et al), Olsen & Olsen, Fredesborg, 347-365.
- GBIF Secretariat (2019):** GBIF Backbone Taxonomy. European-marine-life.org/06/cestum-veneris.php, Downloaded on 29 March 2022.
- Gamulin, T. (1979):** Zooplankton of the eastern Adriatic Sea. Prir. Istr. JAZU, 43, 177-270 (in Croatian).
- Gegenbaur, C. (1856):** Studien über Organisation und Systematik der Ctenophores. Archiv für Naturgeschichte, 2(1), 163-205.
- Graeffe, E. (1884):** Uebersicht der Seethierfauna des Golfes von Triest. III. Coelenteraten. Arbeit. Zool. Inst. Wien, Bd. 5, 333-362.
- Gueroun, S.K.M, S. Schäfer, F. Gizzi, S. Álvarez, J.G. Monteiro, C. Andrade & J. Canning-Clode (2021):** Planktonic Ctenophora of the Madeira Archipelago (Northeastern Atlantic). Zootaxa, 5081(3), 433-443.
- Haddock, S.H.D. (2004):** A golden age of gelata: past and future research on planktonic ctenophores and cnidarians. Hydrobiologia, 530/531, 549-556.
- Haddock, S.H.D. (2007):** Comparative feeding behaviour of planktonic ctenophores. Integrative and Comparative Biology, 47(6), 847-853. <https://doi.org/10.1093/icb/icm088>.
- Harbison, G.R., L.P. Madin & N.R. Swanberg (1978):** On the natural history and distribution of oceanic ctenophores. Deep-Sea Research, 25(3), 233-256.
- Hersbach, H., B. Bell, P. Berrisford, S. Hirahara, A. Horányi, J. Muñoz-Sabater et al. (2020):** The ERA5 global reanalysis. Quarterly Journal of the Royal Meteorological Society, 146(730), 1999-2049.
- Hidaka, M., J. Nishikawa & D.J. Lindsay (2021):** Gelatinous zooplankton community around a hydrothermally active deep-sea caldera: Results from ROV video records. Plankton and Benthos Research, 16(1), 40-58. <https://doi.org/10.3800/pbr.16.40>.
- Krumbach, T. (1911):** Notizen über die Fauna der Adria bei Rovigno. IV. Die Ctenophorenfauna von Rovigno nach den Novemberstürmen 1910. Zool. Anz., 37, 315-319.
- Lesueur, C.A. (1813):** Mémoire sur quelques nouvelles espèces d'animaux mollusques et radiaires recueillis dans la Méditerranée près de Nice. Nouveau Bulletin des Sciences, par la Société Philomatique de Paris, 3(69), 281-285.
- Lindsay, D.J., M. Umetsu, M. Grossmann, H. Miyake & H. Yamamoto (2015):** The gelatinous macroplankton community at the Hatoma Knoll hydrothermal vent. In: Ishibashi, J., Okino, K. & Sunamura, M. (Eds.), Subseafloor Biosphere Linked to Hydrothermal Systems: TAIGA Concept. Springer, Tokyo, pp. 639-666.
- Malačič, V., B. Petelin & M. Vodopivec (2012):** Topographic control of wind-driven circulation in the northern Adriatic. Journal of Geophysical Research: Oceans, 117(C6).
- Malej, A., V. Tirelli, P. Paliaga, M. Vodopivec, A. Goruppi, S. Ancona, M. Benzi, N. Bettoso, E. Camatti, M. Ercolessi, C.R. Ferrari & T. Shiganova (2017):** *Mnemiopsis leidyi* in the northern Adriatic: here to stay? J. Sea Res., 124, 10-16.
- Matsumoto, G.I. & G.R. Harbison (1993):** *In situ* observations of foraging, feeding, and escape behavior in three orders of oceanic ctenophores: Lobata, Cestida, and Beroida. Marine Biology, 117, 279-287.
- Mayer, A.G. (1912):** Ctenophores of the Atlantic coast of North America. Carnegie Institution, Washington, 58 pp. <https://doi.org/10.5962/bhl.title.5968>.
- Mer et Littoral - European Marine Life: <https://www.european-marine-life.org/06/cestum-veneris.php>**, Downloaded, 31 March 2022
- Mills, C.E. (1998-present):** Phylum Ctenophora: list of all valid species names. Electronic internet document. *Cestum veneris* Lesueur, 1813. Accessed through: World Register of Marine Species at: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=106363> on 2022-04-03.
- Mills, C.E. (2008):** Ctenophora. Checklist of Fauna Italiana. Biol. Mar. Mediterr., 15 (suppl.), 102-104.
- Miloš, Č. & A. Malej (2005):** Gelatinous plankton assemblages in temperate coastal waters – seasonal variations (Gulf of Trieste, Adriatic Sea). Annales, Ser. Hist. Nat., 15(1), 11-20.

Pestorić, B., D. Lučić, N. Bojanić, M. Vodopivec, T. Kogovšek, I. Violić, P. Paliaga & A. Malej (2021): Scyphomedusae and Ctenophora of the Eastern Adriatic: Historical Overview and New Data. *Diversity*, 13, 186.

Purcell, J.E. (2012): Jellyfish and Ctenophore Blooms Coincide with Human Proliferations and Environmental Perturbations. *Annu. Rev. Mar. Sci.*, 4, 209-235.

Rossi, L. (1971): Guida a cnidari e ctenofori della fauna italiana. *Quad. Civ. Staz. Idrob. Milano*, 2, 91-95.

Violić, I., D. Lučić, N. Bojanić, B. Pestorić, B. Gangai Zovko, I. Onofri & M. Hure (2022): Long-term Monitoring of Carnivorous Macrozooplankton Dubrovnik-Neretva County. *Naše more*, 69(1), 22-29.

Vodopivec, M., Á.J. Peliz & A. Malej (2017): Offshore marine constructions as propagators of moon jellyfish dispersal. *Environmental Research Letters*, 12(8), 084003.

Zavatarelli, M. & N. Pinardi (2003): The Adriatic Sea modelling system: a nested approach. *Annales Geophysicae*, 21(1), 345-364.